

Full wwPDB X-ray Structure Validation Report (i)

Nov 24, 2024 – 06:07 pm GMT

PDB ID	:	9F7L
Title	:	Bartonella henselae NrnC bound to deoxy-pGG
Authors	:	Mortensen, S.; Sondermann, H.
Deposited on	:	2024-05-04
Resolution	:	2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motria	Whole archive	Similar resolution
	$(\# {\rm Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R _{free}	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	207	2% 91%	7% •
1	С	207	.% 91%	8%
1	Е	207	94%	5%•
1	G	207	93%	6%
1	Ι	207	3% 	•



Mol	Chain	Length	Quality of chain	
1	К	207	89%	10%
1	М	207	% 95%	5%
1	0	207	88%	12%
2	В	2	50% 50%	
2	D	2	50% 50%	
2	F	2	100%	
2	Н	2	50% 50%	
2	J	2	50% 50%	
2	L	2	50% 50%	
2	Ν	2	50% 50%	
2	Р	2	50% 50%	

Continued from previous page...



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 14342 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Δ	202	Total	С	Ν	0	S	0	0	0
	A	203	1614	1017	288	303	6	0	0	0
1	С	206	Total	С	Ν	0	S	0	0	0
1		200	1641	1032	293	309	7	0	0	0
1	F	205	Total	С	Ν	0	S	0	0	0
1		205	1633	1027	292	308	6	0	0	0
1	С	206	Total	С	Ν	0	S	0	0	0
1	G	200	1638	1030	293	309	6	0	0	0
1	т	206	Total	С	Ν	0	S	0	0	0
1	1	200	1628	1024	289	309	6	0	0	
1	K	206	Total	С	Ν	0	S	0	0	0
	Γ	200	1638	1030	293	309	6	0	0	0
1	М	206	Total	С	Ν	0	S	0	0	0
	111	200	1641	1032	293	309	7	0	0	0
1	0	207	Total	С	Ν	0	S	0	0	0
		207	1647	1035	294	311	7		0	

• Molecule 1 is a protein called 3'-5' exonuclease.

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	0	SER	-	expression tag	UNP X5MEI1
С	0	SER	-	expression tag	UNP X5MEI1
Е	0	SER	-	expression tag	UNP X5MEI1
G	0	SER	-	expression tag	UNP X5MEI1
Ι	0	SER	-	expression tag	UNP X5MEI1
K	0	SER	-	expression tag	UNP X5MEI1
М	0	SER	-	expression tag	UNP X5MEI1
0	0	SER	-	expression tag	UNP X5MEI1

• Molecule 2 is a DNA chain called DNA (5'-R(P*DGP*DG)-3').



Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
0	D	0	Total	С	Ν	0	Р	0	0	0
	D	Δ	45	20	10	13	2	0	0	0
9	П	9	Total	С	Ν	Ο	Р	0	0	0
	D	2	45	20	10	13	2	0	0	0
9	F	9	Total	С	Ν	Ο	Р	0	0	0
	Г	2	45	20	10	13	2	0	0	0
9	ц	9	Total	С	Ν	Ο	Р	0	0	0
	11	2	45	20	10	13	2	0		0
9	т	9	Total	С	Ν	Ο	Р	0	0	0
	J	2	45	20	10	13	2	0	0	0
0	т	2	Total	С	Ν	Ο	Р	0	0	0
		2	45	20	10	13	2	0	0	U
0	N	0	Total	С	Ν	Ο	Р	0	0	0
	IN	Δ	45	20	10	13	2	0	0	0
0	D	2	Total	С	Ν	Ο	Р	0	0	0
	L L		45	20	10	13	2			U

• Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	3	Total Na 3 3	0	0
3	С	2	Total Na 2 2	0	0
3	Ε	2	Total Na 2 2	0	0
3	G	2	Total Na 2 2	0	0
3	Ι	2	Total Na 2 2	0	0
3	K	2	Total Na 2 2	0	0
3	М	2	Total Na 2 2	0	0
3	0	2	Total Na 2 2	0	0

• Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	Total Cl 1 1	0	0
4	С	1	Total Cl 1 1	0	0



• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	141	Total O 141 141	0	0
5	В	7	Total O 7 7	0	0
5	С	127	Total O 127 127	0	0
5	D	4	Total O 4 4	0	0
5	Е	132	Total O 132 132	0	0
5	F	5	Total O 5 5	0	0
5	G	103	Total O 103 103	0	0
5	Н	7	Total O 7 7	0	0
5	Ι	125	Total O 125 125	0	0
5	J	7	Total O 7 7	0	0
5	K	80	Total O 80 80	0	0
5	L	1	Total O 1 1	0	0
5	М	76	Total O 76 76	0	0
5	Ν	6	Total O 6 6	0	0
5	О	57	Total O 57 57	0	0
5	Р	5	Total O 5 5	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



 \bullet Molecule 1: 3'-5' exonuclease



Chain K:	89%	10%
SER M1 E3 E3 E3 E3 A2 A2 C42 L43	T49 K60 H79 C1117 C1117 L118 N128 S131 K122 S131 W139 W139 W139 W139 W139 W139 W139 W	E150 7151 164 164 164 164 164 188 M188 M188 M188 206
• Molecule 1:	3'-5' exonuclease	
Chain M:	95%	5%
SER N1 12 119 020 042 042	H79 H79 E173 M188 M188 M188 S206	
• Molecule 1:	3'-5' exonuclease	
Chain O:	88%	12%
• Molecule 2:	DNA (5'-R(P*DGP*DG)-3')	K132 K132 Q135 Q135 S136 S145 S145 S145 P151 P151 P151 P151 P151 P181 S206 S206
Chain B:	50%	50%
<mark>6604</mark>		
• Molecule 2:	DNA (5'-R(P*DGP*DG)-3')	
Chain D:	50%	50%
G6 03 G6 04		
• Molecule 2:	DNA $(5'-R(P*DGP*DG)-3')$	
Chain F:	100%	
G603 G604		
• Molecule 2:	DNA (5'-R(P*DGP*DG)-3')	
Chain H:	50%	50%
6603 6604		
• Molecule 2:	DNA (5'-R(P*DGP*DG)-3')	
Chain J:	50%	50%
	W O PROT	

G603 G604

• Molecule 2: DNA $(5'-R(P*DGP*DG)-3')$						
Chain L:	50%	50%				
G603 G604						
• Molecule	e 2: DNA (5'-R(P*DGP*DG)-3')					
Chain N:	50%	50%				
G603 G604						
• Molecule 2: DNA $(5'-R(P*DGP*DG)-3')$						
Chain P:	50%	50%				
G604						



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	71.33Å 128.05Å 128.71Å	Depositor
a, b, c, α , β , γ	90.00° 94.82° 90.00°	Depositor
Besolution(Å)	47.57 - 2.20	Depositor
Resolution (A)	47.57 - 2.20	EDS
% Data completeness	96.9(47.57-2.20)	Depositor
(in resolution range)	87.5(47.57-2.20)	EDS
R _{merge}	0.06	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$0.86 (at 2.20 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.19.2_4158, PHENIX 1.19.2_4158	Depositor
D D	0.176 , 0.210	Depositor
$\mathbf{n}, \mathbf{n}_{free}$	0.176 , 0.210	DCC
R_{free} test set	113344 reflections $(1.75%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	36.6	Xtriage
Anisotropy	0.190	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34, 39.6	EDS
L-test for twinning ²	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14342	wwPDB-VP
Average B, all atoms $(Å^2)$	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bo	nd lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.42	0/1641	0.62	0/2216	
1	С	0.40	0/1669	0.61	0/2254	
1	Е	0.40	0/1661	0.61	0/2244	
1	G	0.39	0/1666	0.60	0/2251	
1	Ι	0.40	0/1656	0.61	0/2240	
1	Κ	0.37	0/1666	0.59	0/2251	
1	М	0.38	0/1669	0.60	0/2254	
1	0	0.35	0/1675	0.59	0/2262	
2	В	1.72	1/50~(2.0%)	0.67	0/74	
2	D	1.83	1/50~(2.0%)	0.68	0/74	
2	F	1.74	1/50~(2.0%)	0.70	0/74	
2	Н	1.83	1/50~(2.0%)	0.77	0/74	
2	J	1.80	1/50~(2.0%)	0.75	0/74	
2	L	1.81	1/50~(2.0%)	0.79	0/74	
2	N	1.75	1/50~(2.0%)	0.74	0/74	
2	Р	1.74	1/50~(2.0%)	0.72	0/74	
All	All	0.49	8/13703~(0.1%)	0.61	0/18564	

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms		Observed(Å)	Ideal(Å)
2	D	603	DG	OP3-P	-11.36	1.47	1.61
2	L	603	DG	OP3-P	-11.01	1.48	1.61
2	Н	603	DG	OP3-P	-10.98	1.48	1.61
2	J	603	DG	OP3-P	-10.95	1.48	1.61
2	N	603	DG	OP3-P	-10.88	1.48	1.61
2	F	603	DG	OP3-P	-10.79	1.48	1.61
2	В	603	DG	OP3-P	-10.68	1.48	1.61
2	Р	603	DG	OP3-P	-10.40	1.48	1.61

There are no bond angle outliers.



There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1614	0	1614	9	0
1	С	1641	0	1643	12	0
1	Е	1633	0	1631	6	0
1	G	1638	0	1636	7	0
1	Ι	1628	0	1614	2	0
1	K	1638	0	1636	15	0
1	М	1641	0	1643	5	0
1	0	1647	0	1648	10	0
2	В	45	0	23	0	0
2	D	45	0	23	0	0
2	F	45	0	23	1	0
2	Н	45	0	23	2	0
2	J	45	0	23	0	0
2	L	45	0	23	1	0
2	Ν	45	0	23	0	0
2	Р	45	0	23	0	0
3	А	3	0	0	0	0
3	С	2	0	0	0	0
3	Ε	2	0	0	0	0
3	G	2	0	0	0	0
3	Ι	2	0	0	0	0
3	Κ	2	0	0	0	0
3	М	2	0	0	0	0
3	0	2	0	0	0	0
4	А	1	0	0	0	0
4	С	1	0	0	0	0
5	А	141	0	0	2	0
5	В	7	0	0	0	0
5	С	127	0	0	5	0
5	D	4	0	0	0	0
5	Е	132	0	0	4	0
5	F	5	0	0	0	0
5	G	103	0	0	2	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	Н	7	0	0	2	0
5	Ι	125	0	0	0	0
5	J	7	0	0	1	0
5	Κ	80	0	0	2	0
5	L	1	0	0	0	0
5	М	76	0	0	0	0
5	Ν	6	0	0	0	0
5	0	57	0	0	0	0
5	Р	5	0	0	0	0
All	All	14342	0	13249	59	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:K:132:LYS:NZ	5:K:402:HOH:O	2.24	0.71
1:E:25:ASP:OD1	5:E:401:HOH:O	2.10	0.70
1:C:191:ASN:ND2	5:C:402:HOH:O	2.27	0.68
1:C:178:VAL:HG22	1:K:188:MET:HE1	1.78	0.66
1:A:167:GLU:OE2	5:A:401:HOH:O	2.14	0.65
1:A:181:ALA:HB3	1:G:188:MET:HG2	1.79	0.65
1:K:3:GLU:HB3	1:K:49:THR:HG22	1.78	0.65
1:G:171:LYS:NZ	5:G:402:HOH:O	2.30	0.64
1:O:115:ARG:NH2	1:O:120:GLU:OE2	2.31	0.64
2:H:604:DG:OP1	5:H:701:HOH:O	2.15	0.63
1:E:167:GLU:OE2	5:E:402:HOH:O	2.16	0.62
1:A:205:HIS:NE2	2:H:603:DG:H5"	2.15	0.61
1:E:188:MET:HG2	1:M:181:ALA:HB3	1.84	0.60
5:E:401:HOH:O	2:F:604:DG:OP1	2.17	0.59
1:0:146:ARG:0	1:O:150:GLU:HG3	2.04	0.58
1:A:60:LYS:NZ	5:A:405:HOH:O	2.36	0.58
1:O:27:GLU:HG2	1:O:42:GLN:HE22	1.69	0.57
1:K:145:SER:O	1:K:149:ILE:HD12	2.05	0.57
1:G:191:ASN:ND2	5:G:403:HOH:O	2.34	0.55
1:I:188:MET:HG2	1:O:181:ALA:HB3	1.89	0.54
1:G:25:ASP:OD1	5:H:701:HOH:O	2.17	0.53
1:A:79:HIS:CD2	1:A:118:LEU:HB2	2.46	0.51
1:C:40:VAL:HG12	1:C:42:GLN:HE21	1.77	0.49
1:O:27:GLU:HG2	1:O:42:GLN:NE2	2.27	0.49



	A L C	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:O:69:LEU:HB3	1:O:96:PRO:HG3	1.93	0.49
1:O:19:ILE:HD12	1:0:21:ALA:O	2.13	0.48
1:K:145:SER:OG	1:K:148:GLN:HG3	2.14	0.48
1:M:19:ILE:O	1:M:74:ILE:HG12	2.14	0.47
1:C:169:ARG:HD2	5:C:457:HOH:O	2.13	0.47
1:K:117:GLY:HA2	2:L:603:DG:OP2	2.15	0.47
1:K:4:ILE:HG21	1:K:149:ILE:HG22	1.95	0.46
1:C:178:VAL:HG23	5:C:462:HOH:O	2.16	0.46
1:E:169:ARG:HD2	5:E:429:HOH:O	2.16	0.46
1:K:79:HIS:CD2	1:K:118:LEU:HB2	2.51	0.45
1:K:132:LYS:HA	1:K:151:TYR:HE2	1.82	0.45
1:E:60:LYS:HA	1:E:60:LYS:HD2	1.70	0.45
1:C:92:PHE:HB2	5:C:419:HOH:O	2.16	0.45
1:E:181:ALA:HB3	1:M:188:MET:HG2	1.98	0.45
1:K:164:ASP:OD2	5:K:401:HOH:O	2.21	0.45
1:0:25:ASP:OD2	1:0:155:ASP:HB3	2.17	0.45
1:M:79:HIS:CD2	1:M:118:LEU:HB2	2.53	0.44
1:C:193:ASP:OD1	1:C:198:SER:HA	2.17	0.44
1:A:19:ILE:HD12	1:A:21:ALA:O	2.18	0.44
1:C:167:GLU:OE2	5:C:401:HOH:O	2.20	0.44
1:A:188:MET:HG2	1:G:181:ALA:HB3	2.00	0.43
1:O:64:ASN:OD1	1:O:67:ARG:NH2	2.51	0.43
1:C:188:MET:HG2	1:K:181:ALA:HB3	2.00	0.43
1:I:115:ARG:HE	1:I:115:ARG:HB3	1.67	0.43
1:K:60:LYS:HA	1:K:60:LYS:HD2	1.82	0.42
1:C:181:ALA:CB	1:K:188:MET:HG3	2.50	0.42
1:G:40:VAL:HG12	1:G:42:GLN:HE21	1.85	0.42
1:O:54:GLN:OE1	1:0:144:LEU:HD23	2.20	0.41
1:K:23:ALA:O	1:K:43:LEU:HA	2.19	0.41
1:A:181:ALA:CB	1:G:188:MET:HG2	2.48	0.41
1:M:19:ILE:HD12	1:M:21:ALA:O	2.21	0.41
1:A:146:ARG:O	1:A:150:GLU:HG3	2.20	0.41
1:K:139:TRP:CZ3	1:K:148:GLN:HB3	2.56	0.41
1:C:114:ASP:HB3	5:J:706:HOH:O	2.21	0.40
1:C:180:LYS:HE3	1:C:184:GLN:HE22	1.87	0.40

Continued from previous page...

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percer	ntiles
1	А	199/207~(96%)	198 (100%)	1 (0%)	0	100	100
1	С	204/207~(99%)	203 (100%)	1 (0%)	0	100	100
1	Е	203/207~(98%)	201 (99%)	2(1%)	0	100	100
1	G	204/207~(99%)	203 (100%)	1 (0%)	0	100	100
1	Ι	204/207~(99%)	202 (99%)	2(1%)	0	100	100
1	K	204/207~(99%)	201 (98%)	3~(2%)	0	100	100
1	М	204/207~(99%)	202 (99%)	2(1%)	0	100	100
1	Ο	205/207~(99%)	$199 \ (97\%)$	5(2%)	1 (0%)	25	28
All	All	1627/1656~(98%)	1609 (99%)	17 (1%)	1 (0%)	48	57

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	0	131	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	175/180~(97%)	171 (98%)	4 (2%)	45	59
1	С	179/180~(99%)	176 (98%)	3 (2%)	56	71
1	Ε	178/180~(99%)	174 (98%)	4 (2%)	47	61
1	G	178/180~(99%)	171 (96%)	7 (4%)	27	37



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	Ι	176/180~(98%)	174 (99%)	2(1%)	70 82
1	Κ	178/180~(99%)	175~(98%)	3~(2%)	56 71
1	М	179/180~(99%)	176~(98%)	3(2%)	56 71
1	Ο	180/180~(100%)	174 (97%)	6 (3%)	33 44
All	All	1423/1440~(99%)	1391~(98%)	32 (2%)	47 61

Continued from previous page...

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	68	LEU
1	А	171	LYS
1	А	173	GLU
1	А	206	SER
1	С	142	GLU
1	С	173	GLU
1	С	206	SER
1	Е	42	GLN
1	Е	97	ASP
1	Е	173	GLU
1	Е	206	SER
1	G	42	GLN
1	G	68	LEU
1	G	72	ARG
1	G	97	ASP
1	G	131	SER
1	G	173	GLU
1	G	184	GLN
1	Ι	42	GLN
1	Ι	173	GLU
1	Κ	42	GLN
1	Κ	173	GLU
1	Κ	184	GLN
1	М	2	THR
1	М	42	GLN
1	М	173	GLU
1	0	0	SER
1	0	3	GLU
1	Ο	61	SER
1	0	119	LYS
1	0	173	GLU
1	0	206	SER



Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	42	GLN
1	С	34	HIS
1	С	42	GLN
1	С	184	GLN
1	G	42	GLN
1	Κ	16	ASN
1	Κ	135	GLN
1	М	34	HIS
1	М	184	GLN
1	0	34	HIS
1	0	42	GLN
1	0	191	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 19 ligands modelled in this entry, 19 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	# RSRZ > 2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	А	203/207~(98%)	-0.41	4 (1%) 64 61	32, 41, 61, 95	0
1	С	206/207~(99%)	-0.44	2 (0%) 79 76	35, 43, 58, 88	0
1	Ε	205/207~(99%)	-0.41	1 (0%) 87 85	34, 43, 64, 80	0
1	G	206/207~(99%)	-0.25	3 (1%) 71 68	34, 49, 65, 80	0
1	Ι	206/207~(99%)	-0.24	7 (3%) 48 45	36, 44, 75, 110	0
1	Κ	206/207~(99%)	-0.07	4 (1%) 66 62	38, 55, 79, 104	0
1	М	206/207~(99%)	-0.13	2 (0%) 79 76	37, 53, 73, 92	0
1	Ο	207/207~(100%)	0.14	7 (3%) 48 45	38, 59, 86, 117	0
2	В	2/2~(100%)	-1.00	0 100 100	43, 43, 43, 43	0
2	D	2/2~(100%)	-0.92	0 100 100	47, 47, 47, 48	0
2	F	2/2~(100%)	-1.16	0 100 100	45, 45, 45, 47	0
2	Η	2/2~(100%)	-0.98	0 100 100	47, 47, 47, 49	0
2	J	2/2~(100%)	-0.94	0 100 100	47, 47, 47, 51	0
2	L	2/2~(100%)	-0.74	0 100 100	51,51,51,54	0
2	Ν	2/2~(100%)	-0.96	0 100 100	49, 49, 49, 50	0
2	Р	2/2~(100%)	-0.80	0 100 100	57, 57, 57, 59	0
All	All	1661/1672~(99%)	-0.23	30 (1%) 67 64	32, 47, 74, 117	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	0	130	ILE	5.9
1	Ι	2	THR	4.3
1	М	2	THR	3.7
1	С	2	THR	3.6
1	Ι	1	MET	3.5



9F7	L
-----	---

Mol	Chain	Res	Type	RSRZ
1	М	1	MET	3.4
1	G	134	GLN	3.3
1	Ι	130	ILE	3.2
1	0	1	MET	3.1
1	G	1	MET	3.1
1	0	135	GLN	3.0
1	А	1	MET	3.0
1	G	136	SER	3.0
1	К	1	MET	2.8
1	А	133	GLN	2.6
1	Ι	135	GLN	2.5
1	Ι	151	TYR	2.5
1	А	151	TYR	2.5
1	0	132	LYS	2.5
1	0	136	SER	2.3
1	Κ	136	SER	2.3
1	Κ	128	VAL	2.2
1	0	2	THR	2.2
1	С	1	MET	2.2
1	Е	136	SER	2.1
1	0	151	TYR	2.1
1	Ι	129	ASN	2.1
1	К	130	ILE	2.1
1	А	137	SER	2.1
1	Ι	155	ASP	2.0

Continued from previous page...

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



9F7L	1
------	---

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
3	NA	0	302	1/1	0.71	0.11	56, 56, 56, 56	1
3	NA	А	303	1/1	0.81	0.14	44,44,44,44	1
3	NA	Е	301	1/1	0.83	0.12	$50,\!50,\!50,\!50$	1
3	NA	С	302	1/1	0.87	0.09	45,45,45,45	1
3	NA	0	301	1/1	0.89	0.08	63,63,63,63	1
3	NA	Ι	302	1/1	0.90	0.07	45,45,45,45	1
3	NA	М	302	1/1	0.90	0.13	$51,\!51,\!51,\!51$	1
3	NA	Ι	301	1/1	0.92	0.07	$51,\!51,\!51,\!51$	1
3	NA	K	301	1/1	0.93	0.10	$61,\!61,\!61,\!61$	1
3	NA	А	301	1/1	0.94	0.08	43,43,43,43	1
3	NA	С	301	1/1	0.94	0.12	$50,\!50,\!50,\!50$	1
3	NA	G	301	1/1	0.95	0.06	$54,\!54,\!54,\!54$	1
3	NA	М	301	1/1	0.96	0.05	56, 56, 56, 56	1
3	NA	Е	302	1/1	0.96	0.06	43,43,43,43	1
3	NA	А	302	1/1	0.96	0.06	48,48,48,48	0
3	NA	G	302	1/1	0.96	0.06	45,45,45,45	1
3	NA	K	302	1/1	0.98	0.04	$51,\!51,\!51,\!51$	1
4	CL	A	304	1/1	0.99	0.05	41,41,41,41	0
4	CL	С	303	1/1	0.99	0.05	46,46,46,46	0

6.5 Other polymers (i)

There are no such residues in this entry.

