



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 27, 2023 – 02:31 AM EDT

PDB ID : 3F9Y  
Title : Structural Insights into Lysine Multiple Methylation by SET Domain Methyltransferases, SET8-Y334F / H4-Lys20me1 / AdoHcy  
Authors : Couture, J.-F.; Dirk, L.M.A.; Brunzelle, J.S.; Houtz, R.L.; Trievel, R.C.  
Deposited on : 2008-11-14  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	166	
1	B	166	
2	E	10	
2	F	10	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2978 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-lysine N-methyltransferase SETD8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	159	1271	795	223	247	6	0	1	0
1	B	160	1295	807	230	252	6	0	6	0


There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	187	GLY	-	expression tag	UNP Q9NQR1
A	188	ALA	-	expression tag	UNP Q9NQR1
A	189	MET	-	expression tag	UNP Q9NQR1
A	190	GLY	-	expression tag	UNP Q9NQR1
A	334	PHE	TYR	engineered mutation	UNP Q9NQR1
B	187	GLY	-	expression tag	UNP Q9NQR1
B	188	ALA	-	expression tag	UNP Q9NQR1
B	189	MET	-	expression tag	UNP Q9NQR1
B	190	GLY	-	expression tag	UNP Q9NQR1
B	334	PHE	TYR	engineered mutation	UNP Q9NQR1

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	E	7	66	42	17	7	0	0	0
2	F	8	67	42	17	8	0	0	0

- Molecule 3 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: C<sub>14</sub>H<sub>20</sub>N<sub>6</sub>O<sub>5</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	A	1	26	14	6	5	1	0	0
3	B	1	26	14	6	5	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	B	227	227	227	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone-lysine N-methyltransferase SETD8




- Molecule 1: Histone-lysine N-methyltransferase SETD8



- Molecule 2: Histone H4



- Molecule 2: Histone H4



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	44.40Å 45.00Å 52.40Å 91.40° 115.30° 89.80°	Depositor
Resolution (Å)	23.68 – 1.50 23.68 – 1.65	Depositor EDS
% Data completeness (in resolution range)	100.0 (23.68-1.50) 96.3 (23.68-1.65)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	12.01 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.194 , 0.220 0.184 , 0.213	Depositor DCC
$R_{free}$ test set	2125 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.1	Xtriage
Anisotropy	0.073	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 55.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for h,-k,-h-l 0.014 for -h,k,-l 0.009 for -h,-k,h+l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	2978	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.34% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAH, MLZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.94	1/1299 (0.1%)	1.01	3/1743 (0.2%)
1	B	0.92	1/1354 (0.1%)	1.01	6/1813 (0.3%)
2	E	1.01	0/55	0.84	0/69
2	F	0.90	0/56	0.97	0/72
All	All	0.93	2/2764 (0.1%)	1.00	9/3697 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	194	SER	CB-OG	6.29	1.50	1.42
1	B	273	TYR	CE2-CZ	6.00	1.46	1.38

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	339	ARG	NE-CZ-NH1	9.43	125.01	120.30
1	A	339	ARG	NE-CZ-NH2	-8.84	115.88	120.30
1	B	324	ARG	NE-CZ-NH2	7.29	123.94	120.30
1	A	335	ASP	CB-CG-OD2	-5.82	113.07	118.30
1	B	318	LEU	CB-CG-CD1	-5.68	101.34	111.00
1	B	289	GLU	OE1-CD-OE2	-5.35	116.89	123.30
1	B	245	TYR	CZ-CE2-CD2	-5.29	115.04	119.80
1	B	292	ARG	NE-CZ-NH2	-5.21	117.69	120.30
1	B	325	ASP	CB-CG-OD1	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1271	0	1249	19	0
1	B	1295	0	1267	17	0
2	E	66	0	78	2	0
2	F	67	0	69	0	0
3	A	26	0	19	0	0
3	B	26	0	19	0	0
4	B	227	0	0	16	1
All	All	2978	0	2701	36	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:313:ASP:HB2	4:B:378:HOH:O	1.59	1.03
1:B:214:LYS:HB3	4:B:373:HOH:O	1.63	0.98
1:A:313:ASP:HB2	4:B:111:HOH:O	1.68	0.92
1:A:288:ARG:HH11	1:A:288:ARG:CG	1.86	0.88
1:A:288:ARG:HH11	1:A:288:ARG:HG3	1.40	0.84
1:A:288:ARG:HD3	4:B:170:HOH:O	1.78	0.82
1:A:346:ALA:HB2	2:E:16:LYS:HE3	1.62	0.80
1:B:309:LEU:HD23	1:B:318:LEU:HD23	1.67	0.75
4:B:374:HOH:O	2:E:19:ARG:HD3	1.91	0.70
1:B:339[B]:ARG:NH2	4:B:183:HOH:O	2.27	0.67
1:A:288:ARG:HG3	1:A:288:ARG:NH1	2.12	0.65
1:B:235:GLN:OE1	4:B:112:HOH:O	2.16	0.62
1:A:197:GLU:O	1:A:201:GLU:HG2	2.04	0.57
1:B:214:LYS:HG2	1:B:216:GLU:HG3	1.87	0.55
1:A:222:LEU:HD13	1:B:274:TYR:CG	2.40	0.55
1:A:277:TYR:OH	1:A:311:ASP:OD2	2.14	0.54
1:B:319:ILE:HD12	1:B:321:ILE:HD11	1.90	0.54
1:B:214:LYS:CB	4:B:373:HOH:O	2.36	0.53
1:B:308[B]:LYS:HG2	1:B:319:ILE:HG13	1.90	0.53
1:A:280:LYS:HD3	4:B:381:HOH:O	2.09	0.52

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288:ARG:CG	1:A:288:ARG:NH1	2.57	0.51
1:B:231:ILE:HD13	1:B:331:GLU:HA	1.93	0.51
1:B:308[B]:LYS:HE2	1:B:309:LEU:O	2.11	0.50
1:A:266:PRO:HG3	4:B:173:HOH:O	2.12	0.49
1:A:335:ASP:OD2	1:A:339:ARG:NH2	2.30	0.49
1:A:200:SER:O	1:A:204:LYS:HG2	2.15	0.47
1:B:309:LEU:HD23	1:B:309:LEU:HA	1.77	0.47
1:B:256:LYS:HD2	4:B:150:HOH:O	2.15	0.47
1:B:235:GLN:CD	4:B:112:HOH:O	2.55	0.45
1:B:251:GLU:HB2	4:B:361:HOH:O	2.16	0.45
1:A:352:HIS:CD2	4:B:88:HOH:O	2.71	0.43
1:A:313:ASP:CB	4:B:111:HOH:O	2.46	0.43
1:B:280:LYS:HE2	4:B:46:HOH:O	2.18	0.41
1:A:346:ALA:O	1:A:348:PRO:HD3	2.20	0.40
1:A:277:TYR:OH	1:A:314:GLY:HA2	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:121:HOH:O	4:B:127:HOH:O[1_556]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	158/166 (95%)	154 (98%)	4 (2%)	0	100 100
1	B	164/166 (99%)	160 (98%)	4 (2%)	0	100 100
2	E	4/10 (40%)	4 (100%)	0	0	100 100
2	F	5/10 (50%)	5 (100%)	0	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	331/352 (94%)	323 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	136/139 (98%)	132 (97%)	4 (3%)	42	13
1	B	142/139 (102%)	140 (99%)	2 (1%)	67	42
2	E	6/8 (75%)	6 (100%)	0	100	100
2	F	5/8 (62%)	5 (100%)	0	100	100
All	All	289/294 (98%)	283 (98%)	6 (2%)	53	23

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	200	SER
1	A	251	GLU
1	A	273	TYR
1	A	288	ARG
1	B	273	TYR
1	B	341	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	MLZ	F	20	2	8,9,10	1.28	1 (12%)	4,9,11	1.23	1 (25%)
2	MLZ	E	20	2	8,9,10	0.58	0	4,9,11	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLZ	F	20	2	-	0/7/8/10	-
2	MLZ	E	20	2	-	1/7/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	20	MLZ	O-C	2.00	1.27	1.19

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	20	MLZ	CM-NZ-CE	2.15	118.17	111.95

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	20	MLZ	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SAH	B	801	-	24,28,28	1.24	3 (12%)	25,40,40	1.60	4 (16%)
3	SAH	A	801	-	24,28,28	1.59	4 (16%)	25,40,40	1.65	5 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SAH	B	801	-	-	2/11/31/31	0/3/3/3
3	SAH	A	801	-	-	2/11/31/31	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	801	SAH	C8-N7	-4.09	1.27	1.34
3	A	801	SAH	O4'-C1'	3.65	1.46	1.41
3	B	801	SAH	C4-N3	2.96	1.39	1.35
3	A	801	SAH	C4-N3	2.84	1.39	1.35
3	B	801	SAH	O4'-C1'	2.45	1.44	1.41
3	A	801	SAH	OXT-C	-2.33	1.22	1.30
3	B	801	SAH	OXT-C	-2.29	1.23	1.30

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	801	SAH	C5-C6-N6	5.15	128.18	120.35
3	B	801	SAH	C5-C6-N6	4.35	126.97	120.35
3	A	801	SAH	C5-C6-N1	-3.49	112.43	120.35
3	B	801	SAH	C4-C5-N7	-3.39	105.86	109.40
3	B	801	SAH	OXT-C-CA	2.45	121.73	113.38
3	B	801	SAH	C5-C6-N1	-2.38	114.95	120.35
3	A	801	SAH	OXT-C-CA	2.32	121.28	113.38
3	A	801	SAH	CB-CG-SD	-2.16	108.45	113.31
3	A	801	SAH	C4-C5-N7	-2.03	107.28	109.40

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	801	SAH	CB-CG-SD-C5'
3	A	801	SAH	CB-CG-SD-C5'
3	B	801	SAH	C-CA-CB-CG
3	A	801	SAH	O-C-CA-N

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	159/166 (95%)	1.15	38 (23%) 0 0	26, 31, 42, 47	1 (0%)
1	B	160/166 (96%)	0.84	28 (17%) 1 1	25, 31, 40, 44	0
2	E	6/10 (60%)	0.93	1 (16%) 1 1	31, 34, 35, 42	0
2	F	7/10 (70%)	1.43	2 (28%) 0 0	29, 32, 39, 41	0
All	All	332/352 (94%)	1.00	69 (20%) 1 1	25, 31, 41, 47	1 (0%)

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	266	PRO	6.1
1	A	320[A]	LEU	5.7
1	A	313	ASP	5.5
1	A	194	SER	5.2
1	A	267	SER	5.1
1	A	297	ILE	5.1
1	A	263	ALA	4.8
1	A	314	GLY	4.7
1	A	196	ALA	4.5
1	B	263	ALA	4.3
1	A	197	GLU	4.2
1	A	261	LEU	4.1
1	B	214	LYS	3.9
1	A	199	GLN	3.8
1	A	265	ASP	3.7
1	B	261	LEU	3.7
1	B	266	PRO	3.7
1	A	268	THR	3.7
2	F	21	VAL	3.7
1	B	320[A]	LEU	3.6
2	E	16	LYS	3.6

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	245	TYR	3.5
1	A	264	GLN	3.5
1	B	194	SER	3.4
2	F	16	LYS	3.4
1	A	277	TYR	3.3
1	B	196	ALA	3.3
1	A	198	LEU	3.3
1	B	273	TYR	3.2
1	A	334	PHE	3.2
1	B	238	ARG	3.2
1	B	336	TYR	3.2
1	A	284	VAL	3.1
1	A	333	LEU	3.0
1	B	297	ILE	2.9
1	B	264	GLN	2.9
1	A	214	LYS	2.9
1	B	195	LYS	2.8
1	B	313[A]	ASP	2.8
1	A	312	ILE	2.8
1	B	260	ALA	2.8
1	A	273	TYR	2.8
1	B	334	PHE	2.7
1	A	315	VAL	2.7
1	B	302	CYS	2.7
1	A	242	VAL	2.7
1	B	284	VAL	2.6
1	A	211	GLU	2.5
1	A	230	VAL	2.5
1	A	245	TYR	2.4
1	A	319	ILE	2.4
1	A	296	LEU	2.4
1	A	274	TYR	2.4
1	B	199	GLN	2.3
1	B	197	GLU	2.3
1	B	345[A]	GLU	2.3
1	A	318	LEU	2.3
1	B	294	GLY	2.3
1	B	272	MET	2.2
1	A	332	LEU	2.2
1	B	296	LEU	2.2
1	A	342	ALA	2.2
1	B	254	ASP	2.2

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	223	ILE	2.1
1	B	352	HIS	2.1
1	A	254	ASP	2.1
1	B	268	THR	2.0
1	A	257	LYS	2.0
1	A	311	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MLZ	F	20	10/11	0.92	0.27	27,29,32,32	0
2	MLZ	E	20	10/11	0.95	0.21	28,29,32,32	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SAH	B	801	26/26	0.95	0.08	20,24,27,29	0
3	SAH	A	801	26/26	0.96	0.08	17,21,26,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around SAH B 801:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.