



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 6, 2023 – 09:54 PM EDT

PDB ID : 4FB0
Title : Structure of *Oceanobacillus iheyensis* group II intron C377G mutant in a ligand-free state in the presence of K⁺ and Mg²⁺
Authors : Marcia, M.; Pyle, A.M.
Deposited on : 2012-05-22
Resolution : 3.22 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1335 (3.24-3.20)
Clashscore	141614	1460 (3.24-3.20)
RSRZ outliers	127900	1291 (3.24-3.20)
RNA backbone	3102	1023 (3.54-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	393	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MG	A	413	-	-	-	X
4	EPE	A	443	-	X	-	-
4	EPE	A	447	-	X	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EPE	A	449	-	X	-	-
5	SPM	A	445	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 8532 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called Group IIC intron.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	393	8415	3755	1561	2707	392	0	0	0

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	25	Total	Mg	0	0
			25	25		

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	18	Total	K	0	0
			18	18		

- Molecule 4 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O S 6 2 3 1	0	0
4	A	1	Total C N O S 12 6 2 3 1	0	0
4	A	1	Total C O S 6 2 3 1	0	0
4	A	1	Total O S 4 3 1	0	0
4	A	1	Total C O S 5 1 3 1	0	0

- Molecule 5 is SPERMINE (three-letter code: SPM) (formula: $C_{10}H_{26}N_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	N	0	0
			14	10	4		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	27	Total	O	0	0
			27	27		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	89.19Å 95.44Å 224.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.16 – 3.22 43.74 – 3.22	Depositor EDS
% Data completeness (in resolution range)	99.6 (49.16-3.22) 99.7 (43.74-3.22)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 3.25Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.184 , 0.230 0.182 , 0.230	Depositor DCC
R_{free} test set	1600 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	115.7	Xtrriage
Anisotropy	0.197	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 82.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8532	wwPDB-VP
Average B, all atoms (Å ²)	132.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K, SPM, MG, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	4/9431 (0.0%)	1.47	282/14712 (1.9%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	64	G	O3'-P	5.96	1.68	1.61
1	A	245	A	O3'-P	-5.54	1.54	1.61
1	A	168	U	O3'-P	-5.38	1.54	1.61
1	A	49	A	O3'-P	-5.31	1.54	1.61

The worst 5 of 282 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	38	C	O5'-P-OP2	-15.36	91.88	105.70
1	A	245	A	P-O3'-C3'	-14.73	102.03	119.70
1	A	222	A	P-O3'-C3'	14.61	137.23	119.70
1	A	47	U	O5'-P-OP1	-14.36	92.78	105.70
1	A	286	C	P-O3'-C3'	13.74	136.19	119.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8415	0	4223	275	0
2	A	25	0	0	0	0
3	A	18	0	0	0	0
4	A	33	0	20	3	0
5	A	14	0	26	2	0
6	A	27	0	0	1	0
All	All	8532	0	4269	276	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 276 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:104:G:H5''	1:A:105:A:H5'	1.40	1.01
1:A:183:A:H2'	1:A:184:A:H8	1.29	0.98
1:A:387:U:C2'	1:A:388:G:H5'	1.99	0.92
1:A:114:G:C6	1:A:115:A:C2	2.58	0.91
1:A:63:A:O2'	1:A:64:G:H5'	1.72	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	391/393 (99%)	142 (36%)	37 (9%)

5 of 142 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	-1	G
1	A	0	G
1	A	1	G
1	A	2	U
1	A	13	U

5 of 37 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	297	A
1	A	366	A
1	A	311	G
1	A	334	U
1	A	115	A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 49 ligands modelled in this entry, 43 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EPE	A	447	-	5,5,15	2.37	2 (40%)	7,7,20	2.03	3 (42%)
4	EPE	A	448	-	0,3,15	-	-	0,3,20	-	-
4	EPE	A	446	-	12,12,15	2.25	2 (16%)	14,16,20	2.07	4 (28%)
5	SPM	A	445	-	13,13,13	0.31	0	12,12,12	0.85	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EPE	A	443	-	5,5,15	2.32	1 (20%)	7,7,20	9.53	5 (71%)
4	EPE	A	449	-	4,4,15	1.45	1 (25%)	5,6,20	5.24	4 (80%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EPE	A	447	-	-	1/3/3/19	-
5	SPM	A	445	-	-	8/11/11/11	-
4	EPE	A	446	-	-	5/6/14/19	1/1/1/1
4	EPE	A	443	-	-	2/3/3/19	-

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	446	EPE	C10-S	-6.19	1.68	1.77
4	A	443	EPE	O3S-S	4.62	1.63	1.47
4	A	447	EPE	O3S-S	4.54	1.63	1.47
4	A	446	EPE	O3S-S	4.34	1.63	1.47
4	A	447	EPE	C10-S	-2.66	1.67	1.77

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	443	EPE	O3S-S-C10	-12.35	85.80	105.77
4	A	443	EPE	O1S-S-C10	11.73	121.04	106.92
4	A	443	EPE	O3S-S-O1S	-10.98	84.44	111.27
4	A	443	EPE	O2S-S-C10	10.88	120.02	106.92
4	A	443	EPE	O3S-S-O2S	-10.20	86.36	111.27

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	443	EPE	C9-C10-S-O2S
4	A	446	EPE	C10-C9-N1-C2
4	A	446	EPE	C10-C9-N1-C6
4	A	446	EPE	C9-C10-S-O1S

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	446	EPE	C9-C10-S-O2S

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	446	EPE	C2-C3-C5-C6-N1-N4

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	446	EPE	2	0
5	A	445	SPM	2	0
4	A	449	EPE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	393/393 (100%)	-0.15	6 (1%) 73 62	62, 116, 265, 326	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	390	G	18.6
1	A	214	U	3.3
1	A	237	A	3.2
1	A	213	G	2.5
1	A	181	A	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	K	A	441	1/1	0.54	0.40	135,135,135,135	0
2	MG	A	413	1/1	0.60	1.36	85,85,85,85	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	K	A	439	1/1	0.63	0.13	128,128,128,128	0
2	MG	A	425	1/1	0.66	0.16	105,105,105,105	0
2	MG	A	407	1/1	0.71	0.29	76,76,76,76	0
5	SPM	A	445	14/14	0.71	0.58	61,99,116,126	14
3	K	A	433	1/1	0.76	0.24	119,119,119,119	0
3	K	A	430	1/1	0.76	0.16	117,117,117,117	0
3	K	A	434	1/1	0.79	0.12	108,108,108,108	0
3	K	A	442	1/1	0.81	0.14	100,100,100,100	0
2	MG	A	424	1/1	0.82	0.54	86,86,86,86	0
3	K	A	440	1/1	0.84	0.36	133,133,133,133	0
3	K	A	436	1/1	0.84	0.44	135,135,135,135	0
4	EPE	A	448	4/15	0.86	0.73	61,78,86,87	4
4	EPE	A	443	6/15	0.86	0.22	74,109,132,141	6
2	MG	A	415	1/1	0.87	0.33	89,89,89,89	0
3	K	A	431	1/1	0.88	0.09	155,155,155,155	0
3	K	A	428	1/1	0.89	0.31	109,109,109,109	0
2	MG	A	418	1/1	0.90	1.04	87,87,87,87	0
3	K	A	435	1/1	0.90	0.17	117,117,117,117	0
3	K	A	427	1/1	0.91	0.38	101,101,101,101	0
2	MG	A	410	1/1	0.91	0.24	78,78,78,78	0
3	K	A	429	1/1	0.91	0.29	120,120,120,120	0
2	MG	A	422	1/1	0.91	0.38	67,67,67,67	0
3	K	A	432	1/1	0.92	0.34	124,124,124,124	0
2	MG	A	412	1/1	0.92	0.66	81,81,81,81	0
3	K	A	426	1/1	0.93	0.30	148,148,148,148	0
4	EPE	A	446	12/15	0.93	0.32	60,72,83,90	12
2	MG	A	414	1/1	0.94	0.24	69,69,69,69	0
3	K	A	438	1/1	0.94	0.19	110,110,110,110	0
2	MG	A	405	1/1	0.94	0.16	84,84,84,84	0
3	K	A	444	1/1	0.94	0.11	101,101,101,101	0
2	MG	A	403	1/1	0.95	0.22	82,82,82,82	0
2	MG	A	419	1/1	0.95	0.21	88,88,88,88	0
4	EPE	A	447	6/15	0.95	0.73	65,79,87,87	6
2	MG	A	420	1/1	0.95	0.23	65,65,65,65	0
2	MG	A	421	1/1	0.95	0.13	129,129,129,129	0
2	MG	A	416	1/1	0.96	0.50	81,81,81,81	0
2	MG	A	408	1/1	0.97	0.67	90,90,90,90	0
2	MG	A	404	1/1	0.97	0.28	79,79,79,79	0
4	EPE	A	449	5/15	0.97	0.39	93,101,118,119	5
2	MG	A	423	1/1	0.97	0.30	97,97,97,97	0
2	MG	A	409	1/1	0.98	0.11	96,96,96,96	0
2	MG	A	401	1/1	0.98	0.42	69,69,69,69	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	K	A	437	1/1	0.98	0.15	119,119,119,119	0
2	MG	A	417	1/1	0.98	0.25	79,79,79,79	0
2	MG	A	411	1/1	0.98	0.30	60,60,60,60	0
2	MG	A	406	1/1	0.99	0.38	74,74,74,74	0
2	MG	A	402	1/1	1.00	0.33	81,81,81,81	0

6.5 Other polymers [i](#)

There are no such residues in this entry.