



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 25, 2022 – 08:26 PM EDT

PDB ID : 7FPB
Title : PanDDA analysis group deposition – Aar2/RNaseH in complex with fragment P09C03 from the F2X-Universal Library
Authors : Barthel, T.; Wollenhaupt, J.; Lima, G.M.A.; Wahl, M.C.; Weiss, M.S.
Deposited on : 2022-08-26
Resolution : 2.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.31.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	10434 (2.04-2.00)
Clashscore	141614	11643 (2.04-2.00)
Ramachandran outliers	138981	11493 (2.04-2.00)
Sidechain outliers	138945	11492 (2.04-2.00)
RSRZ outliers	127900	10220 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	
2	B	308	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	W4L	A	2101	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9189 atoms, of which 4524 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-splicing factor 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	237	4068	1287	2060	336	373	12	18	21	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1833	GLY	-	expression tag	UNP P33334
A	1834	ALA	-	expression tag	UNP P33334
A	1835	MET	-	expression tag	UNP P33334

- Molecule 2 is a protein called A1 cistron-splicing factor AAR2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
2	B	300	5044	1654	2464	421	485	20	0	17	0

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P32357
B	-2	ALA	-	expression tag	UNP P32357
B	-1	MET	-	expression tag	UNP P32357
B	0	ALA	-	expression tag	UNP P32357
B	166	SER	LEU	conflict	UNP P32357
B	167	SER	LYS	conflict	UNP P32357
B	?	-	LEU	deletion	UNP P32357
B	?	-	GLN	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	GLY	deletion	UNP P32357
B	?	-	SER	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	MET	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ASN	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	170	SER	ASP	conflict	UNP P32357

- Molecule 3 is N²-[(6-chloropyridin-3-yl)methyl]-N,N²-diethylglycinamide (three-letter code: W4L) (formula: C₁₂H₁₈ClN₃O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Cl	N	O		
3	A	1	17	12	1	3	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	26	Total	O	0	0
			26	26		
4	B	34	Total	O	0	0
			34	34		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pre-mRNA-splicing factor 8



- Molecule 2: A1 cistron-splicing factor AAR2



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	89.20Å 81.73Å 92.84Å 90.00° 108.32° 90.00°	Depositor
Resolution (Å)	29.40 – 2.02 29.40 – 2.02	Depositor EDS
% Data completeness (in resolution range)	99.4 (29.40-2.02) 99.5 (29.40-2.02)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 2.03Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.237 , 0.289 0.263 , 0.310	Depositor DCC
R_{free} test set	2000 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	45.4	Xtrriage
Anisotropy	0.658	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 60.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9189	wwPDB-VP
Average B, all atoms (Å ²)	83.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: W4L

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.75	3/2149 (0.1%)	0.98	9/2911 (0.3%)
2	B	0.75	1/2739 (0.0%)	0.86	2/3699 (0.1%)
All	All	0.75	4/4888 (0.1%)	0.92	11/6610 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	62	CYS	CB-SG	-6.58	1.71	1.82
1	A	1918[A]	SER	C-N	5.79	1.47	1.34
1	A	1918[B]	SER	C-N	5.79	1.47	1.34
1	A	2007	ARG	CZ-NH2	5.41	1.40	1.33

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2033	THR	CA-C-N	-7.37	100.98	117.20
1	A	2008	LEU	CB-CG-CD2	-6.28	100.33	111.00
1	A	1888	HIS	CB-CA-C	5.97	122.33	110.40
1	A	2038	HIS	CB-CA-C	-5.70	98.99	110.40
2	B	63	ARG	NE-CZ-NH2	-5.65	117.47	120.30
1	A	2034	ILE	N-CA-C	-5.54	96.05	111.00
2	B	199	LEU	CB-CG-CD1	-5.53	101.60	111.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1937	ARG	NE-CZ-NH2	-5.24	117.68	120.30
1	A	1994	ASP	CB-CG-OD1	5.24	123.01	118.30
1	A	1931[A]	LYS	C-N-CA	-5.13	108.88	121.70
1	A	1931[B]	LYS	C-N-CA	-5.13	108.88	121.70

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1930	PRO	Mainchain
1	A	1931[A]	LYS	Mainchain
1	A	1931[B]	LYS	Mainchain
1	A	2033	THR	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2008	2060	1974	55	2
2	B	2580	2464	2396	47	2
3	A	17	0	0	0	0
4	A	26	0	0	5	0
4	B	34	0	0	3	0
All	All	4665	4524	4370	94	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1947:HIS:CE1	2:B:194:GLU:HB2	1.73	1.22
1:A:1947:HIS:CE1	2:B:194:GLU:CB	2.24	1.21
1:A:1885:LYS:HD2	1:A:2001[A]:SER:OG	1.47	1.14
1:A:1885:LYS:CE	1:A:2001[A]:SER:OG	2.05	1.05

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1885:LYS:CD	1:A:2001[A]:SER:OG	2.04	1.04
1:A:1947:HIS:HE1	2:B:194:GLU:HB2	1.18	0.95
1:A:1993:ASP:O	4:A:2202:HOH:O	1.92	0.87
1:A:1885:LYS:HD2	1:A:2001[B]:SER:CB	2.04	0.87
1:A:1885:LYS:HD2	1:A:2001[A]:SER:CB	2.05	0.87
1:A:1947:HIS:CE1	2:B:194:GLU:CG	2.64	0.80
2:B:287:ARG:O	2:B:291:ILE:HD13	1.82	0.79
1:A:1885:LYS:HE3	1:A:2001[A]:SER:OG	1.81	0.79
1:A:2039:LEU:HD23	1:A:2039:LEU:N	2.01	0.76
1:A:2027:LEU:CD2	1:A:2034:ILE:HD11	2.15	0.75
1:A:1885:LYS:HD2	1:A:2001[B]:SER:HB3	1.66	0.75
1:A:1947:HIS:CE1	2:B:194:GLU:HB3	2.23	0.74
2:B:1:MET:N	4:B:401:HOH:O	2.23	0.72
1:A:1973:LYS:NZ	4:A:2203:HOH:O	2.16	0.71
2:B:131:ARG:NH2	2:B:177[B]:ASN:OD1	2.28	0.66
1:A:2064:GLY:O	1:A:2068:ASN:N	2.30	0.65
1:A:1843:LEU:HD23	1:A:1849:LYS:HD2	1.78	0.65
1:A:2039:LEU:HD23	1:A:2039:LEU:H	1.63	0.62
1:A:1947:HIS:ND1	2:B:194:GLU:HG2	2.17	0.59
1:A:1996:LEU:HD23	1:A:1996:LEU:N	2.17	0.59
1:A:1885:LYS:HD2	1:A:2001[A]:SER:HG	1.67	0.59
1:A:2045:ASP:O	1:A:2049:ILE:HG12	2.03	0.58
2:B:1:MET:N	2:B:38:ILE:HD11	2.19	0.58
1:A:2027:LEU:HD21	1:A:2034:ILE:HD11	1.85	0.58
1:A:2058:LEU:C	1:A:2058:LEU:HD23	2.25	0.58
1:A:1931[B]:LYS:HE2	1:A:1931[B]:LYS:HA	1.85	0.57
1:A:2069:VAL:HG12	1:A:2069:VAL:O	2.03	0.57
2:B:206:ASN:O	2:B:211[A]:GLN:HG3	2.06	0.55
2:B:2:ASN:OD1	2:B:62:CYS:HB3	2.07	0.55
2:B:300:ASN:O	2:B:303:HIS:NE2	2.40	0.55
1:A:1885:LYS:NZ	1:A:2001[A]:SER:OG	2.40	0.53
1:A:1940:MET:C	1:A:1943:PRO:HD2	2.29	0.53
2:B:70:GLN:HB3	2:B:81:MET:HE1	1.90	0.53
1:A:1895:HIS:O	1:A:1898[A]:VAL:HG22	2.09	0.52
2:B:50:ASP:OD1	2:B:51:ASN:N	2.42	0.52
1:A:1976:ASP:HB3	4:A:2211:HOH:O	2.08	0.51
1:A:1885:LYS:CD	1:A:2001[B]:SER:OG	2.58	0.51
2:B:70:GLN:HB3	2:B:81:MET:CE	2.42	0.50
1:A:1913:THR:HB	1:A:1940:MET:HE1	1.94	0.50
2:B:77:LEU:HD21	2:B:79:LYS:HE3	1.94	0.49
2:B:44[A]:ILE:O	2:B:44[A]:ILE:HG23	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:43:VAL:HG13	2:B:43:VAL:O	2.12	0.49
1:A:1855[A]:THR:HG22	1:A:1855[A]:THR:O	2.12	0.49
1:A:1996:LEU:N	1:A:1996:LEU:CD2	2.76	0.49
2:B:251:CYS:O	2:B:296:SER:HB2	2.13	0.49
1:A:1885:LYS:HD2	1:A:2001[B]:SER:OG	2.06	0.48
2:B:126:ILE:HG12	2:B:226:PHE:CE1	2.48	0.48
2:B:214:PHE:O	2:B:215:LYS:HB2	2.12	0.48
2:B:126:ILE:HG12	2:B:226:PHE:CZ	2.48	0.47
1:A:1947:HIS:CE1	2:B:194:GLU:HG2	2.45	0.47
1:A:2027:LEU:HD22	1:A:2034:ILE:HD11	1.92	0.47
1:A:1947:HIS:ND1	2:B:194:GLU:CG	2.76	0.47
2:B:17:ILE:HD12	2:B:44[B]:ILE:HG13	1.95	0.47
2:B:190:ARG:HG3	2:B:203[B]:TYR:CZ	2.50	0.47
1:A:2053[A]:SER:OG	1:A:2056[A]:ARG:NH2	2.48	0.47
2:B:6:PHE:CD1	2:B:32:GLY:HA2	2.50	0.47
1:A:2034:ILE:HG22	4:A:2203:HOH:O	2.15	0.46
1:A:1878:CYS:HA	1:A:1892:LYS:O	2.17	0.45
1:A:1942:ASP:N	1:A:1943:PRO:CD	2.79	0.45
1:A:1885:LYS:HE3	1:A:2001[A]:SER:HG	1.81	0.45
2:B:110:GLU:O	2:B:110:GLU:HG2	2.17	0.45
2:B:281:ASP:OD1	2:B:281:ASP:N	2.48	0.45
2:B:1:MET:HB3	2:B:35:ASP:HA	1.99	0.44
2:B:247:ILE:HG22	2:B:292:CYS:SG	2.58	0.44
2:B:1:MET:H2	2:B:38:ILE:HD11	1.81	0.44
2:B:37:PRO:HD3	2:B:105:TYR:CD1	2.53	0.44
2:B:104:SER:O	2:B:105:TYR:C	2.54	0.43
1:A:1844:PHE:CD2	1:A:1844:PHE:N	2.86	0.43
1:A:2028:SER:OG	1:A:2029:ASP:N	2.51	0.43
1:A:1859:ARG:HH12	1:A:1979[A]:MET:CE	2.31	0.43
2:B:293:LEU:HD22	2:B:306:GLU:HB2	2.01	0.43
1:A:2047:GLN:O	1:A:2051:ILE:HG12	2.19	0.42
2:B:40:HIS:CD2	4:B:421:HOH:O	2.71	0.42
2:B:190:ARG:HG3	2:B:203[B]:TYR:CE2	2.55	0.42
2:B:250:ILE:HG21	2:B:264:LEU:HD22	2.02	0.41
1:A:2011:LEU:HD23	1:A:2011:LEU:HA	1.80	0.41
1:A:2058:LEU:HD23	1:A:2058:LEU:O	2.20	0.41
1:A:2036:SER:HA	4:A:2203:HOH:O	2.19	0.41
2:B:291:ILE:O	2:B:296:SER:HB3	2.21	0.41
1:A:1846:ASN:HD22	1:A:1846:ASN:HA	1.71	0.41
1:A:1922:ARG:O	1:A:1922:ARG:HG2	2.21	0.41
2:B:44[A]:ILE:O	2:B:44[A]:ILE:CG2	2.69	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:51:ASN:OD1	2:B:53:SER:HB2	2.21	0.41
2:B:300:ASN:HB2	4:B:405:HOH:O	2.20	0.41
2:B:280:SER:HB3	2:B:313:TYR:CZ	2.56	0.41
2:B:96:PHE:HB2	2:B:102:MET:HE3	2.02	0.40
2:B:2:ASN:HD22	2:B:2:ASN:HA	1.76	0.40
1:A:1850:LEU:HD22	1:A:1881:THR:HG22	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1846:ASN:OD1	2:B:110:GLU:O[4_555]	1.92	0.28
1:A:1846:ASN:H	2:B:112:ASP:OD2[4_555]	1.42	0.18

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	258/258 (100%)	245 (95%)	12 (5%)	1 (0%)	34	28
2	B	315/308 (102%)	298 (95%)	12 (4%)	5 (2%)	9	4
All	All	573/566 (101%)	543 (95%)	24 (4%)	6 (1%)	17	9

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2068	ASN
2	B	62	CYS
2	B	54[A]	MET
2	B	54[B]	MET
2	B	132	LYS
2	B	215	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/233 (102%)	223 (94%)	14 (6%)	19	14
2	B	294/284 (104%)	290 (99%)	4 (1%)	67	70
All	All	531/517 (103%)	513 (97%)	18 (3%)	38	35

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1921	VAL
1	A	1924	LEU
1	A	1927	GLU
1	A	1929	GLN
1	A	1962	ARG
1	A	1973	LYS
1	A	1979[A]	MET
1	A	1979[B]	MET
1	A	1979[C]	MET
1	A	1988	LEU
1	A	1996	LEU
1	A	2035	LYS
1	A	2036	SER
1	A	2039	LEU
2	B	2	ASN
2	B	174	HIS
2	B	258	LYS
2	B	317	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1846	ASN
1	A	1947	HIS
2	B	47	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	W4L	A	2101	-	17,17,17	1.83	2 (11%)	20,21,21	1.61	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	W4L	A	2101	-	-	5/13/13/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2101	W4L	C6-N1	4.99	1.57	1.47
3	A	2101	W4L	C10-CL	-4.17	1.65	1.74

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2101	W4L	C7-C6-N1	-5.53	102.39	113.12
3	A	2101	W4L	C2-C3-N1	-3.32	105.66	113.36

There are no chirality outliers.


All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	2101	W4L	C-C1-N-C2
3	A	2101	W4L	C5-C4-N1-C6
3	A	2101	W4L	C5-C4-N1-C3
3	A	2101	W4L	C7-C6-N1-C4
3	A	2101	W4L	C7-C6-N1-C3

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/258 (91%)	1.64	80 (33%) 0 0	41, 71, 114, 171	0
2	B	300/308 (97%)	1.40	73 (24%) 0 0	43, 70, 114, 171	0
All	All	537/566 (94%)	1.51	153 (28%) 0 0	41, 71, 114, 171	0

All (153) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2068	ASN	8.7
1	A	2036	SER	8.3
2	B	108	ILE	8.1
2	B	243	TRP	7.4
1	A	1848	ILE	7.3
2	B	224	LEU	7.2
2	B	228	PHE	6.8
1	A	2037	TYR	6.6
2	B	109	ASP	6.4
2	B	53	SER	6.3
2	B	246	MET	6.2
2	B	52	SER	6.1
1	A	1878	CYS	5.9
1	A	1965	PHE	5.9
1	A	2034	ILE	5.8
2	B	172	PRO	5.5
2	B	226	PHE	5.5
2	B	247	ILE	5.5
1	A	2060	LEU	5.2
1	A	2063	TYR	5.1
1	A	2044	THR	5.1
1	A	2027	LEU	5.1
2	B	227	ALA	5.0
1	A	1865	THR	4.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1953	ASN	4.8
1	A	1925	PRO	4.8
1	A	1885	LYS	4.7
1	A	1951	PHE	4.7
1	A	2032	ILE	4.6
1	A	2069	VAL	4.6
1	A	2065	ARG	4.6
1	A	2038	HIS	4.5
2	B	316	LEU	4.5
2	B	110	GLU	4.5
1	A	1888	HIS	4.5
2	B	245	ALA	4.4
2	B	58	TYR	4.4
1	A	1926	LYS	4.4
2	B	121	VAL	4.2
1	A	1852	VAL	4.2
1	A	1917	VAL	4.2
2	B	117	LEU	4.1
1	A	1988	LEU	4.1
2	B	1	MET	4.1
2	B	229	LEU	4.0
1	A	1924	LEU	4.0
1	A	1931[A]	LYS	3.8
1	A	2046	GLU	3.8
2	B	43	VAL	3.8
2	B	267	ILE	3.7
2	B	59	TRP	3.7
2	B	281	ASP	3.7
1	A	2057[A]	ASP	3.6
2	B	57	GLY	3.6
2	B	133	ASP	3.6
2	B	171	ASP	3.5
2	B	221	PHE	3.5
2	B	151	GLU	3.5
1	A	2039	LEU	3.4
1	A	1927	GLU	3.4
1	A	1866	PHE	3.4
1	A	2008	LEU	3.4
1	A	2047	GLN	3.3
2	B	181	ILE	3.3
2	B	254	ALA	3.3
1	A	1928	GLU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	242	GLN	3.3
2	B	268	LEU	3.3
2	B	294	TYR	3.2
2	B	233	PHE	3.2
2	B	295	SER	3.2
2	B	231	ALA	3.2
1	A	1948	MET	3.2
2	B	220	TYR	3.2
2	B	178[A]	TYR	3.1
2	B	264	LEU	3.1
1	A	1979[A]	MET	3.1
2	B	234	PHE	3.1
2	B	279	TYR	3.1
1	A	1833	GLY	3.0
1	A	2035	LYS	3.0
1	A	2066	LYS	2.9
1	A	2064	GLY	2.9
2	B	249	LEU	2.9
2	B	317	LEU	2.9
1	A	1861	THR	2.9
1	A	1860	VAL	2.8
1	A	2040	TRP	2.8
1	A	1890	PHE	2.8
2	B	170	SER	2.8
2	B	92	ILE	2.7
1	A	1884	PRO	2.7
1	A	1870	VAL	2.7
1	A	1836	ASN	2.7
1	A	1846	ASN	2.7
2	B	17	ILE	2.7
1	A	1853	ASP	2.7
2	B	54[A]	MET	2.6
2	B	230[A]	ASN	2.6
1	A	1847	ASP	2.6
1	A	1871	ALA	2.6
2	B	277	GLU	2.6
2	B	223	GLU	2.6
1	A	2004	ALA	2.6
2	B	289	TRP	2.6
1	A	2001[A]	SER	2.5
2	B	173	ALA	2.5
1	A	2067	TYR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	44[A]	ILE	2.5
2	B	271	GLN	2.5
1	A	1996	LEU	2.5
2	B	123[A]	MET	2.5
1	A	1886	THR	2.5
1	A	1869	ASN	2.5
2	B	191	PRO	2.5
1	A	1879	ILE	2.5
2	B	84	ARG	2.4
2	B	225	GLN	2.4
1	A	1837	SER	2.4
1	A	2028	SER	2.4
2	B	274	THR	2.4
2	B	193	HIS	2.4
2	B	244	HIS	2.4
1	A	2049	ILE	2.4
1	A	1889	LEU	2.4
1	A	1929	GLN	2.4
2	B	122[A]	GLN	2.4
1	A	1868	GLY	2.4
1	A	1857	VAL	2.3
1	A	1940	MET	2.3
2	B	260	MET	2.3
2	B	120	PHE	2.3
2	B	86	GLY	2.3
1	A	1881	THR	2.3
2	B	69	ILE	2.3
2	B	80	MET	2.2
2	B	211[A]	GLN	2.2
1	A	1947	HIS	2.2
1	A	1954	ILE	2.2
1	A	1964	PRO	2.2
1	A	1991	ILE	2.1
2	B	183	PHE	2.1
2	B	301	SER	2.1
1	A	1961[A]	LEU	2.1
1	A	1851	PHE	2.1
1	A	1867	GLU	2.1
1	A	1835	MET	2.0
1	A	1850	LEU	2.0
1	A	1855[A]	THR	2.0
1	A	1913	THR	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	89	PHE	2.0
1	A	1963	LEU	2.0
2	B	313	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	W4L	A	2101	17/17	0.56	0.57	20,20,20,20	17

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.