

wwPDB EM Validation Summary Report (i)

Nov 13, 2023 – 10:13 AM EST

PDB ID :	ID : 8FPF
EMDB ID :	ID : EMD-29362
Title :	tle : Heterodimeric ABC transporter BmrCD in the inward-facing conformation
	bound to ATP: BmrCD_IF-ATP
Authors :	ors : Qingyu, T.; Hassane, M.
	on : 2023-01-04
Resolution :	on : $3.27 \text{ Å}(\text{reported})$
Authors : posited on :	bound to ATP: BmrCD_IF-ATP ors : Qingyu, T.; Hassane, M.

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


EMDB validation analysis		
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4.02b-467
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.9
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\# \textbf{Entries}) \end{array}$	${f EM\ structures}\ (\#{f Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	D	681	81%	17%	·
2	С	607	83%	9%	8%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 21749 atoms, of which 11125 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Probable multidrug resistance ABC transporter ATP-binding/permease protein YheH.

Mol	Chain	Residues	Atoms					AltConf	Trace	
1	П	666	Total	С	Η	N	0	S	0	0
	D	000	10708	3423	5387	907	974	17	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	154	ALA	CYS	engineered mutation	UNP 007549
D	256	ALA	CYS	engineered mutation	UNP 007549
D	351	ALA	CYS	engineered mutation	UNP 007549
D	592	GLN	GLU	engineered mutation	UNP 007549
D	674	LEU	-	expression tag	UNP 007549
D	675	GLU	-	expression tag	UNP 007549
D	676	HIS	-	expression tag	UNP 007549
D	677	HIS	-	expression tag	UNP 007549
D	678	HIS	-	expression tag	UNP 007549
D	679	HIS	-	expression tag	UNP 007549
D	680	HIS	-	expression tag	UNP 007549
D	681	HIS	-	expression tag	UNP 007549

• Molecule 2 is a protein called Probable multidrug resistance ABC transporter ATP-binding/permease protein YheI.

Mol	Chain	Residues	Atoms					AltConf	Trace	
2	С	558	Total 8861	C 2844	Н 4467	N 732	0 796	S 22	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	-21	MET	-	initiating methionine	UNP 007550
С	-20	GLY	-	expression tag	UNP 007550
С	-19	SER	-	expression tag	UNP 007550


Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
С	-18	SER	-	expression tag	UNP 007550
С	-17	HIS	-	- expression tag	
С	-16	HIS	-	expression tag	UNP 007550
С	-15	HIS	-	expression tag	UNP 007550
С	-14	HIS	-	expression tag	UNP 007550
С	-13	HIS	-	expression tag	UNP 007550
С	-12	HIS	-	expression tag	UNP 007550
С	-11	SER	-	expression tag	UNP 007550
С	-10	SER	-	expression tag	UNP 007550
С	-9	GLY	-	expression tag	UNP 007550
С	-8	LEU	-	expression tag	UNP 007550
С	-7	VAL	-	expression tag	UNP 007550
С	-6	PRO	-	expression tag	UNP 007550
С	-5	ARG	-	expression tag	UNP 007550
С	-4	GLY	-	expression tag	UNP 007550
С	-3	SER	-	expression tag	UNP 007550
С	-2	HIS	-	expression tag	UNP 007550
С	-1	MET	-	expression tag	UNP 007550
С	0	LEU	-	expression tag	UNP 007550
С	1	GLU	-	expression tag	UNP 007550
С	500	GLN	ASP	engineered mutation	UNP 007550

Continued from previous page...

• Molecule 3 is (2S)-3-(hexadecanoyloxy)-2-[(9Z)-octadec-9-enoyloxy]propyl 2-(trimethylamm onio)ethyl phosphate (three-letter code: POV) (formula: C₄₂H₈₂NO₈P) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	AltConf
3	D	1	Total C H O P 63 21 33 8 1	0
3	D	1	Total C H 39 15 24	0
3	D	1	Total C H O P 100 33 58 8 1	0
3	D	1	Total C H 71 25 46	0
3	D	1	Total C H O 53 18 33 2	0
3	D	1	Total C H N O P 119 38 71 1 8 1	0
3	D	1	Total C H N O P 101 33 58 1 8 1	0
3	D	1	Total C H N O P 73 23 40 1 8 1	0
3	D	1	Total C H N O P 103 32 61 1 8 1	0
3	D	1	Total C H 51 18 33	0
3	D	1	Total C H N O P 113 36 67 1 8 1	0
3	D	1	Total C H 71 25 46	0
3	D	1	Total C H 71 25 46	0
3	D	1	Total C H N O P 125 40 75 1 8 1	0
3	С	1	Total C H O P 76 26 41 8 1	0
3	С	1	Total C H N O P 90 29 51 1 8 1	0
3	С	1	Total C H O 39 14 24 1	0
3	С	1	Total C H N O P 101 33 58 1 8 1	0
3	С	1	Total C H O P 115 37 69 8 1	0
3	С	1	Total C H 98 34 64	0
3	С	1	Total C H N O P 95 31 54 1 8 1	0
3	С	1	Total C H O 81 28 49 4	0

Continued on next page...



3

AltConf

0

0

0

Contr	Continuea from previous page									
Mol	Chain	Residues	Atoms							
3	С	1	Total C H							
5	U	1	98 34 64							
2	С	1	Total C H O P							
0	U	1	84 27 48 8 1							

1

Total

64

С

20

Η

34

Continued from previous page


С

• Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$ (labeled as "Ligand of Interest" by depositor).

1

N O P

8 1




Mol	Chain	Residues	Atoms						AltConf
4	р	1	Total	С	Η	Ν	0	Р	0
4	D	1	43	10	12	5	13	3	0
4	С	1	Total	С	Η	Ν	Ο	Р	0
4	U	1	43	10	12	5	13	3	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

 \bullet Molecule 1: Probable multidrug resistance ABC transporter ATP-binding/permease protein YheH



• Molecule 2: Probable multidrug resistance ABC transporter ATP-binding/permease protein YheI





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	255381	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	54	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1600	Depositor
Magnification	Not provided	
Image detector	GATAN K3 $(6k \ge 4k)$	Depositor
Maximum map value	5.439	Depositor
Minimum map value	-0.368	Depositor
Average map value	0.027	Depositor
Map value standard deviation	0.067	Depositor
Recommended contour level	0.43	Depositor
Map size (Å)	330.0, 330.0, 330.0	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.1, 1.1, 1.1	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, POV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	D	0.32	0/5422	0.53	0/7319	
2	С	0.33	0/4486	0.52	0/6072	
All	All	0.32	0/9908	0.52	0/13391	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	5321	5387	5387	79	0
2	С	4394	4467	4465	38	0
3	С	385	556	556	7	0
3	D	462	691	691	9	0
4	С	31	12	12	0	0
4	D	31	12	12	0	0
All	All	10624	11125	11123	118	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 118 close contacts within the same asymmetric unit are listed below, sorted by



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)	
1:D:429:ARG:O	1:D:489:ILE:HD12	2.03	0.59	
1:D:515:LEU:HD13	1:D:523:ASN:HD21	1.73	0.55	
1:D:628:ILE:HG22	1:D:628:ILE:O	2.07	0.54	
1:D:577:PHE:CE2	1:D:590:LEU:HD11	2.43	0.54	
1:D:405:ARG:O	1:D:408:GLU:HG3	2.08	0.54	

their clash magnitude.

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Favoured Allowed		Outliers	Percentiles		
1	D	664/681~(98%)	605~(91%)	59~(9%)	0	100	100
2	С	554/607~(91%)	529~(96%)	25~(4%)	0	100	100
All	All	1218/1288~(95%)	1134 (93%)	84 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Rotameric Outliers		Percentiles		
1	D	567/581~(98%)	567~(100%)	0	100 100	
2	С	471/508~(93%)	471 (100%)	0	100 100	

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed Rotameric Outliers				
All	All	1038/1089~(95%)	1038 (100%)	0	100 100	

There are no protein residues with a non-rotameric sidechain to report.

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such side chains are listed below:

Mol	Chain	Res	Type
1	D	143	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

27 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mal Turna Chain		in Res Li	Link	Bo	Bond lengths			Bond angles					
	Type	Chain	nes	nes	nes	nes	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
3	POV	D	702	-	14,14,51	0.34	0	13, 13, 59	0.16	0				
3	POV	С	605	-	45,45,51	0.40	0	49,50,59	0.53	1 (2%)				
3	POV	D	701	-	29,29,51	0.44	0	33,34,59	0.67	1 (3%)				
3	POV	D	714	-	49,49,51	0.34	0	$55,\!57,\!59$	0.37	0				



Mol	Trune	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	les
10101	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
3	POV	D	708	-	32,32,51	0.41	0	$38,\!40,\!59$	0.41	0
3	POV	С	608	-	30,30,51	0.34	0	$29,\!30,\!59$	0.28	0
3	POV	С	610	-	$35,\!35,\!51$	0.42	0	$39,\!40,\!59$	0.53	1 (2%)
3	POV	С	602	-	38,38,51	0.37	0	44,46,59	0.55	1 (2%)
3	POV	D	704	-	23,23,51	0.21	0	21,21,59	0.11	0
3	POV	С	606	-	32,32,51	0.24	0	$30,\!30,\!59$	0.15	0
3	POV	D	703	-	41,41,51	0.42	0	$45,\!46,\!59$	0.47	1 (2%)
3	POV	С	609	-	32,32,51	0.24	0	30,30,59	0.14	0
4	ATP	С	612	-	26,33,33	0.73	0	31,52,52	0.99	1 (3%)
3	POV	С	601	-	34,34,51	0.44	0	38,39,59	0.68	1 (2%)
3	POV	D	706	-	47,47,51	0.33	0	$53,\!55,\!59$	0.41	0
3	POV	D	709	-	41,41,51	0.36	0	47,49,59	0.42	0
3	POV	С	611	-	29,29,51	0.37	0	$35,\!37,\!59$	0.64	1 (2%)
4	ATP	D	715	-	26,33,33	0.71	0	31,52,52	0.89	1 (3%)
3	POV	D	713	-	23,23,51	0.23	0	21,21,59	0.14	0
3	POV	D	712	-	23,23,51	0.26	0	$21,\!21,\!59$	0.14	0
3	POV	С	604	-	42,42,51	0.43	0	$48,\!50,\!59$	0.48	0
3	POV	С	607	-	$40,\!40,\!51$	0.34	0	$46,\!48,\!59$	0.39	0
3	POV	D	710	-	17,17,51	0.19	0	16, 16, 59	0.20	0
3	POV	D	711	-	45,45,51	0.33	0	50, 53, 59	0.33	0
3	POV	D	707	-	42,42,51	0.36	0	48,50,59	0.46	0
3	POV	С	603	-	14,14,51	0.48	0	13, 13, 59	0.15	0
3	POV	D	705	-	19,19,51	0.41	0	19,19,59	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	POV	D	702	-	-	4/12/12/55	-
3	POV	С	605	-	-	22/47/47/55	-
3	POV	D	701	-	-	8/31/31/55	-
3	POV	D	714	-	-	15/53/53/55	-
3	POV	D	708	-	-	12/36/36/55	-
3	POV	С	608	-	-	5/26/26/55	-
3	POV	С	610	-	-	11/37/37/55	-
3	POV	С	602	-	-	16/42/42/55	-
3	POV	D	704	-	-	6/19/19/55	-

Continued on next page...



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	POV	С	606	-	-	7/28/28/55	-
3	POV	D	703	-	-	7/43/43/55	-
3	POV	С	609	-	-	17/28/28/55	-
4	ATP	С	612	-	-	2/18/38/38	0/3/3/3
3	POV	С	601	-	-	11/36/36/55	-
3	POV	D	706	-	-	16/51/51/55	-
3	POV	D	709	-	-	19/45/45/55	-
3	POV	С	611	-	-	13/33/33/55	-
4	ATP	D	715	-	-	2/18/38/38	0/3/3/3
3	POV	D	713	-	-	2/19/19/55	-
3	POV	D	712	-	-	5/19/19/55	-
3	POV	С	604	-	-	11/46/46/55	-
3	POV	С	607	-	-	16/44/44/55	-
3	POV	D	710	-	-	4/15/15/55	-
3	POV	D	711	-	-	19/49/49/55	-
3	POV	D	707	-	-	13/46/46/55	-
3	POV	С	603	-	-	4/12/12/55	-
3	POV	D	705	-	-	4/17/17/55	-

Continued from previous page...

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	С	612	ATP	PB-O3B-PG	-3.07	122.28	132.83
3	С	610	POV	O13-P-O14	2.52	120.56	110.68
3	С	611	POV	O21-C21-C22	2.44	117.61	110.80
3	С	605	POV	O13-P-O14	2.40	120.07	110.68
3	D	701	POV	O13-P-O14	2.37	119.97	110.68

There are no chirality outliers.

5	of 271	$\operatorname{torsion}$	outliers	are	listed	below:	
---	----------	--------------------------	----------	-----	--------	--------	--

Mol	Chain	Res	Type	Atoms
3	D	701	POV	O22-C21-O21-C2
3	D	703	POV	C22-C21-O21-C2
3	D	707	POV	C2-C1-O11-P
3	D	707	POV	C3-C2-O21-C21

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	D	707	POV	C22-C21-O21-C2


There are no ring outliers.

11 monomers are involved in 16 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	702	POV	3	0
3	D	701	POV	1	0
3	D	708	POV	1	0
3	С	608	POV	1	0
3	D	704	POV	1	0
3	С	601	POV	1	0
3	С	604	POV	2	0
3	С	607	POV	2	0
3	D	710	POV	1	0
3	D	711	POV	2	0
3	Ċ	603	POV	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient the outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

















































5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-29362. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map




The images above show the map projected in three orthogonal directions.




6.2 Central slices (i)


6.2.1 Primary map



X Index: 150




Y Index: 150



Z Index: 150

6.2.2 Raw map



X Index: 150

Y Index: 150

Z Index: 150

The images above show central slices of the map in three orthogonal directions.



6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 146




Y Index: 137



Z Index: 171

6.3.2 Raw map



X Index: 146

Y Index: 137




The images above show the largest variance slices of the map in three orthogonal directions.




6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



6.4.2 Raw map




The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.43. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is 107 nm^3 ; this corresponds to an approximate mass of 97 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.306 ${\rm \AA^{-1}}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.306 $\mathrm{\AA^{-1}}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estim	ation	criterion (FSC cut-off)
Resolution estimate (A)	0.143	0.5	Half-bit
Reported by author	3.27	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.13	6.89	4.19


*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.13 differs from the reported value 3.27 by more than 10 %



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-29362 and PDB model 8FPF. Per-residue inclusion information can be found in section 3 on page 7.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.43 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.43).



9.4 Atom inclusion (i)



At the recommended contour level, 88% of all backbone atoms, 88% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.43) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8770	0.4590
\mathbf{C}	0.8870	0.4660
D	0.8720	0.4540



