



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 8, 2024 – 04:07 PM EDT

PDB ID : 8FPO
Title : PCSK9 in complex with an inhibitor
Authors : Xu, M.; Chopra, R.
Deposited on : 2023-01-05
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| R_{free} | 164625 | 2511 (3.00-3.00) |
| Clashscore | 180529 | 2866 (3.00-3.00) |
| Ramachandran outliers | 177936 | 2778 (3.00-3.00) |
| Sidechain outliers | 177891 | 2781 (3.00-3.00) |
| RSRZ outliers | 164620 | 2523 (3.00-3.00) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | A | 152 | |
| 2 | B | 540 | |
| 3 | C | 9 | |

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4353 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proprotein convertase subtilisin/kexin type 9.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 1 | A | 92 | 740 | 474 | 133 | 131 | 2 | 0 | 0 | 0 |

- Molecule 2 is a protein called Proprotein convertase subtilisin/kexin type 9.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 2 | B | 477 | 3537 | 2185 | 651 | 670 | 31 | 0 | 0 | 0 |

There are 2 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|---------|------------|
| B | 474 | ILE | VAL | variant | UNP Q8NBP7 |
| B | 670 | GLU | GLY | variant | UNP Q8NBP7 |

- Molecule 3 is a protein called MCR-ALA-7T2-GLY-004-7T2-SER-7T2-0NC inhibitor.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace | |
|-----|-------|----------|-------|----|----|---|----|---------|---------|-------|---|
| | | | Total | C | Cl | N | O | | | | S |
| 3 | C | 9 | 75 | 52 | 3 | 9 | 10 | 1 | 0 | 0 | 0 |

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| 4 | B | 1 | Total | Ca | 0 | 0 |
| | | | 1 | 1 | | |

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proprotein convertase subtilisin/kexin type 9

Chain A:  59% 39%

MET
GLY
THR
VAL
SER
SER
ARG
ARG
SER
TRP
TRP
PRO
LEU
PRO
LEU
LEU
LEU
LEU
LEU
LEU
GLY
PRO
ALA
GLY
ALA
ALA
ALA
GLN
GLU
ASP
GLU
ASP
GLY
ASP
TYR
GLU
GLU
VAL
VAL
LEU
LEU
ALA
LEU
ARG
SER
SER
GLU
GLU
GLU
ASP
GLY
LEU
ALA
GLU
ALA
PRO
GLU
GLU
HIS
GLY
THR

T81
R96
Q101
L133
Q152

- Molecule 2: Proprotein convertase subtilisin/kexin type 9

Chain B:  4% 78% 10% 12%

S153
P163
Y166
R167
ALA
ASP
GLU
TYR
GLN
PRO
PRO
ASP
G176
G177
S178
L179
Y183
I189
R194
V200
R201
D212
THR
ARG
PHE
HIS
ARG
GLN
A220
R237
G244
M247
R248
Q278
P279
V280
Y285
L311
D343
Q344
P345
C375
T377

S401
E410
V423
I424
P446
S447
T448
H449
GLY
ALA
G452
V460
P467
T468
R469
T472
A473
I474
S490
R491
R510
F515
V520
A524
L528
Q531
H537
H542
GLU
ALA
SER
THR
MET
GLY
SER
THR
V550
H553
GLN
GLN
GLY
H557
L571

GLY
THR
HIS
LYS
PRO
PRO
VAL
LEU
ARG
PRO
ARG
ARG
GLN
GLN
P565
C588
V589
I596
G605
L606
P616
ALA
PRO
Q619
T623
E628
L632
L638
P639
GLY
THR
S642
L645
R659
ASP
VAL
SER
THR
THR
GLY
SER
THR
GLU
GLU
A671
V672
T673
S681
ARG

HIS
LEU
ALA
GLN
ALA
SER
GLN
GLU
LEU
GLN

- Molecule 3: MCR-ALA-7T2-GLY-004-7T2-SER-7T2-0NC inhibitor

Chain C:  33% 67%

KCH1
A2
F3
G4
O045
F6
S7
F8
A9

4 Data and refinement statistics

| Property | Value | Source |
|---|---|------------------|
| Space group | P 21 21 21 | Depositor |
| Cell constants a, b, c, α , β , γ | 62.95Å 70.75Å 150.09Å 90.00° 90.00° 90.00° | Depositor |
| Resolution (Å) | 58.05 – 3.00 58.05 – 3.00 | Depositor EDS |
| % Data completeness (in resolution range) | 99.7 (58.05-3.00) 99.7 (58.05-3.00) | Depositor EDS |
| R_{merge} | 0.17 | Depositor |
| R_{sym} | (Not available) | Depositor |
| $\langle I/\sigma(I) \rangle$ ¹ | 3.25 (at 3.01Å) | Xtrriage |
| Refinement program | PHENIX 1 | Depositor |
| R, R_{free} | 0.182 , 0.250 0.189 , 0.250 | Depositor DCC |
| R_{free} test set | 687 reflections (4.87%) | wwPDB-VP |
| Wilson B-factor (Å ²) | 40.2 | Xtrriage |
| Anisotropy | 0.387 | Xtrriage |
| Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²) | 0.34 , 33.9 | EDS |
| L-test for twinning ² | $\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$ | Xtrriage |
| Estimated twinning fraction | No twinning to report. | Xtrriage |
| F_o, F_c correlation | 0.92 | EDS |
| Total number of atoms | 4353 | wwPDB-VP |
| Average B, all atoms (Å ²) | 20.0 | wwPDB-VP |

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, 7T2, MCR, 0NC, 004

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|---------|-------------|---------------|
| | | RMSZ | # Z >5 | RMSZ | # Z >5 |
| 1 | A | 0.65 | 0/757 | 0.77 | 0/1023 |
| 2 | B | 0.63 | 0/3599 | 0.80 | 1/4882 (0.0%) |
| 3 | C | 1.30 | 0/12 | 2.28 | 1/11 (9.1%) |
| All | All | 0.64 | 0/4368 | 0.80 | 2/5916 (0.0%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 2 | B | 0 | 1 |

There are no bond length outliers.

All (2) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|----------|------|-------------|----------|
| 2 | B | 179 | LEU | CA-CB-CG | 6.04 | 129.21 | 115.30 |
| 3 | C | 2 | ALA | CB-CA-C | 5.60 | 118.49 | 110.10 |

There are no chirality outliers.

All (1) planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|---------|
| 2 | B | 244 | GLY | Peptide |

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 740 | 0 | 750 | 1 | 0 |
| 2 | B | 3537 | 0 | 3451 | 29 | 0 |
| 3 | C | 75 | 0 | 25 | 0 | 0 |
| 4 | B | 1 | 0 | 0 | 0 | 0 |
| All | All | 4353 | 0 | 4226 | 30 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 30 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|------------------|------------------|--------------------------|-------------------|
| 2:B:638:LEU:HD23 | 2:B:639:PRO:HD2 | 1.48 | 0.94 |
| 2:B:167:ARG:HB2 | 2:B:447:SER:OG | 1.89 | 0.73 |
| 2:B:490:SER:HB2 | 2:B:520:VAL:HG12 | 1.69 | 0.72 |
| 2:B:194:ARG:O | 2:B:237:ARG:NH1 | 2.30 | 0.64 |
| 2:B:446:PRO:HB2 | 2:B:448:THR:HG22 | 1.80 | 0.63 |

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles |
|-----|-------|---------------|-----------|---------|----------|-------------|
| 1 | A | 90/152 (59%) | 88 (98%) | 2 (2%) | 0 | 100 100 |
| 2 | B | 457/540 (85%) | 447 (98%) | 9 (2%) | 1 (0%) | 44 77 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|---------------|-----------|---------|----------|-------------|-----|
| 3 | C | 3/9 (33%) | 2 (67%) | 1 (33%) | 0 | 100 | 100 |
| All | All | 550/701 (78%) | 537 (98%) | 12 (2%) | 1 (0%) | 44 | 77 |

All (1) Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2 | B | 280 | VAL |

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|---------------|-----------|----------|-------------|-----|
| 1 | A | 79/127 (62%) | 78 (99%) | 1 (1%) | 65 | 85 |
| 2 | B | 381/431 (88%) | 370 (97%) | 11 (3%) | 37 | 70 |
| 3 | C | 1/1 (100%) | 1 (100%) | 0 | 100 | 100 |
| All | All | 461/559 (82%) | 449 (97%) | 12 (3%) | 41 | 72 |

5 of 12 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2 | B | 623 | THR |
| 2 | B | 638 | LEU |
| 2 | B | 672 | VAL |
| 2 | B | 645 | LEU |
| 2 | B | 375 | CYS |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 3 | 004 | C | 5 | 3 | 9,10,11 | 1.02 | 0 | 9,12,14 | 1.17 | 1 (11%) |
| 3 | 0NC | C | 9 | 3 | 5,6,6 | 1.19 | 1 (20%) | 6,7,7 | 1.11 | 1 (16%) |
| 3 | 7T2 | C | 3 | 3 | 12,13,14 | 0.83 | 0 | 15,16,18 | 2.12 | 5 (33%) |
| 3 | 7T2 | C | 6 | 3 | 12,13,14 | 0.81 | 0 | 15,16,18 | 2.07 | 2 (13%) |
| 3 | 7T2 | C | 8 | 3 | 12,13,14 | 0.72 | 0 | 15,16,18 | 2.93 | 4 (26%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|----------|---------|
| 3 | 004 | C | 5 | 3 | - | 0/4/6/8 | 0/1/1/1 |
| 3 | 0NC | C | 9 | 3 | - | 2/6/6/6 | - |
| 3 | 7T2 | C | 3 | 3 | - | 0/5/8/10 | 0/1/1/1 |
| 3 | 7T2 | C | 6 | 3 | - | 0/5/8/10 | 0/1/1/1 |
| 3 | 7T2 | C | 8 | 3 | - | 0/5/8/10 | 0/1/1/1 |

All (1) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|------|-------------|----------|
| 3 | C | 9 | 0NC | C-N1 | 2.43 | 1.36 | 1.33 |

The worst 5 of 13 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|----------|-------|-------------|----------|
| 3 | C | 8 | 7T2 | CG-CB-CA | -9.15 | 100.49 | 113.51 |
| 3 | C | 6 | 7T2 | CB-CA-N | -6.15 | 101.42 | 110.48 |
| 3 | C | 3 | 7T2 | CG-CB-CA | -5.88 | 105.14 | 113.51 |
| 3 | C | 8 | 7T2 | O-C-CA | -3.96 | 114.58 | 124.77 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|---------|------|-------------|----------|
| 3 | C | 8 | 7T2 | CB-CA-N | 3.87 | 116.17 | 110.48 |

There are no chirality outliers.

All (2) torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|------------|
| 3 | C | 9 | 0NC | O-C-CA-CB |
| 3 | C | 9 | 0NC | N1-C-CA-CB |

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed | <RSRZ> | #RSRZ>2 | OWAB(Å ²) | Q<0.9 |
|-----|-------|---------------|--------|---|-----------------------|-------|
| 1 | A | 92/152 (60%) | -0.64 | 0 100 100 | 4, 12, 29, 48 | 0 |
| 2 | B | 477/540 (88%) | -0.24 | 19 (3%) 43 25 | 3, 16, 52, 91 | 0 |
| 3 | C | 3/9 (33%) | -0.24 | 0 100 100 | 14, 14, 14, 28 | 0 |
| All | All | 572/701 (81%) | -0.31 | 19 (3%) 49 29 | 3, 15, 51, 91 | 0 |

The worst 5 of 19 RSRZ outliers are listed below:

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 2 | B | 448 | THR | 5.0 |
| 2 | B | 177 | GLY | 4.1 |
| 2 | B | 571 | LEU | 3.8 |
| 2 | B | 515 | PHE | 3.7 |
| 2 | B | 449 | HIS | 3.6 |

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(Å ²) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|----------------------------|-------|
| 3 | 0NC | C | 9 | 7/7 | 0.91 | 0.09 | 14,24,28,35 | 0 |
| 3 | 7T2 | C | 8 | 13/14 | 0.93 | 0.09 | 9,14,26,45 | 0 |
| 3 | 7T2 | C | 3 | 13/14 | 0.93 | 0.10 | 7,13,15,60 | 0 |
| 3 | 7T2 | C | 6 | 13/14 | 0.95 | 0.08 | 12,15,23,69 | 0 |
| 3 | 004 | C | 5 | 10/11 | 0.99 | 0.04 | 13,15,17,19 | 0 |

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(Å ²) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|----------------------------|-------|
| 4 | CA | B | 701 | 1/1 | 0.84 | 0.16 | 38,38,38,38 | 0 |

6.5 Other polymers [i](#)

There are no such residues in this entry.