



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 20, 2023 – 05:57 AM EDT

PDB ID : 2FR4
Title : Structure of Fab DNA-1 complexed with a stem-loop DNA ligand
Authors : Tanner, J.J.; Ou, Z.
Deposited on : 2006-01-18
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	M	10	
1	N	10	
2	A	214	
2	L	214	
3	B	230	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	H	230	 <p>14% 83% 8% • 7%</p>

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 7283 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*CP*TP*GP*CP*CP*TP*TP*CP*AP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	M	10	198	96	33	60	9	0	0	0
1	N	10	198	96	33	60	9	0	0	0

- Molecule 2 is a protein called antibody light chain FAB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	212	1628	1020	271	331	6	0	0	0
2	A	213	1614	1011	268	329	6	0	0	0

- Molecule 3 is a protein called antibody heavy chain FAB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	H	214	1622	1036	259	318	9	0	0	0
3	B	214	1615	1033	257	316	9	0	0	0

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
H	1	GLN	-	cloning artifact	GB 3399661
H	2	VAL	-	cloning artifact	GB 3399661
H	3	LYS	-	cloning artifact	GB 3399661
H	4	LEU	-	cloning artifact	GB 3399661
H	218	HIS	-	expression tag	GB 3399661
H	219	HIS	-	expression tag	GB 3399661
H	220	HIS	-	expression tag	GB 3399661
H	221	HIS	-	expression tag	GB 3399661

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	222	HIS	-	expression tag	GB 3399661
H	223	HIS	-	expression tag	GB 3399661
B	1	GLN	-	cloning artifact	GB 3399661
B	2	VAL	-	cloning artifact	GB 3399661
B	3	LYS	-	cloning artifact	GB 3399661
B	4	LEU	-	cloning artifact	GB 3399661
B	218	HIS	-	expression tag	GB 3399661
B	219	HIS	-	expression tag	GB 3399661
B	220	HIS	-	expression tag	GB 3399661
B	221	HIS	-	expression tag	GB 3399661
B	222	HIS	-	expression tag	GB 3399661
B	223	HIS	-	expression tag	GB 3399661

- Molecule 4 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C₈H₁₈O₅).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	H	1	Total	0	0
			C		
			8	5	3

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	M	1	Total	0	0
			O		
			1	1	
5	N	3	Total	0	0
			O		
			3	3	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	L	51	Total 51	O 51	0	0
5	H	146	Total 146	O 146	0	0
5	A	57	Total 57	O 57	0	0
5	B	142	Total 142	O 142	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(*CP*TP*GP*CP*CP*TP*TP*CP*AP*G)-3'

Chain M: 





- Molecule 1: 5'-D(*CP*TP*GP*CP*CP*TP*TP*CP*AP*G)-3'

Chain N: 





- Molecule 2: antibody light chain FAB

Chain L: 





- Molecule 2: antibody light chain FAB

Chain A: 








• Molecule 3: antibody heavy chain FAB



• Molecule 3: antibody heavy chain FAB



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	84.76Å 90.45Å 128.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	22.30 – 1.95 22.29 – 1.95	Depositor EDS
% Data completeness (in resolution range)	100.0 (22.30-1.95) 97.5 (22.29-1.95)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.73 (at 1.95Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.207 , 0.256 0.216 , 0.262	Depositor DCC
R_{free} test set	3596 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	30.8	Xtrriage
Anisotropy	0.370	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 49.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7283	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	M	0.93	0/220	1.73	2/337 (0.6%)
1	N	1.02	1/220 (0.5%)	1.77	5/337 (1.5%)
2	A	0.57	0/1652	0.67	0/2251
2	L	0.54	0/1666	0.67	2/2265 (0.1%)
3	B	0.74	2/1661 (0.1%)	0.74	1/2269 (0.0%)
3	H	0.73	1/1668 (0.1%)	0.79	1/2279 (0.0%)
All	All	0.68	4/7087 (0.1%)	0.83	11/9738 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	140	CYS	CB-SG	-7.66	1.69	1.82
3	H	148	GLU	CD-OE2	5.91	1.32	1.25
3	B	195	CYS	CB-SG	-5.13	1.73	1.81
1	N	10	DT	C2-N3	-5.07	1.33	1.37

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	N	6	DT	O4'-C1'-N1	8.75	114.12	108.00
3	H	177	LEU	CA-CB-CG	8.37	134.56	115.30
1	N	10	DT	O4'-C1'-N1	-7.54	102.72	108.00
1	N	6	DT	C6-C5-C7	-7.00	118.70	122.90
3	B	177	LEU	CA-CB-CG	6.48	130.21	115.30
1	M	7	DG	O4'-C1'-N9	5.75	112.03	108.00
1	N	8	DC	C5-C4-N4	5.50	124.05	120.20
2	L	108	ARG	NE-CZ-NH2	-5.41	117.59	120.30
2	L	108	ARG	NE-CZ-NH1	5.35	122.98	120.30
1	M	11	DT	O3'-P-O5'	-5.33	93.87	104.00
1	N	14	DG	O4'-C1'-N9	-5.10	104.43	108.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	M	198	0	115	0	0
1	N	198	0	115	0	0
2	A	1614	0	1501	13	0
2	L	1628	0	1534	10	0
3	B	1615	0	1572	7	0
3	H	1622	0	1579	13	0
4	H	8	0	9	2	0
5	A	57	0	0	1	0
5	B	142	0	0	1	0
5	H	146	0	0	0	0
5	L	51	0	0	0	0
5	M	1	0	0	0	0
5	N	3	0	0	0	0
All	All	7283	0	6425	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:136:LEU:HD11	2:A:146:VAL:HG22	1.76	0.68
3:H:12:VAL:HG21	3:H:82(C):LEU:HD13	1.76	0.67
3:B:143:LYS:NZ	3:B:171:GLN:OE1	2.32	0.61
3:H:41:PRO:O	3:H:43:GLN:NE2	2.33	0.61
2:L:113:PRO:HG2	2:L:205:ILE:HD12	1.81	0.60
3:B:73:LYS:NZ	5:B:358:HOH:O	2.32	0.58
3:H:43:GLN:HE21	3:H:43:GLN:N	2.01	0.58
3:H:5:LEU:HD12	3:H:105:GLN:HE21	1.72	0.55
2:L:11:LEU:HD23	2:L:104:LEU:HD13	1.89	0.55
2:L:103:LYS:HD3	2:L:105:GLU:HG3	1.87	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:135:PHE:C	2:A:136:LEU:HD23	2.28	0.54
2:A:161:ASN:ND2	2:A:177:SER:OG	2.41	0.54
2:L:161:ASN:ND2	2:L:177:SER:OG	2.41	0.54
2:A:2:LEU:HD22	2:A:90:HIS:CE1	2.45	0.51
2:A:136:LEU:HD23	2:A:136:LEU:N	2.27	0.50
3:H:11:LEU:HD22	3:H:110:THR:HB	1.93	0.50
2:L:161:ASN:HD22	2:L:177:SER:HA	1.77	0.50
3:H:18:VAL:HG12	3:H:82(C):LEU:HD11	1.94	0.49
3:H:154:TRP:CZ3	3:H:195:CYS:HB3	2.49	0.48
2:A:136:LEU:HD11	2:A:146:VAL:CG2	2.43	0.46
2:L:52:LYS:HB2	2:L:64:GLY:O	2.15	0.46
2:A:161:ASN:HD22	2:A:177:SER:HA	1.80	0.46
3:B:189:PRO:HB3	3:B:212:PRO:HG3	1.98	0.45
2:A:115:VAL:HG13	2:A:136:LEU:HD22	1.99	0.45
3:H:96:GLY:HA3	3:H:100(A):TYR:HB2	2.00	0.44
2:L:103:LYS:CD	2:L:105:GLU:HG3	2.48	0.44
2:L:138:ASN:HA	2:L:172:THR:HB	1.99	0.44
3:H:5:LEU:HD12	3:H:105:GLN:NE2	2.33	0.44
3:B:154:TRP:CZ3	3:B:195:CYS:HB3	2.53	0.43
3:H:12:VAL:HG21	3:H:82(C):LEU:CD1	2.47	0.43
3:H:149:PRO:HB3	4:H:501:PG4:H41	2.01	0.43
3:H:212:PRO:O	3:H:213:ARG:HB2	2.18	0.43
3:B:47:TRP:CZ2	3:B:49:GLY:HA2	2.55	0.42
2:A:76:ASN:O	2:A:76:ASN:ND2	2.54	0.41
3:H:148:GLU:HB3	4:H:501:PG4:H31	2.02	0.41
3:B:96:GLY:HA3	3:B:100(A):TYR:HB2	2.02	0.41
2:L:48:VAL:HG22	2:L:54:LEU:HD12	2.02	0.41
2:A:106:LEU:HD23	2:A:107:LYS:N	2.35	0.41
2:L:108:ARG:HD3	2:L:109:ALA:O	2.20	0.40
2:A:2:LEU:HD13	5:A:252:HOH:O	2.22	0.40
2:A:149:LYS:HA	2:A:153:SER:O	2.20	0.40
2:A:44:PRO:HD2	3:B:103:TRP:CE3	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	211/214 (99%)	205 (97%)	5 (2%)	1 (0%)	29	17
2	L	210/214 (98%)	202 (96%)	8 (4%)	0	100	100
3	B	210/230 (91%)	206 (98%)	4 (2%)	0	100	100
3	H	210/230 (91%)	206 (98%)	4 (2%)	0	100	100
All	All	841/888 (95%)	819 (97%)	21 (2%)	1 (0%)	51	43

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	77	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	175/188 (93%)	166 (95%)	9 (5%)	24	11
2	L	179/188 (95%)	168 (94%)	11 (6%)	18	7
3	B	182/200 (91%)	175 (96%)	7 (4%)	33	21
3	H	184/200 (92%)	177 (96%)	7 (4%)	33	21
All	All	720/776 (93%)	686 (95%)	34 (5%)	26	13

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	L	2	LEU
2	L	24	ARG
2	L	42	LYS
2	L	52	LYS
2	L	54	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	L	77	SER
2	L	103	LYS
2	L	108	ARG
2	L	142	LYS
2	L	143	ASP
2	L	185	GLU
3	H	11	LEU
3	H	43	GLN
3	H	58	LYS
3	H	73	LYS
3	H	133	ASN
3	H	170	LEU
3	H	177	LEU
2	A	52	LYS
2	A	76	ASN
2	A	96	LEU
2	A	105	GLU
2	A	136	LEU
2	A	142	LYS
2	A	145	ASN
2	A	169	LYS
2	A	191	SER
3	B	5	LEU
3	B	11	LEU
3	B	62	LYS
3	B	105	GLN
3	B	177	LEU
3	B	190	SER
3	B	203	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
2	L	3	GLN
2	L	137	ASN
2	L	161	ASN
2	L	212	ASN
3	H	1	GLN
3	H	43	GLN
3	H	105	GLN
2	A	76	ASN
2	A	89	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	A	137	ASN
2	A	157	ASN
2	A	161	ASN
2	A	210	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PG4	H	501	-	7,7,12	0.54	0	6,6,11	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PG4	H	501	-	-	2/5/5/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	H	501	PG4	C1-C2-O2-C3
4	H	501	PG4	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	501	PG4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	M	10/10 (100%)	1.02	0 100 100	28, 37, 42, 43	0
1	N	10/10 (100%)	1.64	1 (10%) 7 11	30, 37, 44, 44	0
2	A	213/214 (99%)	1.55	69 (32%) 0 0	35, 41, 46, 55	0
2	L	212/214 (99%)	1.28	49 (23%) 0 0	35, 40, 45, 49	0
3	B	214/230 (93%)	1.10	37 (17%) 1 2	34, 40, 45, 52	0
3	H	214/230 (93%)	0.97	32 (14%) 2 3	32, 40, 48, 55	0
All	All	873/908 (96%)	1.23	188 (21%) 0 1	28, 40, 46, 55	0

All (188) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	B	134	SER	6.1
3	H	141	LEU	5.9
2	A	9	ALA	5.8
2	L	133	VAL	5.5
2	A	203	SER	5.4
2	A	135	PHE	5.2
2	A	168	SER	5.0
2	A	133	VAL	4.8
3	H	172	SER	4.8
2	A	8	PRO	4.8
3	H	133	ASN	4.7
3	H	142	VAL	4.7
2	A	212	ASN	4.7
3	B	124	LEU	4.5
2	L	35	TRP	4.5
3	B	37	VAL	4.5
3	B	141	LEU	4.4
2	A	41	GLY	4.3
2	L	213	GLU	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	A	7	SER	4.1
2	A	213	GLU	4.1
3	H	41	PRO	4.1
3	B	41	PRO	4.0
2	L	132	VAL	4.0
2	L	156	GLN	4.0
2	L	36	TYR	3.9
2	A	34	ALA	3.8
2	A	169	LYS	3.7
2	A	136	LEU	3.7
3	B	135	MET	3.6
3	H	1	GLN	3.6
3	B	121	VAL	3.6
2	A	199	LYS	3.6
2	A	30	TYR	3.6
3	B	127	GLY	3.6
3	H	93	VAL	3.6
3	B	172	SER	3.6
3	B	36	TRP	3.5
2	A	35	TRP	3.5
2	L	46	LEU	3.5
2	L	148	TRP	3.4
3	B	186	SER	3.4
2	A	165	ASP	3.4
2	A	73	LEU	3.4
1	N	12	DC	3.3
3	B	140	CYS	3.3
2	L	184	ASP	3.3
3	B	51	ILE	3.3
2	A	88	CYS	3.3
2	L	21	ILE	3.3
2	A	33	LEU	3.2
3	B	100(B)	ALA	3.2
3	B	139	GLY	3.2
3	H	109	VAL	3.2
2	A	69	THR	3.2
3	H	42	GLY	3.1
2	A	134	CYS	3.1
2	A	11	LEU	3.1
2	L	86	TYR	3.1
2	A	75	ILE	3.1
2	A	80	PRO	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	134	CYS	3.1
2	L	34	ALA	3.1
2	A	157	ASN	3.1
3	B	184	PRO	3.0
2	L	178	THR	3.0
3	H	124	LEU	3.0
2	L	56	GLU	3.0
2	A	40	GLN	3.0
2	A	156	GLN	3.0
2	L	33	LEU	2.9
2	L	135	PHE	2.9
2	L	58	VAL	2.9
2	A	36	TYR	2.9
3	B	136	VAL	2.9
2	A	146	VAL	2.9
2	A	26	SER	2.8
2	L	41	GLY	2.8
2	A	78	LEU	2.8
2	A	202	THR	2.8
2	A	205	ILE	2.8
2	A	74	LYS	2.8
2	A	170	ASP	2.8
2	L	157	ASN	2.8
2	L	182	THR	2.8
2	A	148	TRP	2.8
2	L	202	THR	2.7
2	A	204	PRO	2.7
3	B	42	GLY	2.7
3	H	36	TRP	2.7
3	H	97	TYR	2.7
3	B	109	VAL	2.7
2	A	106	LEU	2.7
2	A	98	PHE	2.7
2	L	73	LEU	2.7
2	A	60	SER	2.7
2	A	109	ALA	2.6
3	H	43	GLN	2.6
2	A	25	ALA	2.6
2	A	117	ILE	2.6
2	L	48	VAL	2.6
3	B	33	VAL	2.6
3	B	171	GLN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	B	210	ILE	2.6
3	B	93	VAL	2.6
2	L	57	GLY	2.6
2	L	77	SER	2.6
3	H	90	TYR	2.5
2	A	87	TYR	2.5
3	B	52(A)	PRO	2.5
3	B	82	LEU	2.5
3	B	162	GLY	2.5
2	A	77	SER	2.5
2	A	59	PRO	2.5
3	H	170	LEU	2.5
2	L	188	ARG	2.5
3	B	95	GLY	2.4
2	L	150	ILE	2.4
2	A	58	VAL	2.4
2	L	78	LEU	2.4
2	A	47	LEU	2.4
2	A	71	PHE	2.4
2	L	59	PRO	2.4
2	L	204	PRO	2.4
3	H	171	GLN	2.4
3	H	121	VAL	2.4
2	L	47	LEU	2.4
3	H	78	ALA	2.4
2	A	89	GLN	2.4
2	L	49	TYR	2.3
2	A	86	TYR	2.3
3	B	1	GLN	2.3
3	H	140	CYS	2.3
3	B	142	VAL	2.3
2	A	1	GLU	2.3
2	A	104	LEU	2.3
2	L	160	LEU	2.3
2	A	144	ILE	2.3
3	H	100(C)	MET	2.3
2	L	2	LEU	2.3
2	L	104	LEU	2.3
3	B	90	TYR	2.3
2	L	42	LYS	2.3
2	A	42	LYS	2.3
2	A	132	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	A	196	ALA	2.3
3	H	119	PRO	2.2
2	A	153	SER	2.2
2	L	24	ARG	2.2
3	H	213	ARG	2.2
2	A	70	GLN	2.2
2	A	49	TYR	2.2
2	A	118	PHE	2.2
3	H	80	MET	2.2
2	L	154	GLU	2.2
2	L	196	ALA	2.2
2	A	197	THR	2.2
3	H	100(B)	ALA	2.2
3	H	102	TYR	2.2
3	H	197	VAL	2.2
3	B	206	VAL	2.2
2	A	46	LEU	2.1
2	A	81	GLU	2.1
2	A	139	PHE	2.1
2	L	122	SER	2.1
2	L	88	CYS	2.1
2	L	117	ILE	2.1
2	A	28	ASN	2.1
2	A	150	ILE	2.1
3	H	51	ILE	2.1
3	B	69	LEU	2.1
2	L	87	TYR	2.1
3	B	102	TYR	2.1
3	B	190	SER	2.1
2	A	198	HIS	2.1
3	B	197	VAL	2.1
3	H	48	ILE	2.1
2	L	126	THR	2.1
3	H	34	MET	2.1
2	L	54	LEU	2.1
3	H	177	LEU	2.1
2	L	106	LEU	2.1
3	H	92	CYS	2.1
3	B	80	MET	2.0
3	H	4	LEU	2.0
3	B	53	TYR	2.0
2	L	40	GLN	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	89	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PG4	H	501	8/13	0.62	0.40	56,57,58,58	0

6.5 Other polymers [i](#)

There are no such residues in this entry.