



wwPDB X-ray Structure Validation Summary Report ⓘ

May 17, 2020 – 05:20 am BST

PDB ID : 1FSK
Title : COMPLEX FORMATION BETWEEN A FAB FRAGMENT OF A MONOCLONAL IGG ANTIBODY AND THE MAJOR ALLERGEN FROM BIRCH POLLEN BET V 1
Authors : Mirza, O.; Henriksen, A.; Ipsen, H.; Larsen, J.; Wissenbach, M.; Spangfort, M.; Gajhede, M.
Deposited on : 2000-09-11
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	159	 47% 46% 7%
1	D	159	 48% 45% 7%
1	G	159	 47% 47% 7% 4%
1	J	159	 48% 45% 8% 11%
2	B	214	 37% 54% 9% 3%
2	E	214	 40% 51% 9%
2	H	214	 38% 52% 10% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	K	214	 2% 38% 52% 10%
3	C	220	 4% 45% 46% 9%
3	F	220	 3% 43% 48% 9%
3	I	220	 4% 42% 49% 9%
3	L	220	 4% 43% 48% 9%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 18308 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MAJOR POLLEN ALLERGEN BET V 1-A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	159	1230	783	202	244	1	0	0	0
1	D	159	1230	783	202	244	1	0	0	0
1	G	159	1230	783	202	244	1	0	0	0
1	J	159	1230	783	202	244	1	0	0	0

- Molecule 2 is a protein called IMMUNOGLOBULIN KAPPA LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	214	1668	1040	277	344	7	0	0	0
2	E	214	1668	1040	277	344	7	0	0	0
2	H	214	1668	1040	277	344	7	0	0	0
2	K	214	1668	1040	277	344	7	0	0	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	4	LEU	MET	CONFLICT	UNP P01837
B	13	VAL	MET	CONFLICT	UNP P01837
B	22	SER	THR	CONFLICT	UNP P01837
B	30	ASP	VAL	CONFLICT	UNP P01837
B	34	PHE	SER	CONFLICT	UNP P01837
B	36	PHE	TYR	CONFLICT	UNP P01837
B	41	ASP	GLU	CONFLICT	UNP P01837
B	48	LEU	ILE	CONFLICT	UNP P01837

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	51	PRO	ALA	CONFLICT	UNP P01837
B	68	THR	ALA	CONFLICT	UNP P01837
B	91	SER	GLY	CONFLICT	UNP P01837
B	149	LYS	ASN	CONFLICT	UNP P01837
E	4	LEU	MET	CONFLICT	UNP P01837
E	13	VAL	MET	CONFLICT	UNP P01837
E	22	SER	THR	CONFLICT	UNP P01837
E	30	ASP	VAL	CONFLICT	UNP P01837
E	34	PHE	SER	CONFLICT	UNP P01837
E	36	PHE	TYR	CONFLICT	UNP P01837
E	41	ASP	GLU	CONFLICT	UNP P01837
E	48	LEU	ILE	CONFLICT	UNP P01837
E	51	PRO	ALA	CONFLICT	UNP P01837
E	68	THR	ALA	CONFLICT	UNP P01837
E	91	SER	GLY	CONFLICT	UNP P01837
E	149	LYS	ASN	CONFLICT	UNP P01837
H	4	LEU	MET	CONFLICT	UNP P01837
H	13	VAL	MET	CONFLICT	UNP P01837
H	22	SER	THR	CONFLICT	UNP P01837
H	30	ASP	VAL	CONFLICT	UNP P01837
H	34	PHE	SER	CONFLICT	UNP P01837
H	36	PHE	TYR	CONFLICT	UNP P01837
H	41	ASP	GLU	CONFLICT	UNP P01837
H	48	LEU	ILE	CONFLICT	UNP P01837
H	51	PRO	ALA	CONFLICT	UNP P01837
H	68	THR	ALA	CONFLICT	UNP P01837
H	91	SER	GLY	CONFLICT	UNP P01837
H	149	LYS	ASN	CONFLICT	UNP P01837
K	4	LEU	MET	CONFLICT	UNP P01837
K	13	VAL	MET	CONFLICT	UNP P01837
K	22	SER	THR	CONFLICT	UNP P01837
K	30	ASP	VAL	CONFLICT	UNP P01837
K	34	PHE	SER	CONFLICT	UNP P01837
K	36	PHE	TYR	CONFLICT	UNP P01837
K	41	ASP	GLU	CONFLICT	UNP P01837
K	48	LEU	ILE	CONFLICT	UNP P01837
K	51	PRO	ALA	CONFLICT	UNP P01837
K	68	THR	ALA	CONFLICT	UNP P01837
K	91	SER	GLY	CONFLICT	UNP P01837
K	149	LYS	ASN	CONFLICT	UNP P01837


- Molecule 3 is a protein called ANTIBODY HEAVY CHAIN FAB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	220	Total 1679	C 1066	N 276	O 330	S 7	0	0	0
3	F	220	Total 1679	C 1066	N 276	O 330	S 7	0	0	0
3	I	220	Total 1679	C 1066	N 276	O 330	S 7	0	0	0
3	L	220	Total 1679	C 1066	N 276	O 330	S 7	0	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MAJOR POLLEN ALLERGEN BET V 1-A



- Molecule 1: MAJOR POLLEN ALLERGEN BET V 1-A




- Molecule 1: MAJOR POLLEN ALLERGEN BET V 1-A




- Molecule 1: MAJOR POLLEN ALLERGEN BET V 1-A






• Molecule 2: IMMUNOGLOBULIN KAPPA LIGHT CHAIN




• Molecule 2: IMMUNOGLOBULIN KAPPA LIGHT CHAIN




• Molecule 2: IMMUNOGLOBULIN KAPPA LIGHT CHAIN







• Molecule 2: IMMUNOGLOBULIN KAPPA LIGHT CHAIN



• Molecule 3: ANTIBODY HEAVY CHAIN FAB




• Molecule 3: ANTIBODY HEAVY CHAIN FAB






- Molecule 3: ANTIBODY HEAVY CHAIN FAB



- Molecule 3: ANTIBODY HEAVY CHAIN FAB



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	91.65Å 99.14Å 108.91Å 105.70° 98.32° 97.62°	Depositor
Resolution (Å)	20.00 – 2.90 20.03 – 2.89	Depositor EDS
% Data completeness (in resolution range)	92.4 (20.00-2.90) 82.3 (20.03-2.89)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.48 (at 2.88Å)	Xtrriage
Refinement program	CNS 0.5	Depositor
R, R_{free}	0.253 , 0.285 0.267 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	53.1	Xtrriage
Anisotropy	0.271	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 78.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	18308	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 31.16 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1592e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/1255	0.71	0/1696
1	D	0.51	0/1255	0.71	0/1696
1	G	0.51	0/1255	0.71	0/1696
1	J	0.51	0/1255	0.71	0/1696
2	B	0.51	0/1707	0.78	1/2319 (0.0%)
2	E	0.51	0/1707	0.78	1/2319 (0.0%)
2	H	0.51	0/1707	0.78	1/2319 (0.0%)
2	K	0.52	0/1707	0.78	1/2319 (0.0%)
3	C	0.54	0/1727	0.77	1/2368 (0.0%)
3	F	0.54	0/1727	0.77	1/2368 (0.0%)
3	I	0.54	0/1727	0.77	1/2368 (0.0%)
3	L	0.54	0/1727	0.77	1/2368 (0.0%)
All	All	0.52	0/18756	0.76	8/25532 (0.0%)

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	K	94	TYR	N-CA-C	-6.08	94.59	111.00
2	E	94	TYR	N-CA-C	-5.92	95.02	111.00
2	H	94	TYR	N-CA-C	-5.85	95.20	111.00
2	B	94	TYR	N-CA-C	-5.84	95.23	111.00
3	I	2	VAL	N-CA-C	-5.11	97.20	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1230	0	1221	74	0
1	D	1230	0	1221	74	0
1	G	1230	0	1221	80	0
1	J	1230	0	1221	81	0
2	B	1668	0	1592	155	0
2	E	1668	0	1592	156	0
2	H	1668	0	1592	154	0
2	K	1668	0	1592	154	0
3	C	1679	0	1635	114	0
3	F	1679	0	1635	114	0
3	I	1679	0	1635	112	0
3	L	1679	0	1635	125	0
All	All	18308	0	17792	1344	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 38.

The worst 5 of 1344 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:49:TYR:HB2	3:C:103:THR:HG21	1.26	1.16
2:E:49:TYR:HB2	3:F:103:THR:HG21	1.28	1.09
2:K:49:TYR:HB2	3:L:103:THR:HG21	1.38	1.05
2:H:49:TYR:HB2	3:I:103:THR:HG21	1.34	1.04
2:B:49:TYR:HB2	3:C:103:THR:CG2	1.92	0.99

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	157/159 (99%)	120 (76%)	28 (18%)	9 (6%)	1	5
1	D	157/159 (99%)	120 (76%)	28 (18%)	9 (6%)	1	5
1	G	157/159 (99%)	120 (76%)	28 (18%)	9 (6%)	1	5
1	J	157/159 (99%)	120 (76%)	28 (18%)	9 (6%)	1	5
2	B	212/214 (99%)	177 (84%)	29 (14%)	6 (3%)	5	19
2	E	212/214 (99%)	177 (84%)	29 (14%)	6 (3%)	5	19
2	H	212/214 (99%)	176 (83%)	30 (14%)	6 (3%)	5	19
2	K	212/214 (99%)	176 (83%)	30 (14%)	6 (3%)	5	19
3	C	218/220 (99%)	192 (88%)	20 (9%)	6 (3%)	5	19
3	F	218/220 (99%)	192 (88%)	20 (9%)	6 (3%)	5	19
3	I	218/220 (99%)	193 (88%)	19 (9%)	6 (3%)	5	19
3	L	218/220 (99%)	192 (88%)	20 (9%)	6 (3%)	5	19
All	All	2348/2372 (99%)	1955 (83%)	309 (13%)	84 (4%)	3	14

5 of 84 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	2	VAL
3	F	2	VAL
3	I	2	VAL
3	L	2	VAL
1	A	130	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/134 (100%)	122 (91%)	12 (9%)	9	29
1	D	134/134 (100%)	122 (91%)	12 (9%)	9	29
1	G	134/134 (100%)	122 (91%)	12 (9%)	9	29
1	J	134/134 (100%)	121 (90%)	13 (10%)	8	25

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	193/193 (100%)	171 (89%)	22 (11%)	5	17
2	E	193/193 (100%)	171 (89%)	22 (11%)	5	17
2	H	193/193 (100%)	171 (89%)	22 (11%)	5	17
2	K	193/193 (100%)	171 (89%)	22 (11%)	5	17
3	C	192/192 (100%)	163 (85%)	29 (15%)	3	9
3	F	192/192 (100%)	162 (84%)	30 (16%)	2	8
3	I	192/192 (100%)	163 (85%)	29 (15%)	3	9
3	L	192/192 (100%)	162 (84%)	30 (16%)	2	8
All	All	2076/2076 (100%)	1821 (88%)	255 (12%)	4	14

5 of 255 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	F	146	LEU
2	H	48	LEU
3	L	82	GLN
3	F	155	VAL
1	G	47	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 67 such sidechains are listed below:

Mol	Chain	Res	Type
3	F	169	HIS
1	G	159	ASN
3	L	50	ASN
3	F	201	ASN
1	G	118	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	159/159 (100%)	-0.20	1 (0%) 89 89	22, 56, 89, 124	0
1	D	159/159 (100%)	-0.15	1 (0%) 89 89	18, 55, 93, 109	0
1	G	159/159 (100%)	0.27	6 (3%) 40 36	27, 76, 108, 128	0
1	J	159/159 (100%)	0.70	17 (10%) 6 4	27, 89, 116, 133	0
2	B	214/214 (100%)	-0.11	7 (3%) 46 41	19, 52, 92, 137	0
2	E	214/214 (100%)	-0.02	3 (1%) 75 75	19, 56, 93, 132	0
2	H	214/214 (100%)	0.76	23 (10%) 6 4	41, 86, 119, 153	0
2	K	214/214 (100%)	0.07	5 (2%) 60 58	32, 62, 91, 132	0
3	C	220/220 (100%)	-0.25	8 (3%) 42 37	18, 44, 84, 123	0
3	F	220/220 (100%)	-0.19	7 (3%) 47 43	14, 49, 86, 121	0
3	I	220/220 (100%)	0.04	9 (4%) 37 32	33, 60, 106, 134	0
3	L	220/220 (100%)	-0.01	8 (3%) 42 37	25, 56, 93, 123	0
All	All	2372/2372 (100%)	0.07	95 (4%) 38 33	14, 59, 105, 153	0

The worst 5 of 95 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	214	CYS	7.9
2	H	212	ASN	6.5
3	F	220	CYS	5.6
3	L	220	CYS	5.6
3	I	134	ALA	5.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.