



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 16, 2023 – 02:55 AM EDT

PDB ID : 2FTY
Title : Crystal structure of dihydropyrimidinase from *Saccharomyces kluyveri*
Authors : Dobritsch, D.; Lohkamp, B.
Deposited on : 2006-01-25
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	559	 84% 11% 5%
1	B	559	 82% 12% 5%
1	C	559	 85% 9% 5%
1	D	559	 82% 12% 5%

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 17311 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called dihydropyrimidinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	532	4164	2646	680	811	27	0	2	0
1	B	532	4169	2651	680	811	27	0	3	0
1	C	531	4142	2630	677	809	26	0	0	0
1	D	530	4147	2633	678	809	27	0	2	0

There are 76 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	167	KCX	LYS	modified residue	UNP Q9P903
A	543	PRO	-	expression tag	UNP Q9P903
A	544	GLY	-	expression tag	UNP Q9P903
A	545	ASP	-	expression tag	UNP Q9P903
A	546	ASP	-	expression tag	UNP Q9P903
A	547	ASP	-	expression tag	UNP Q9P903
A	548	ASP	-	expression tag	UNP Q9P903
A	549	LYS	-	expression tag	UNP Q9P903
A	550	HIS	-	expression tag	UNP Q9P903
A	551	HIS	-	expression tag	UNP Q9P903
A	552	HIS	-	expression tag	UNP Q9P903
A	553	HIS	-	expression tag	UNP Q9P903
A	554	HIS	-	expression tag	UNP Q9P903
A	555	HIS	-	expression tag	UNP Q9P903
A	556	HIS	-	expression tag	UNP Q9P903
A	557	HIS	-	expression tag	UNP Q9P903
A	558	SER	-	expression tag	UNP Q9P903
A	559	GLY	-	expression tag	UNP Q9P903
A	560	ASP	-	expression tag	UNP Q9P903
B	167	KCX	LYS	modified residue	UNP Q9P903
B	543	PRO	-	expression tag	UNP Q9P903

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	544	GLY	-	expression tag	UNP Q9P903
B	545	ASP	-	expression tag	UNP Q9P903
B	546	ASP	-	expression tag	UNP Q9P903
B	547	ASP	-	expression tag	UNP Q9P903
B	548	ASP	-	expression tag	UNP Q9P903
B	549	LYS	-	expression tag	UNP Q9P903
B	550	HIS	-	expression tag	UNP Q9P903
B	551	HIS	-	expression tag	UNP Q9P903
B	552	HIS	-	expression tag	UNP Q9P903
B	553	HIS	-	expression tag	UNP Q9P903
B	554	HIS	-	expression tag	UNP Q9P903
B	555	HIS	-	expression tag	UNP Q9P903
B	556	HIS	-	expression tag	UNP Q9P903
B	557	HIS	-	expression tag	UNP Q9P903
B	558	SER	-	expression tag	UNP Q9P903
B	559	GLY	-	expression tag	UNP Q9P903
B	560	ASP	-	expression tag	UNP Q9P903
C	167	KCX	LYS	modified residue	UNP Q9P903
C	543	PRO	-	expression tag	UNP Q9P903
C	544	GLY	-	expression tag	UNP Q9P903
C	545	ASP	-	expression tag	UNP Q9P903
C	546	ASP	-	expression tag	UNP Q9P903
C	547	ASP	-	expression tag	UNP Q9P903
C	548	ASP	-	expression tag	UNP Q9P903
C	549	LYS	-	expression tag	UNP Q9P903
C	550	HIS	-	expression tag	UNP Q9P903
C	551	HIS	-	expression tag	UNP Q9P903
C	552	HIS	-	expression tag	UNP Q9P903
C	553	HIS	-	expression tag	UNP Q9P903
C	554	HIS	-	expression tag	UNP Q9P903
C	555	HIS	-	expression tag	UNP Q9P903
C	556	HIS	-	expression tag	UNP Q9P903
C	557	HIS	-	expression tag	UNP Q9P903
C	558	SER	-	expression tag	UNP Q9P903
C	559	GLY	-	expression tag	UNP Q9P903
C	560	ASP	-	expression tag	UNP Q9P903
D	167	KCX	LYS	modified residue	UNP Q9P903
D	543	PRO	-	expression tag	UNP Q9P903
D	544	GLY	-	expression tag	UNP Q9P903
D	545	ASP	-	expression tag	UNP Q9P903
D	546	ASP	-	expression tag	UNP Q9P903
D	547	ASP	-	expression tag	UNP Q9P903

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	548	ASP	-	expression tag	UNP Q9P903
D	549	LYS	-	expression tag	UNP Q9P903
D	550	HIS	-	expression tag	UNP Q9P903
D	551	HIS	-	expression tag	UNP Q9P903
D	552	HIS	-	expression tag	UNP Q9P903
D	553	HIS	-	expression tag	UNP Q9P903
D	554	HIS	-	expression tag	UNP Q9P903
D	555	HIS	-	expression tag	UNP Q9P903
D	556	HIS	-	expression tag	UNP Q9P903
D	557	HIS	-	expression tag	UNP Q9P903
D	558	SER	-	expression tag	UNP Q9P903
D	559	GLY	-	expression tag	UNP Q9P903
D	560	ASP	-	expression tag	UNP Q9P903

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total 2	Zn 2	0	0
2	B	2	Total 2	Zn 2	0	0
2	C	2	Total 2	Zn 2	0	0
2	D	2	Total 2	Zn 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	189	Total 189	O 189	0	0
3	B	188	Total 188	O 188	0	0
3	C	171	Total 171	O 171	0	0
3	D	133	Total 133	O 133	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: dihydropyrimidinase




- Molecule 1: dihydropyrimidinase





- Molecule 1: dihydropyrimidinase





• Molecule 1: dihydropyrimidinase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	91.43Å 73.32Å 162.18Å 90.00° 91.59° 90.00°	Depositor
Resolution (Å)	43.85 – 2.40 43.50 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.0 (43.85-2.40) 99.0 (43.50-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.10 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.217 , 0.274 0.222 , 0.216	Depositor DCC
R_{free} test set	4200 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	35.0	Xtrriage
Anisotropy	0.598	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 27.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.126 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	17311	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/4252	0.59	0/5771
1	B	0.47	0/4260	0.60	0/5782
1	C	0.47	0/4222	0.59	0/5729
1	D	0.45	0/4234	0.57	0/5745
All	All	0.47	0/16968	0.59	0/23027

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4164	0	4093	30	0
1	B	4169	0	4104	38	0
1	C	4142	0	4072	31	0
1	D	4147	0	4077	37	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	189	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	188	0	0	0	0
3	C	171	0	0	3	0
3	D	133	0	0	3	0
All	All	17311	0	16346	132	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 132 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:260:GLN:HE21	1:C:260:GLN:H	1.14	0.92
1:C:140:GLU:O	1:C:146:ALA:HB2	1.78	0.83
1:D:140:GLU:O	1:D:146:ALA:HB2	1.77	0.83
1:A:198:LEU:HD12	1:A:251:ILE:HD11	1.60	0.83
1:B:198:LEU:HD12	1:B:251[A]:ILE:HD11	1.63	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	529/559 (95%)	510 (96%)	15 (3%)	4 (1%)	19 29
1	B	530/559 (95%)	512 (97%)	16 (3%)	2 (0%)	34 48
1	C	526/559 (94%)	506 (96%)	18 (3%)	2 (0%)	34 48
1	D	527/559 (94%)	505 (96%)	19 (4%)	3 (1%)	25 36
All	All	2112/2236 (94%)	2033 (96%)	68 (3%)	11 (0%)	34 41

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	144	VAL
1	A	172[A]	TYR
1	A	172[B]	TYR
1	A	255	HIS
1	B	172[A]	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	457/479 (95%)	443 (97%)	14 (3%)	40 60
1	B	458/479 (96%)	442 (96%)	16 (4%)	36 55
1	C	454/479 (95%)	439 (97%)	15 (3%)	38 57
1	D	456/479 (95%)	442 (97%)	14 (3%)	40 60
All	All	1825/1916 (95%)	1766 (97%)	59 (3%)	39 59

5 of 59 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	476	HIS
1	D	404	LEU
1	C	320	ASP
1	D	388	ARG
1	D	320	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	27	ASN
1	C	202	ASN
1	D	340	GLN
1	C	260	GLN
1	A	318	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	KCX	A	167	1,2	9,11,12	0.93	0	5,12,14	1.48	1 (20%)
1	KCX	C	167	1,2	9,11,12	0.81	0	5,12,14	1.31	1 (20%)
1	KCX	D	167	1,2	9,11,12	0.90	0	5,12,14	1.81	1 (20%)
1	KCX	B	167	1,2	9,11,12	0.93	0	5,12,14	1.92	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	A	167	1,2	-	0/9/10/12	-
1	KCX	C	167	1,2	-	1/9/10/12	-
1	KCX	D	167	1,2	-	0/9/10/12	-
1	KCX	B	167	1,2	-	1/9/10/12	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	167	KCX	OQ1-CX-NZ	-3.85	118.99	124.96
1	B	167	KCX	OQ1-CX-NZ	-3.58	119.40	124.96
1	A	167	KCX	OQ1-CX-NZ	-3.17	120.04	124.96
1	C	167	KCX	OQ1-CX-NZ	-2.84	120.56	124.96
1	B	167	KCX	CE-NZ-CX	-2.17	118.40	121.89

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	167	KCX	C-CA-CB-CG
1	C	167	KCX	C-CA-CB-CG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	531/559 (94%)	-0.21	3 (0%) 89 88	31, 36, 38, 45	0
1	B	531/559 (94%)	-0.20	5 (0%) 84 82	31, 36, 38, 42	0
1	C	530/559 (94%)	-0.13	7 (1%) 77 75	31, 36, 38, 42	0
1	D	529/559 (94%)	-0.13	6 (1%) 80 79	31, 36, 38, 42	0
All	All	2121/2236 (94%)	-0.17	21 (0%) 82 80	31, 36, 38, 45	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	146	ALA	7.2
1	B	294	HIS	3.8
1	C	304	VAL	3.6
1	D	382	ASN	3.2
1	C	457	ASP	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	D	167	12/13	0.90	0.16	35,36,44,44	3
1	KCX	A	167	12/13	0.91	0.14	35,36,44,45	3
1	KCX	B	167	12/13	0.94	0.12	35,35,44,44	3
1	KCX	C	167	12/13	0.95	0.17	35,36,44,44	3

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	D	602	1/1	0.95	0.21	46,46,46,46	1
2	ZN	D	601	1/1	0.96	0.07	44,44,44,44	1
2	ZN	B	602	1/1	0.97	0.17	46,46,46,46	1
2	ZN	C	602	1/1	0.97	0.11	45,45,45,45	1
2	ZN	C	601	1/1	0.98	0.06	44,44,44,44	1
2	ZN	B	601	1/1	0.99	0.08	43,43,43,43	1
2	ZN	A	601	1/1	0.99	0.08	45,45,45,45	1
2	ZN	A	602	1/1	0.99	0.15	45,45,45,45	1

6.5 Other polymers [i](#)

There are no such residues in this entry.