



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 3, 2024 – 07:00 pm GMT

PDB ID : 5FUK
Title : Crystallization of a dimeric heme peroxxygenase from the fungus *Marasmius rotula*
Authors : Piontek, K.; Strittmatter, E.; Plattner, D.A.
Deposited on : 2016-01-27
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1483 (1.56-1.56)
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495 (1.56-1.56)
RSRZ outliers	127900	1465 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	234	 5% 94% 6%
1	B	234	 3% 94% 6%
2	C	5	 100%
3	D	6	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MAN	D	5	-	-	-	X
6	PLM	A	1237[B]	-	-	X	-
6	PLM	B	1237[B]	-	-	X	-
9	NAG	A	1245	-	-	-	X

2 Entry composition i


There are 11 unique types of molecules in this entry. The entry contains 4624 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MROUPO.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	234	Total 1798	C 1130	N 317	O 345	S 6	0	2	0
1	B	234	Total 1817	C 1142	N 322	O 347	S 6	0	5	0

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	5	Total 61	C 34	N 2	O 25	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	6	Total 72	C 40	N 2	O 30	0	0	0

- Molecule 4 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 5 is 1-(ISOPROPYLAMINO)-3-(1-NAPHTHYLOXY)-2-PROPANOL (three-letter code: SNP) (formula: $C_{16}H_{21}NO_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
5	A	1	Total	C	N	O	0	1
			19	16	1	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	B	1	19	16	1	2	0	1

- Molecule 6 is PALMITIC ACID (three-letter code: PLM) (formula: $C_{16}H_{32}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	A	1	18	16	2	0	1
6	B	1	18	16	2	0	1

- Molecule 7 is (1E,2R)-1-(ISOPROPYLIMINO)-3-(1-NAPHTHYLOXY)PROPAN-2-OL (three-letter code: RNP) (formula: $C_{16}H_{19}NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	1
			19	16	1	2		
7	B	1	Total	C	N	O	0	1
			19	16	1	2		

- Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Mg	0	0
			1	1		
8	B	1	Total	Mg	0	0
			1	1		

- Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
9	A	1	14	8	1	5	0	0
9	B	1	14	8	1	5	0	0

- Molecule 10 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
			Total	O			P
10	B	1	5	4	1	0	0


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	333	Total 333	O 333	0	0
11	B	310	Total 310	O 310	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MROUPO



- Molecule 1: MROUPO



- Molecule 2: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.01Å 76.67Å 120.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.36 – 1.55 29.12 – 1.55	Depositor EDS
% Data completeness (in resolution range)	98.3 (32.36-1.55) 98.3 (29.12-1.55)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 1.55Å)	Xtrriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.155 , 0.183 0.168 , 0.195	Depositor DCC
R_{free} test set	3999 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	18.8	Xtrriage
Anisotropy	0.056	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 52.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4624	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, PLM, MG, NAG, PO4, HEM, MAN, RNP, SNP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.98	1/1842 (0.1%)	0.97	7/2507 (0.3%)
1	B	1.09	1/1869 (0.1%)	1.05	10/2542 (0.4%)
All	All	1.04	2/3711 (0.1%)	1.01	17/5049 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	124	SER	CB-OG	-5.86	1.34	1.42
1	B	105	GLU	CG-CD	5.18	1.59	1.51

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	213	ARG	NE-CZ-NH1	8.15	124.38	120.30
1	B	213	ARG	NE-CZ-NH2	-7.83	116.38	120.30
1	A	99	ASP	CB-CG-OD1	7.15	124.73	118.30
1	B	213	ARG	CG-CD-NE	-6.90	97.31	111.80
1	B	133	ARG	NE-CZ-NH2	-6.63	116.99	120.30
1	B	99	ASP	CB-CG-OD1	6.28	123.95	118.30
1	A	71	ASP	CB-CG-OD1	5.80	123.52	118.30
1	A	132	ASP	CB-CG-OD2	-5.54	113.32	118.30
1	A	154	ARG	NE-CZ-NH2	-5.47	117.56	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	14	ARG	NE-CZ-NH1	5.47	123.03	120.30
1	A	136	ASP	CB-CG-OD1	-5.45	113.39	118.30
1	A	54	ASP	CB-CG-OD1	5.42	123.18	118.30
1	B	160[A]	PHE	CB-CG-CD2	-5.28	117.10	120.80
1	B	160[B]	PHE	CB-CG-CD2	-5.28	117.10	120.80
1	B	225	ASP	CB-CA-C	-5.08	100.25	110.40
1	B	154	ARG	NE-CZ-NH1	5.07	122.83	120.30
1	A	133	ARG	NE-CZ-NH2	-5.04	117.78	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	1	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1798	0	1777	6	0
1	B	1817	0	1809	12	0
2	C	61	0	52	0	0
3	D	72	0	61	0	0
4	A	43	0	30	0	0
4	B	43	0	30	0	0
5	A	19	0	21	4	0
5	B	19	0	21	0	0
6	A	18	0	31	9	0
6	B	18	0	31	9	0
7	A	19	0	19	1	0
7	B	19	0	19	1	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
9	A	14	0	13	0	0
9	B	14	0	13	0	0
10	B	5	0	0	0	0
11	A	333	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
11	B	310	0	0	5	0
All	All	4624	0	3927	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:160[B]:PHE:CE2	6:B:1237[B]:PLM:H22	1.41	1.52
1:B:160[B]:PHE:HE2	6:B:1237[B]:PLM:C2	1.46	1.28
1:B:160[B]:PHE:CE2	6:B:1237[B]:PLM:C2	2.24	1.16
1:A:160[B]:PHE:HE2	6:A:1237[B]:PLM:H21	1.16	1.08
1:A:160[B]:PHE:CE2	6:A:1237[B]:PLM:H21	1.87	1.08
6:A:1237[B]:PLM:HD1	6:A:1237[B]:PLM:H82	1.34	1.06
5:A:1236[A]:SNP:O2	5:A:1236[A]:SNP:H161	1.71	0.89
7:B:1238[C]:RNP:H161	11:B:2269:HOH:O	1.73	0.88
1:A:160[B]:PHE:HE2	6:A:1237[B]:PLM:C2	1.86	0.88
6:A:1237[B]:PLM:H82	6:A:1237[B]:PLM:CD	2.14	0.75
6:A:1237[B]:PLM:HD1	6:A:1237[B]:PLM:C8	2.18	0.71
1:A:130:GLN:HE22	1:A:155:SER:HA	1.62	0.64
1:B:11:ASN:C	11:B:2014:HOH:O	2.36	0.64
6:A:1237[B]:PLM:CD	6:A:1237[B]:PLM:C8	2.78	0.62
1:B:160[B]:PHE:HE2	6:B:1237[B]:PLM:H22	0.56	0.61
1:B:160[B]:PHE:CE2	6:B:1237[B]:PLM:H21	2.33	0.60
1:B:160[B]:PHE:CZ	6:B:1237[B]:PLM:H22	2.25	0.60
1:A:160[B]:PHE:CZ	6:A:1237[B]:PLM:H21	2.37	0.58
1:B:128[B]:GLN:OE1	11:B:2174:HOH:O	2.18	0.56
1:A:10[A]:PRO:HD2	11:A:2015:HOH:O	2.05	0.56
1:B:178:LYS:HE2	11:B:2236:HOH:O	2.08	0.54
1:B:12:ASP:N	11:B:2014:HOH:O	2.43	0.52
5:A:1236[A]:SNP:O2	5:A:1236[A]:SNP:C16	2.52	0.49
6:A:1237[B]:PLM:H61	6:A:1237[B]:PLM:HE2	1.95	0.48
11:A:2322:HOH:O	6:B:1237[B]:PLM:HA2	2.14	0.47
6:B:1237[B]:PLM:HB2	6:B:1237[B]:PLM:HE2	1.73	0.46
5:A:1236[A]:SNP:H161	5:A:1236[A]:SNP:H2O1	1.79	0.44
1:B:160[B]:PHE:CZ	6:B:1237[B]:PLM:C2	2.95	0.44
1:B:138:LEU:HD23	1:B:144:VAL:HG11	2.00	0.43
7:A:1238[C]:RNP:H162	7:A:1238[C]:RNP:H13	1.73	0.41
5:A:1236[A]:SNP:O2	11:A:2323:HOH:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	234/234 (100%)	229 (98%)	5 (2%)	0	100	100
1	B	237/234 (101%)	233 (98%)	4 (2%)	0	100	100
All	All	471/468 (101%)	462 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/196 (101%)	194 (98%)	4 (2%)	55	26
1	B	201/196 (103%)	199 (99%)	2 (1%)	76	57
All	All	399/392 (102%)	393 (98%)	6 (2%)	76	37

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	SER
1	A	160[A]	PHE
1	A	160[B]	PHE
1	A	161	PHE
1	B	160[A]	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	160[B]	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	ASN
1	A	130	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

11 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	1,2	14,14,15	1.72	2 (14%)	17,19,21	1.26	1 (5%)
2	NAG	C	2	2	14,14,15	0.80	0	17,19,21	1.66	2 (11%)
2	BMA	C	3	2	11,11,12	0.62	0	15,15,17	1.44	3 (20%)
2	MAN	C	4	2	11,11,12	0.89	0	15,15,17	1.91	4 (26%)
2	MAN	C	5	2	11,11,12	0.78	0	15,15,17	1.01	1 (6%)
3	NAG	D	1	1,3	14,14,15	1.66	3 (21%)	17,19,21	1.90	6 (35%)
3	NAG	D	2	3	14,14,15	0.71	0	17,19,21	1.59	3 (17%)
3	BMA	D	3	3	11,11,12	0.75	0	15,15,17	1.29	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MAN	D	4	3	11,11,12	0.98	1 (9%)	15,15,17	1.54	4 (26%)
3	MAN	D	5	3	11,11,12	0.63	0	15,15,17	1.37	2 (13%)
3	MAN	D	6	3	11,11,12	0.89	0	15,15,17	1.08	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	0/2/19/22	0/1/1/1
2	MAN	C	5	2	-	0/2/19/22	0/1/1/1
3	NAG	D	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	0/2/19/22	0/1/1/1
3	MAN	D	5	3	-	2/2/19/22	1/1/1/1
3	MAN	D	6	3	-	0/2/19/22	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	NAG	C1-C2	5.16	1.60	1.52
3	D	1	NAG	C1-C2	4.25	1.58	1.52
3	D	1	NAG	O5-C1	2.88	1.48	1.43
2	C	1	NAG	O5-C1	-2.72	1.39	1.43
3	D	4	MAN	C4-C5	2.44	1.58	1.53
3	D	1	NAG	O5-C5	-2.24	1.38	1.43

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	4	MAN	C1-O5-C5	4.96	118.92	112.19
3	D	1	NAG	C1-O5-C5	-3.96	106.83	112.19
2	C	3	BMA	O2-C2-C1	-3.56	101.86	109.15
3	D	1	NAG	C3-C4-C5	-3.36	104.25	110.24
2	C	2	NAG	O6-C6-C5	-3.33	99.88	111.29
2	C	2	NAG	C2-N2-C7	-3.27	118.25	122.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1	NAG	O5-C1-C2	-3.07	106.43	111.29
3	D	5	MAN	O5-C5-C6	2.99	111.89	107.20
3	D	1	NAG	C6-C5-C4	-2.91	106.18	113.00
2	C	4	MAN	O4-C4-C3	-2.79	103.89	110.35
2	C	5	MAN	C1-O5-C5	2.76	115.93	112.19
2	C	1	NAG	C4-C3-C2	-2.76	106.97	111.02
3	D	2	NAG	O3-C3-C2	-2.75	103.78	109.47
3	D	4	MAN	O3-C3-C4	2.74	116.69	110.35
3	D	4	MAN	C6-C5-C4	2.65	119.21	113.00
2	C	4	MAN	O5-C5-C6	2.63	111.32	107.20
2	C	3	BMA	O3-C3-C4	-2.55	104.44	110.35
3	D	4	MAN	C1-O5-C5	2.52	115.61	112.19
3	D	1	NAG	O4-C4-C3	-2.50	104.58	110.35
3	D	1	NAG	O5-C5-C6	-2.47	103.34	107.20
3	D	5	MAN	C2-C3-C4	-2.42	106.71	110.89
3	D	2	NAG	O5-C5-C4	-2.25	105.36	110.83
3	D	6	MAN	C1-O5-C5	2.22	115.20	112.19
3	D	4	MAN	C1-C2-C3	2.20	112.37	109.67
3	D	3	BMA	C1-O5-C5	-2.16	109.27	112.19
2	C	3	BMA	O5-C5-C4	-2.06	105.82	110.83
2	C	4	MAN	O4-C4-C5	-2.02	104.27	109.30
3	D	2	NAG	C6-C5-C4	-2.02	108.27	113.00
3	D	3	BMA	O3-C3-C4	-2.01	105.71	110.35

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	5	MAN	C4-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
3	D	5	MAN	O5-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6


All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	5	MAN	C1-C2-C3-C4-C5-O5

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 2 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SNP	A	1236[A]	-	20,20,20	1.11	2 (10%)	26,26,26	0.93	0
6	PLM	B	1237[B]	-	17,17,17	0.63	0	17,17,17	0.97	2 (11%)
10	PO4	B	1247	-	4,4,4	1.34	1 (25%)	6,6,6	1.10	0
4	HEM	B	1235	1,8	41,50,50	1.36	6 (14%)	45,82,82	1.68	7 (15%)
9	NAG	B	1240	1	14,14,15	0.96	0	17,19,21	1.46	2 (11%)
4	HEM	A	1235	1,8	41,50,50	1.48	7 (17%)	45,82,82	1.81	11 (24%)
7	RNP	A	1238[C]	-	19,20,20	1.61	5 (26%)	22,26,26	1.33	1 (4%)
5	SNP	B	1236[A]	-	20,20,20	1.24	2 (10%)	26,26,26	0.92	2 (7%)
6	PLM	A	1237[B]	-	17,17,17	0.64	1 (5%)	17,17,17	1.41	2 (11%)
7	RNP	B	1238[C]	-	19,20,20	1.43	3 (15%)	22,26,26	1.65	6 (27%)
9	NAG	A	1245	1	14,14,15	0.56	0	17,19,21	1.98	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	SNP	A	1236[A]	-	-	7/10/10/10	0/2/2/2
6	PLM	B	1237[B]	-	-	11/15/15/15	-
4	HEM	B	1235	1,8	-	2/12/54/54	-
9	NAG	B	1240	1	-	4/6/23/26	0/1/1/1
4	HEM	A	1235	1,8	-	2/12/54/54	-
7	RNP	A	1238[C]	-	-	3/8/10/10	0/2/2/2
5	SNP	B	1236[A]	-	-	5/10/10/10	0/2/2/2
6	PLM	A	1237[B]	-	-	6/15/15/15	-
7	RNP	B	1238[C]	-	-	6/8/10/10	0/2/2/2
9	NAG	A	1245	1	-	0/6/23/26	0/1/1/1

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	1238[C]	RNP	C12-C13	4.01	1.53	1.49
4	A	1235	HEM	C4D-ND	-3.53	1.34	1.40
4	B	1235	HEM	C1B-NB	-3.47	1.34	1.40
5	B	1236[A]	SNP	C9-C10	3.10	1.48	1.43
4	A	1235	HEM	C3C-C2C	-2.99	1.36	1.40
7	B	1238[C]	RNP	C9-C10	2.80	1.48	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	1238[C]	RNP	C9-C10	2.70	1.47	1.43
7	B	1238[C]	RNP	C1-C9	2.69	1.49	1.42
5	A	1236[A]	SNP	C9-C10	2.60	1.47	1.43
4	B	1235	HEM	O1A-CGA	2.58	1.30	1.22
4	B	1235	HEM	C3C-C2C	-2.58	1.36	1.40
7	B	1238[C]	RNP	C14-N1	-2.57	1.45	1.47
4	A	1235	HEM	C3B-C4B	2.54	1.50	1.44
4	A	1235	HEM	C1B-NB	-2.54	1.35	1.40
4	A	1235	HEM	CMB-C2B	2.47	1.56	1.50
10	B	1247	PO4	P-O1	2.40	1.56	1.50
5	B	1236[A]	SNP	C1-C9	2.39	1.48	1.42
4	A	1235	HEM	C4B-NB	-2.34	1.34	1.38
6	A	1237[B]	PLM	O1-C1	-2.28	1.23	1.30
7	A	1238[C]	RNP	C1-C9	2.27	1.48	1.42
4	A	1235	HEM	CAA-C2A	-2.26	1.48	1.52
4	B	1235	HEM	FE-NB	2.24	2.07	1.96
4	B	1235	HEM	CHA-C4D	2.13	1.40	1.35
4	B	1235	HEM	C1D-C2D	2.12	1.48	1.44
7	A	1238[C]	RNP	C11-C12	2.07	1.54	1.51
5	A	1236[A]	SNP	C1-C9	2.07	1.48	1.42
7	A	1238[C]	RNP	C14-N1	-2.04	1.45	1.47

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	1245	NAG	C1-O5-C5	6.51	121.01	112.19
4	B	1235	HEM	C4B-CHC-C1C	4.96	129.10	122.56
4	B	1235	HEM	C1B-NB-C4B	4.88	110.11	105.07
4	A	1235	HEM	C1B-NB-C4B	4.53	109.75	105.07
7	B	1238[C]	RNP	C16-C14-N1	-4.11	105.02	109.03
7	A	1238[C]	RNP	C15-C14-N1	4.03	112.97	109.03
6	A	1237[B]	PLM	C3-C2-C1	-4.02	104.34	114.47
7	B	1238[C]	RNP	C15-C14-N1	3.59	112.54	109.03
4	A	1235	HEM	CMC-C2C-C3C	3.47	131.18	124.68
4	A	1235	HEM	C2C-C3C-C4C	3.47	109.32	106.90
9	B	1240	NAG	C8-C7-N2	3.29	121.67	116.10
4	A	1235	HEM	CHD-C1D-ND	3.26	127.97	124.43
4	A	1235	HEM	O2A-CGA-CBA	3.26	124.50	114.03
4	B	1235	HEM	O2A-CGA-CBA	3.13	124.08	114.03
4	A	1235	HEM	CMA-C3A-C4A	-3.04	123.80	128.46
9	B	1240	NAG	C4-C3-C2	2.98	115.39	111.02
4	A	1235	HEM	CHC-C4B-NB	2.79	127.46	124.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	1237[B]	PLM	O1-C1-O2	2.63	129.86	123.30
6	A	1237[B]	PLM	O1-C1-O2	2.59	129.77	123.30
4	B	1235	HEM	CAD-CBD-CGD	-2.56	108.10	113.60
4	B	1235	HEM	O1A-CGA-CBA	-2.54	114.91	123.08
4	A	1235	HEM	C4B-CHC-C1C	2.41	125.74	122.56
4	B	1235	HEM	O2D-CGD-CBD	2.37	121.65	114.03
9	A	1245	NAG	C2-N2-C7	-2.37	119.53	122.90
7	B	1238[C]	RNP	C16-C14-C15	2.31	115.54	111.16
4	A	1235	HEM	CHD-C1D-C2D	-2.31	121.37	124.98
4	A	1235	HEM	O2A-CGA-O1A	-2.30	117.56	123.30
5	B	1236[A]	SNP	C13-N1-C14	2.20	121.46	114.44
7	B	1238[C]	RNP	C11-O1-C1	2.19	125.41	118.25
7	B	1238[C]	RNP	O1-C1-C9	2.19	121.12	115.01
4	A	1235	HEM	CMA-C3A-C2A	2.16	129.01	124.94
7	B	1238[C]	RNP	O2-C12-C13	-2.13	104.64	109.20
6	B	1237[B]	PLM	O2-C1-C2	-2.13	116.24	123.08
4	B	1235	HEM	C4B-C3B-C2B	-2.07	105.47	107.11
5	B	1236[A]	SNP	O1-C1-C9	2.05	120.73	115.01

There are no chirality outliers.

All (46) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1236[A]	SNP	C12-C13-N1-C14
5	A	1236[A]	SNP	C15-C14-N1-C13
5	B	1236[A]	SNP	C11-C12-C13-N1
5	B	1236[A]	SNP	O2-C12-C13-N1
5	B	1236[A]	SNP	C12-C13-N1-C14
7	A	1238[C]	RNP	O1-C11-C12-O2
7	A	1238[C]	RNP	O1-C11-C12-C13
7	A	1238[C]	RNP	C16-C14-N1-C13
7	B	1238[C]	RNP	O1-C11-C12-O2
7	B	1238[C]	RNP	O1-C11-C12-C13
7	B	1238[C]	RNP	C16-C14-N1-C13
9	B	1240	NAG	C8-C7-N2-C2
9	B	1240	NAG	O7-C7-N2-C2
6	A	1237[B]	PLM	C1-C2-C3-C4
5	A	1236[A]	SNP	O1-C11-C12-C13
6	B	1237[B]	PLM	C3-C4-C5-C6
6	A	1237[B]	PLM	CC-CD-CE-CF
5	A	1236[A]	SNP	O1-C11-C12-O2
9	B	1240	NAG	O5-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	A	1237[B]	PLM	CA-CB-CC-CD
6	A	1237[B]	PLM	CB-CC-CD-CE
5	A	1236[A]	SNP	C16-C14-N1-C13
5	B	1236[A]	SNP	C16-C14-N1-C13
6	A	1237[B]	PLM	C9-CA-CB-CC
6	B	1237[B]	PLM	C9-CA-CB-CC
6	B	1237[B]	PLM	CB-CC-CD-CE
6	B	1237[B]	PLM	CD-CE-CF-CG
6	B	1237[B]	PLM	C4-C5-C6-C7
7	B	1238[C]	RNP	C9-C1-O1-C11
7	B	1238[C]	RNP	C2-C1-O1-C11
6	B	1237[B]	PLM	CC-CD-CE-CF
5	B	1236[A]	SNP	C15-C14-N1-C13
6	B	1237[B]	PLM	C1-C2-C3-C4
6	B	1237[B]	PLM	CA-CB-CC-CD
5	A	1236[A]	SNP	C9-C1-O1-C11
5	A	1236[A]	SNP	C2-C1-O1-C11
4	B	1235	HEM	CAD-CBD-CGD-O2D
6	B	1237[B]	PLM	O2-C1-C2-C3
4	A	1235	HEM	CAD-CBD-CGD-O2D
6	B	1237[B]	PLM	O1-C1-C2-C3
4	B	1235	HEM	CAD-CBD-CGD-O1D
4	A	1235	HEM	CAD-CBD-CGD-O1D
7	B	1238[C]	RNP	C12-C11-O1-C1
9	B	1240	NAG	C4-C5-C6-O6
6	B	1237[B]	PLM	C6-C7-C8-C9
6	A	1237[B]	PLM	O2-C1-C2-C3


There are no ring outliers.



5 monomers are involved in 24 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1236[A]	SNP	4	0
6	B	1237[B]	PLM	9	0
7	A	1238[C]	RNP	1	0
6	A	1237[B]	PLM	9	0
7	B	1238[C]	RNP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is



within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	234/234 (100%)	0.08	11 (4%) 31 36	15, 20, 36, 70	0
1	B	234/234 (100%)	0.08	7 (2%) 50 58	14, 19, 33, 67	0
All	All	468/468 (100%)	0.08	18 (3%) 40 47	14, 19, 35, 70	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	SER	6.1
1	B	1	SER	5.5
1	B	160[A]	PHE	4.6
1	B	225	ASP	4.5
1	A	225	ASP	4.1
1	A	160[A]	PHE	3.8
1	B	227	CYS	3.6
1	A	10[A]	PRO	3.3
1	A	227	CYS	3.2
1	B	2	ALA	3.1
1	A	2	ALA	3.1
1	B	20	LEU	3.0
1	A	168	GLY	2.9
1	A	11	ASN	2.4
1	B	161	PHE	2.3
1	A	84	ILE	2.2
1	A	90	LEU	2.2
1	A	153	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MAN	C	5	11/12	0.56	0.35	62,66,68,68	0
3	MAN	D	5	11/12	0.56	0.45	83,94,100,101	0
2	MAN	C	4	11/12	0.71	0.25	37,41,46,56	0
3	MAN	D	4	11/12	0.76	0.30	42,51,56,71	0
3	MAN	D	6	11/12	0.85	0.28	42,47,53,57	0
2	BMA	C	3	11/12	0.90	0.21	31,34,43,46	0
2	NAG	C	2	14/15	0.91	0.17	27,32,47,49	0
3	NAG	D	1	14/15	0.92	0.08	16,19,23,25	0
3	NAG	D	2	14/15	0.93	0.18	25,30,35,43	0
3	BMA	D	3	11/12	0.94	0.17	33,37,45,49	0
2	NAG	C	1	14/15	0.95	0.07	19,22,25,27	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	PLM	B	1237[B]	18/18	0.52	0.37	8,24,29,30	18
6	PLM	A	1237[B]	18/18	0.57	0.37	9,29,36,36	18
9	NAG	B	1240	14/15	0.64	0.35	55,68,75,78	0
9	NAG	A	1245	14/15	0.76	0.42	57,67,73,73	0
7	RNP	B	1238[C]	19/19	0.83	0.30	27,30,35,35	19
7	RNP	A	1238[C]	19/19	0.85	0.25	29,35,41,43	19
5	SNP	A	1236[A]	19/19	0.87	0.20	33,36,44,45	19
5	SNP	B	1236[A]	19/19	0.88	0.20	50,52,56,56	19
10	PO4	B	1247	5/5	0.92	0.21	35,35,43,44	0
4	HEM	A	1235	43/43	0.99	0.14	13,14,17,22	0
4	HEM	B	1235	43/43	0.99	0.13	11,13,15,20	0
8	MG	B	1239	1/1	0.99	0.08	15,15,15,15	0
8	MG	A	1239	1/1	1.00	0.10	14,14,14,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around PLM A 1237 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around RNP B 1238 (C):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around RNP A 1238 (C):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around SNP A 1236 (A):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SNP B 1236 (A):


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 1235:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.