

wwPDB X-ray Structure Validation Summary Report (i)

Mar 27, 2023 – 03:32 PM EDT

PDB ID	:	8FUL
Title	:	Heterologous AibH1H2 purified from Lysogeny broth
Authors	:	Powell, M.M.; Rittle, J.
Deposited on	:	2023-01-17
Resolution	:	2.29 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.32.2
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.32.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motria	Whole archive	Similar resolution
Metric	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R_{free}	130704	5042(2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575(2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	А	392	7%	11%		11%
1	С	392	7%	13%		11%
1	Е	392	5%	11%		11%
1	G	392	8%	14%	•	10%
2	В	378	80%	1	6%	•



Continued from previous page...

Mol	Chain	Length	Quality of chain		
2	D	378	10%	18%	• •
3	F	378	74%	21%	•••
3	Н	378	75%	20%	



2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 23276 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	250	Total	С	Ν	0	S	0	1	0
	A	200	2807	1791	491	520	5	0	1	U
1	C	350	Total	С	Ν	0	S	0	1	0
1			2807	1791	491	520	5	0	1	0
1	F	250	Total	С	Ν	0	S	0	1	0
1		330	2791	1782	490	514	5	0		0
1	C	251	Total	С	Ν	0	S	0	ົງ	0
I G	591	2823	1801	495	522	5	0	3	U	

• Molecule 1 is a protein called Amidohydrolase.

There are 76 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-6	MET	-	expression tag	UNP A0A402C2V4
А	-5	GLY	-	expression tag	UNP A0A402C2V4
А	-4	HIS	-	expression tag	UNP A0A402C2V4
А	-3	HIS	-	expression tag	UNP A0A402C2V4
А	-2	HIS	-	expression tag	UNP A0A402C2V4
А	-1	HIS	-	expression tag	UNP A0A402C2V4
А	0	HIS	-	expression tag	UNP A0A402C2V4
А	1	HIS	-	expression tag	UNP A0A402C2V4
А	2	SER	-	expression tag	UNP A0A402C2V4
А	3	GLY	-	expression tag	UNP A0A402C2V4
А	4	GLU	-	expression tag	UNP A0A402C2V4
А	5	ASN	-	expression tag	UNP A0A402C2V4
А	6	LEU	-	expression tag	UNP A0A402C2V4
А	7	TYR	-	expression tag	UNP A0A402C2V4
А	8	PHE	-	expression tag	UNP A0A402C2V4
А	9	GLN	-	expression tag	UNP A0A402C2V4
А	10	SER	-	expression tag	UNP A0A402C2V4
А	11	GLY	-	expression tag	UNP A0A402C2V4
А	12	GLY	-	expression tag	UNP A0A402C2V4
С	-6	MET	-	expression tag	UNP A0A402C2V4
С	-5	GLY	-	expression tag	UNP A0A402C2V4



Chain	Residue	Modelled	Actual	Comment	Reference
С	-4	HIS	-	expression tag	UNP A0A402C2V4
С	-3	HIS	-	expression tag	UNP A0A402C2V4
С	-2	HIS	-	expression tag	UNP A0A402C2V4
С	-1	HIS	-	expression tag	UNP A0A402C2V4
С	0	HIS	-	expression tag	UNP A0A402C2V4
С	1	HIS	-	expression tag	UNP A0A402C2V4
С	2	SER	_	expression tag	UNP A0A402C2V4
С	3	GLY	-	expression tag	UNP A0A402C2V4
С	4	GLU	-	expression tag	UNP A0A402C2V4
С	5	ASN	-	expression tag	UNP A0A402C2V4
С	6	LEU	-	expression tag	UNP A0A402C2V4
С	7	TYR	-	expression tag	UNP A0A402C2V4
С	8	PHE	-	expression tag	UNP A0A402C2V4
С	9	GLN	-	expression tag	UNP A0A402C2V4
С	10	SER	-	expression tag	UNP A0A402C2V4
С	11	GLY	-	expression tag	UNP A0A402C2V4
С	12	GLY	-	expression tag	UNP A0A402C2V4
Е	-6	MET	-	expression tag	UNP A0A402C2V4
Е	-5	GLY	-	expression tag	UNP A0A402C2V4
Е	-4	HIS	-	expression tag	UNP A0A402C2V4
Е	-3	HIS	-	expression tag	UNP A0A402C2V4
Е	-2	HIS	-	expression tag	UNP A0A402C2V4
Е	-1	HIS	-	expression tag	UNP A0A402C2V4
Е	0	HIS	-	expression tag	UNP A0A402C2V4
Е	1	HIS	-	expression tag	UNP A0A402C2V4
Е	2	SER	-	expression tag	UNP A0A402C2V4
E	3	GLY	-	expression tag	UNP A0A402C2V4
E	4	GLU	-	expression tag	UNP A0A402C2V4
E	5	ASN	-	expression tag	UNP A0A402C2V4
E	6	LEU	-	expression tag	UNP A0A402C2V4
E	7	TYR	-	expression tag	UNP A0A402C2V4
E	8	PHE	-	expression tag	UNP A0A402C2V4
E	9	GLN	-	expression tag	UNP A0A402C2V4
E	10	SER	-	expression tag	UNP A0A402C2V4
E	11	GLY	-	expression tag	UNP A0A402C2V4
E	12	GLY	-	expression tag	UNP A0A402C2V4
G	-6	MET	-	expression tag	UNP A0A402C2V4
G	-5	GLY	-	expression tag	UNP A0A402C2V4
G	-4	HIS	-	expression tag	UNP A0A402C2V4
G	-3	HIS	-	expression tag	UNP A0A402C2V4
G	-2	HIS	-	expression tag	UNP A0A402C2V4
G	-1	HIS	-	expression tag	UNP A0A402C2V4

Continued from previous page...



Chain	Residue	Modelled	Actual	Comment	Reference
G	0	HIS	-	expression tag	UNP A0A402C2V4
G	1	HIS	-	expression tag	UNP A0A402C2V4
G	2	SER	-	expression tag	UNP A0A402C2V4
G	3	GLY	-	expression tag	UNP A0A402C2V4
G	4	GLU	-	expression tag	UNP A0A402C2V4
G	5	ASN	-	expression tag	UNP A0A402C2V4
G	6	LEU	-	expression tag	UNP A0A402C2V4
G	7	TYR	-	expression tag	UNP A0A402C2V4
G	8	PHE	-	expression tag	UNP A0A402C2V4
G	9	GLN	-	expression tag	UNP A0A402C2V4
G	10	SER	-	expression tag	UNP A0A402C2V4
G	11	GLY	-	expression tag	UNP A0A402C2V4
G	12	GLY	_	expression tag	UNP A0A402C2V4

Continued from previous page...

• Molecule 2 is a protein called Amidohydrolase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
9	В	363	Total	С	Ν	Ο	S	0	1	0
	D		2866	1824	493	540	9	0	L I	
0	а	260	Total	С	Ν	0	\mathbf{S}	0	1	0
	D	502	2842	1812	486	535	9	0		0

• Molecule 3 is a protein called Amidohydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	262	Total	С	Ν	Ο	S	0	2	0
0	Ľ	505	2860	1826	484	541	9	0		
3	ц	262	Total	С	Ν	0	S	0	9	0
5	11	505	2862	1827	487	539	9	0		

• Molecule 4 is FE (III) ION (three-letter code: FE) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	Total Fe 1 1	0	0
4	В	2	Total Fe 2 2	0	0
4	С	1	Total Fe 1 1	0	0
4	D	2	Total Fe 2 2	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	Е	1	Total Fe 1 1	0	0
4	F	2	Total Fe 2 2	0	0
4	G	1	Total Fe 1 1	0	0
4	Н	2	Total Fe 2 2	0	0

• Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
5	D	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0

• Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	В	1	Total Mg 1 1	0	0

• Molecule 7 is water.



8FUL

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	А	74	Total O 74 74	0	0
7	В	87	Total O 87 87	0	0
7	С	83	Total O 83 83	0	0
7	D	76	Total O 76 76	0	0
7	Е	86	Total O 86 86	0	0
7	F	57	$\begin{array}{cc} {\rm Total} & {\rm O} \\ 57 & 57 \end{array}$	0	0
7	G	71	Total O 71 71	0	0
7	Н	59	Total O 59 59	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Amidohydrolase



V378 PRO SER PRO GLY LYS VAL GLY GLY • Molecule 1: Amidohydrolase 8% Chain G: 74% 14% 10% • MET GLY GLY HIS HIS HIS HIS HIS SER HIS SER ALA SER CU VAL CLY VAL ALA RO THR SER ASN PRO GLY VAL PRO PRO ASP GLU GLU ASP PRO SER PRO GLY LYS VAL GLY • Molecule 2: Amidohydrolase Chain B: 80% 16% MET THR ILE GLU GLU GLY SER SER CLY GLY <mark>V373</mark> THR ASN GLN ALA ASP • Molecule 2: Amidohydrolase 10% Chain D: 76% . . 18% MET THR ILE GLU GLY SER LEU GLY THR 82





• Molecule 3: Amidohydrolase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants	82.77Å 231.40Å 145.00Å	Deperitor
a, b, c, α , β , γ	90.00° 92.17° 90.00°	Depositor
$\mathbf{P}_{\text{assolution}}(\hat{\mathbf{A}})$	47.28 - 2.29	Depositor
Resolution (A)	47.28 - 2.29	EDS
% Data completeness	97.9 (47.28-2.29)	Depositor
(in resolution range)	97.9 (47.28-2.29)	EDS
R _{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.18 (at 2.29 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
D D.	0.190 , 0.258	Depositor
$\mathbf{n}, \mathbf{n}_{free}$	0.190 , 0.258	DCC
R_{free} test set	2000 reflections $(1.68%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	46.1	Xtriage
Anisotropy	0.071	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.39, 43.6	EDS
L-test for twinning ²	$< L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	0.059 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	23276	wwPDB-VP
Average B, all atoms $(Å^2)$	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.86% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FE, MG, CSO, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles	
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.46	0/2894	0.61	0/3955
1	С	0.41	0/2894	0.58	0/3955
1	Е	0.42	0/2878	0.59	0/3936
1	G	0.42	0/2914	0.60	1/3982~(0.0%)
2	В	0.46	0/2953	0.62	1/4032~(0.0%)
2	D	0.53	0/2929	0.68	4/4002~(0.1%)
3	F	0.43	0/2944	0.61	1/4023~(0.0%)
3	Н	0.43	0/2946	0.63	2/4025~(0.0%)
All	All	0.45	0/23352	0.62	9/31910~(0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	1
3	F	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	D	115	LEU	CA-CB-CG	6.25	129.68	115.30
2	В	12	LEU	CA-CB-CG	6.15	129.45	115.30
3	Н	212	LEU	CA-CB-CG	6.13	129.40	115.30
3	Н	78	PRO	N-CA-C	6.06	127.86	112.10
3	F	115	LEU	C-N-CA	-6.03	106.64	121.70



There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	F	170	VAL	Mainchain
1	G	97	GLU	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2807	0	2705	39	0
1	С	2807	0	2705	37	0
1	Е	2791	0	2682	26	1
1	G	2823	0	2716	45	1
2	В	2866	0	2749	54	0
2	D	2842	0	2713	58	0
3	F	2860	0	2726	79	0
3	Н	2862	0	2733	54	0
4	А	1	0	0	0	0
4	В	2	0	0	0	0
4	С	1	0	0	0	0
4	D	2	0	0	0	0
4	Е	1	0	0	0	0
4	F	2	0	0	0	0
4	G	1	0	0	0	0
4	Н	2	0	0	0	0
5	В	6	0	7	0	0
5	D	6	0	7	0	0
6	В	1	0	0	0	0
7	А	74	0	0	1	0
7	В	87	0	0	1	0
7	С	83	0	0	1	0
7	D	76	0	0	1	0
7	Е	86	0	0	0	0
7	F	57	0	0	1	0
7	G	71	0	0	1	0
7	Н	59	0	0	1	0
All	All	23276	0	21743	357	1





The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 357 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:110:PRO:HD2	2:B:116:ASN:ND2	1.69	1.06
3:H:72:MET:CE	3:H:340:HIS:CD2	2.43	1.02
1:C:194:GLU:OE2	1:G:163:ARG:HD2	1.65	0.97
3:F:105:LEU:HD21	3:F:149:MET:SD	2.04	0.96
3:H:72:MET:HE2	3:H:340:HIS:CD2	2.01	0.96

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:171:PHE:O	1:G:288:ARG:NH1[4_545]	2.13	0.07

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	349/392~(89%)	340 (97%)	8 (2%)	1 (0%)	41	50
1	С	349/392~(89%)	338~(97%)	10 (3%)	1 (0%)	41	50
1	Е	349/392~(89%)	338 (97%)	10 (3%)	1 (0%)	41	50
1	G	352/392~(90%)	341 (97%)	10 (3%)	1 (0%)	41	50
2	В	362/378~(96%)	340 (94%)	22 (6%)	0	100	100
2	D	361/378~(96%)	331 (92%)	29 (8%)	1 (0%)	41	50
3	F	362/378~(96%)	340 (94%)	21 (6%)	1 (0%)	41	50
3	Н	362/378~(96%)	335 (92%)	25 (7%)	2 (1%)	25	31
All	All	2846/3080~(92%)	2703 (95%)	135 (5%)	8 (0%)	41	50



5 of 8 Ramachandran outliers are listed below:

Mol	Chain	\mathbf{Res}	Type
3	Н	31	GLN
1	С	270	PHE
2	D	328	CYS
1	Е	270	PHE
1	G	270	PHE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percen	tiles
1	А	300/332~(90%)	296~(99%)	4 (1%)	69	82
1	С	300/332~(90%)	294 (98%)	6 (2%)	55	72
1	Ε	296/332~(89%)	290~(98%)	6 (2%)	55	72
1	G	301/332~(91%)	296~(98%)	5 (2%)	60	76
2	В	305/318~(96%)	303~(99%)	2 (1%)	84	92
2	D	300/318~(94%)	298~(99%)	2 (1%)	84	92
3	F	302/317~(95%)	299~(99%)	3 (1%)	76	87
3	Н	302/317~(95%)	297 (98%)	5 (2%)	60	76
All	All	2406/2598~(93%)	2373~(99%)	33 (1%)	67	81

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	Н	42	ARG
3	Н	90	LEU
3	Н	341	PHE
2	D	90	LEU
1	С	333	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:



Mol	Chain	Res	Type
3	Н	340	HIS
3	Н	99	ASN
3	F	116	ASN
2	D	113	GLN
3	Н	98	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Type Chain B	Dec	Tink	B	Bond lengths			Bond angles		
	туре	Unam	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	CSO	F	328	3	4,5,7	1.08	0	$1,\!5,\!8$	0.43	0
3	CSO	Н	328	3	4,5,7	0.92	0	$1,\!5,\!8$	1.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CSO	F	328	3	-	1/1/4/7	-
3	CSO	Н	328	3	-	1/1/4/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	F	328	CSO	N-CA-CB-SG
3	Н	328	CSO	N-CA-CB-SG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 15 ligands modelled in this entry, 13 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	nin Dec Link		Chain Bos Link Bond lengths				E	Bond ang	gles
	туре	Unam	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2	
5	GOL	В	401	4	5,5,5	0.94	0	5,5,5	0.82	0	
5	GOL	D	401	4	5,5,5	0.85	0	5,5,5	1.00	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	В	401	4	-	3/4/4/4	-
5	GOL	D	401	4	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 7 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
5	D	401	GOL	O1-C1-C2-C3
5	D	401	GOL	C1-C2-C3-O3
5	В	401	GOL	O1-C1-C2-C3
5	D	401	GOL	O2-C2-C3-O3
5	D	401	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2			$OWAB(Å^2)$	Q<0.9
1	А	350/392~(89%)	0.45	26 (7%)	14	19	31, 42, 58, 82	0
1	С	350/392~(89%)	0.51	27 (7%)	13	17	33, 46, 63, 88	0
1	Е	350/392~(89%)	0.46	19 (5%)	25	32	35, 45, 58, 70	0
1	G	351/392~(89%)	0.60	31 (8%)	10	13	34, 49, 66, 87	0
2	В	363/378~(96%)	0.43	26 (7%)	15	20	33, 44, 69, 92	0
2	D	362/378~(95%)	0.64	36 (9%)	7	10	33, 45, 75, 94	0
3	F	362/378~(95%)	0.69	46 (12%) 3	5	35, 48, 79, 99	0
3	Н	362/378~(95%)	0.85	57 (15%) 2	2	33, 48, 81, 96	0
All	All	2850/3080~(92%)	0.58	268 (9%)	8	11	31, 46, 72, 99	0

The worst 5 of 268 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	80	SER	9.3
2	D	112	GLY	8.8
2	D	115	LEU	8.6
2	D	114	THR	8.2
1	G	28	GLY	7.7

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
3	CSO	F	328	6/8	0.93	0.14	44,46,47,52	0



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q < 0.9
3	CSO	Н	328	6/8	0.94	0.14	$46,\!47,\!51,\!57$	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
5	GOL	D	401	6/6	0.77	0.26	$60,\!63,\!82,\!83$	0
5	GOL	В	401	6/6	0.80	0.26	46,54,67,68	0
6	MG	В	404	1/1	0.87	0.13	$57,\!57,\!57,\!57$	0
4	FE	Н	402	1/1	0.93	0.06	80,80,80,80	0
4	FE	D	402	1/1	0.93	0.07	74,74,74,74	0
4	FE	F	402	1/1	0.96	0.05	80,80,80,80	0
4	FE	D	403	1/1	0.97	0.04	58, 58, 58, 58	0
4	FE	Н	401	1/1	0.98	0.06	58, 58, 58, 58	0
4	FE	В	402	1/1	0.98	0.07	68,68,68,68	0
4	FE	В	403	1/1	0.98	0.07	$55,\!55,\!55,\!55$	0
4	FE	F	401	1/1	0.98	0.05	60,60,60,60	0
4	FE	С	401	1/1	0.98	0.04	48,48,48,48	0
4	FE	G	401	1/1	0.99	0.05	52, 52, 52, 52, 52	0
4	FE	А	401	1/1	0.99	0.05	42,42,42,42	0
4	FE	E	401	1/1	0.99	0.07	49,49,49,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

















































6.5 Other polymers (i)

There are no such residues in this entry.

