



wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 02:55 am BST

PDB ID : 3FZ3
Title : Crystal Structure of almond Pru1 protein
Authors : Jin, T.C.; Zhang, Y.Z.
Deposited on : 2009-01-23
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	531	 48% 20% 5% 27%
1	B	531	 51% 21% 5% 23%
1	C	531	 53% 19% 5% 23%
1	D	531	 52% 18% 5% 27%
1	E	531	 50% 18% 5% 27%
1	F	531	 52% 20% 5% 23%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 19957 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Prunin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	388	Total 3104	C 1926	N 583	O 591	S 4	0	2	0
1	B	410	Total 3296	C 2034	N 626	O 632	S 4	0	1	0
1	C	410	Total 3298	C 2035	N 628	O 631	S 4	0	1	0
1	D	388	Total 3102	C 1924	N 583	O 591	S 4	0	1	0
1	E	388	Total 3097	C 1922	N 580	O 591	S 4	0	1	0
1	F	410	Total 3302	C 2038	N 627	O 633	S 4	0	2	0

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total Ca 1 1	0	0
2	E	1	Total Ca 1 1	0	0
2	B	1	Total Ca 1 1	0	0
2	C	1	Total Ca 1 1	0	0
2	A	1	Total Ca 1 1	0	0
2	F	1	Total Ca 1 1	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total Na 1 1	0	0
3	E	2	Total Na 2 2	0	0
3	B	1	Total Na 1 1	0	0
3	C	1	Total Na 1 1	0	0
3	A	2	Total Na 2 2	0	0
3	F	2	Total Na 2 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	90	Total O 90 90	0	0
4	B	162	Total O 162 162	0	0
4	C	123	Total O 123 123	0	0
4	D	136	Total O 136 136	0	0
4	E	102	Total O 102 102	0	0
4	F	130	Total O 130 130	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	151.02Å 151.02Å 165.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.08 – 2.40 36.09 – 2.40	Depositor EDS
% Data completeness (in resolution range)	88.7 (36.08-2.40) 88.0 (36.09-2.40)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.172 , 0.229 0.175 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	37.3	Xtrriage
Anisotropy	0.575	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 38.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.478 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	19957	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/3168	0.90	1/4291 (0.0%)
1	B	0.78	0/3358	1.01	6/4542 (0.1%)
1	C	0.74	0/3360	1.00	9/4544 (0.2%)
1	D	0.75	0/3163	0.98	3/4284 (0.1%)
1	E	0.72	0/3158	0.94	1/4278 (0.0%)
1	F	0.80	0/3367	1.02	3/4554 (0.1%)
All	All	0.75	0/19574	0.97	23/26493 (0.1%)

There are no bond length outliers.

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	321	ARG	NE-CZ-NH1	-7.18	116.71	120.30
1	D	195	ARG	NE-CZ-NH1	-6.43	117.09	120.30
1	C	189	ARG	NE-CZ-NH1	5.96	123.28	120.30
1	A	194	ARG	NE-CZ-NH1	5.93	123.27	120.30
1	C	236	ASP	CB-CG-OD2	5.87	123.58	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3104	0	2974	95	0
1	B	3296	0	3143	97	0
1	C	3298	0	3148	88	0
1	D	3102	0	2972	66	0
1	E	3097	0	2963	97	0
1	F	3302	0	3151	111	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	2	0	0	0	0
3	F	2	0	0	0	0
4	A	90	0	0	7	0
4	B	162	0	0	11	0
4	C	123	0	0	5	0
4	D	136	0	0	4	0
4	E	102	0	0	5	0
4	F	130	0	0	4	0
All	All	19957	0	18351	494	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 494 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:250:ASN:HB3	1:B:253:ASN:ND2	1.43	1.30
1:B:250:ASN:CB	1:B:253:ASN:HD21	1.49	1.23
1:A:195:ARG:HH21	1:A:195:ARG:HG3	1.05	1.11
1:F:195:ARG:CG	1:F:195:ARG:HH21	1.67	1.08
1:F:195:ARG:HG3	1:F:195:ARG:NH2	1.53	1.05

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	382/531 (72%)	342 (90%)	36 (9%)	4 (1%)	15	23
1	B	403/531 (76%)	371 (92%)	32 (8%)	0	100	100
1	C	403/531 (76%)	375 (93%)	24 (6%)	4 (1%)	15	23
1	D	381/531 (72%)	356 (93%)	23 (6%)	2 (0%)	29	41
1	E	381/531 (72%)	354 (93%)	22 (6%)	5 (1%)	12	17
1	F	404/531 (76%)	372 (92%)	30 (7%)	2 (0%)	29	41
All	All	2354/3186 (74%)	2170 (92%)	167 (7%)	17 (1%)	22	32

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	328[A]	ARG
1	A	328[B]	ARG
1	C	340	ARG
1	E	328	ARG
1	E	330	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	330/452 (73%)	272 (82%)	58 (18%)	2	2
1	B	349/452 (77%)	293 (84%)	56 (16%)	2	3
1	C	349/452 (77%)	312 (89%)	37 (11%)	6	9

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	330/452 (73%)	289 (88%)	41 (12%)	4	5
1	E	329/452 (73%)	281 (85%)	48 (15%)	3	3
1	F	350/452 (77%)	304 (87%)	46 (13%)	4	4
All	All	2037/2712 (75%)	1751 (86%)	286 (14%)	3	4

5 of 286 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	347	GLU
1	D	195	ARG
1	F	347	GLU
1	C	358	GLN
1	C	483	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 89 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	350	GLN
1	D	78	ASN
1	F	302	GLN
1	C	358	GLN
1	D	29	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 15 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	388/531 (73%)	-0.06	1 (0%) 94 93	27, 45, 66, 84	0
1	B	410/531 (77%)	-0.04	5 (1%) 79 77	21, 39, 81, 96	0
1	C	410/531 (77%)	-0.09	3 (0%) 87 86	22, 38, 80, 103	0
1	D	388/531 (73%)	-0.11	0 100 100	21, 37, 61, 87	0
1	E	388/531 (73%)	-0.06	1 (0%) 94 93	26, 45, 67, 80	0
1	F	410/531 (77%)	-0.11	0 100 100	21, 38, 75, 86	0
All	All	2394/3186 (75%)	-0.08	10 (0%) 92 91	21, 41, 68, 103	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	359	GLN	5.0
1	B	27	SER	2.9
1	B	357	ARG	2.8
1	C	353	LEU	2.7
1	B	353	LEU	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CA	A	1	1/1	0.92	0.21	52,52,52,52	0
3	NA	D	552	1/1	0.94	0.35	58,58,58,58	0
3	NA	E	552	1/1	0.94	0.25	67,67,67,67	0
3	NA	C	552	1/1	0.96	0.31	51,51,51,51	0
3	NA	A	552	1/1	0.96	0.26	53,53,53,53	0
3	NA	F	552	1/1	0.97	0.23	43,43,43,43	0
3	NA	F	7	1/1	0.97	0.23	53,53,53,53	0
2	CA	B	2	1/1	0.97	0.15	53,53,53,53	0
2	CA	F	6	1/1	0.98	0.14	51,51,51,51	0
3	NA	A	9	1/1	0.98	0.18	50,50,50,50	0
2	CA	D	4	1/1	0.99	0.17	45,45,45,45	0
3	NA	B	552	1/1	0.99	0.29	40,40,40,40	0
3	NA	E	8	1/1	0.99	0.20	44,44,44,44	0
2	CA	E	5	1/1	0.99	0.18	51,51,51,51	0
2	CA	C	3	1/1	0.99	0.18	45,45,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.