



Full wwPDB X-ray Structure Validation Report i

May 25, 2020 – 05:36 am BST

PDB ID : 4G8X
Title : G1 ORF67 / Staphylococcus aureus sigmaA domain 4 complex
Authors : Darst, S.A.; Osmundson, J.S.; Montero-Diaz, C.M.; Hochschild, A.
Deposited on : 2012-07-23
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.1.3
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 3325 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA polymerase sigma factor rpoD.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	61	Total	C	N	O	0	0	0
			460	283	90	87			

1	C	54	Total	C	N	O	0	0	0
			302	181	63	58			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	296	MSE	LEU	CONFLICT	UNP P0A0J0
C	296	MSE	LEU	CONFLICT	UNP P0A0J0

- Molecule 2 is a protein called ORF067.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	197	Total	C	N	O	S	Se	0	0	0
			1526	964	266	291	1	4			

2	D	177	Total	C	N	O	Se		0	0	0
			1037	627	192	215	3				


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RNA polymerase sigma factor rpoD




- Molecule 1: RNA polymerase sigma factor rpoD



- Molecule 2: ORF067



- Molecule 2: ORF067



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	219.14 Å 56.95 Å 57.03 Å 90.00° 104.79° 90.00°	Depositor
Resolution (Å)	37.71 – 3.00 37.71 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (37.71-3.00) 99.8 (37.71-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	2.41 (at 3.01 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R , R_{free}	0.281 , 0.328 0.281 , 0.329	Depositor DCC
R_{free} test set	1382 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	86.8	Xtriage
Anisotropy	0.156	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 92.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.029 for -h+k-l,-l,-k 0.000 for -h-k-l,l,k 0.020 for -h-2*l,-k,l	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	3325	wwPDB-VP
Average B, all atoms (Å ²)	123.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.24	0/461	0.36	0/618
1	C	0.32	0/301	0.57	0/410
2	B	0.29	1/1549 (0.1%)	0.45	1/2091 (0.0%)
2	D	0.28	0/1043	0.41	0/1401
All	All	0.29	1/3354 (0.0%)	0.44	1/4520 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	92	PRO	N-CD	5.23	1.55	1.47

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	29	PRO	N-CA-CB	5.90	110.38	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	460	0	439	7	0
1	C	302	0	182	3	0
2	B	1526	0	1441	39	0
2	D	1037	0	673	11	0
All	All	3325	0	2735	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:92:PRO:O	2:B:94:VAL:N	1.68	1.26
2:B:81:ASN:C	2:B:92:PRO:CD	2.35	0.95
2:B:81:ASN:O	2:B:92:PRO:CD	2.22	0.88
2:B:81:ASN:O	2:B:92:PRO:CG	2.22	0.87
2:B:81:ASN:C	2:B:92:PRO:HD3	1.97	0.83
2:B:81:ASN:O	2:B:92:PRO:HG2	1.79	0.83
2:B:81:ASN:HA	2:B:92:PRO:HG3	1.62	0.80
2:B:91:ASN:CB	2:B:92:PRO:HA	2.18	0.73
2:B:81:ASN:O	2:B:92:PRO:HD2	1.90	0.71
2:B:81:ASN:C	2:B:92:PRO:CG	2.59	0.71
2:D:129:ILE:HD11	2:D:190:LEU:HD13	1.74	0.69
2:B:92:PRO:O	2:B:93:ALA:C	2.31	0.69
2:B:81:ASN:CA	2:B:92:PRO:HG3	2.24	0.68
2:B:82:HIS:N	2:B:92:PRO:HD3	2.10	0.66
2:B:92:PRO:C	2:B:94:VAL:N	2.52	0.60
2:B:18:ARG:HD2	2:B:49:ARG:NH1	2.18	0.59
2:D:113:ILE:HG23	2:D:114:THR:H	1.68	0.58
2:B:7:ASP:O	2:B:9:ASP:N	2.36	0.56
2:B:37:TYR:O	2:B:98:ASN:ND2	2.37	0.56
2:B:18:ARG:HD2	2:B:49:ARG:HH12	1.70	0.56
2:B:82:HIS:HA	2:B:92:PRO:HD2	1.88	0.56
2:B:1:MSE:SE	2:B:45:TYR:H	2.39	0.55
2:B:91:ASN:CB	2:B:92:PRO:CA	2.85	0.55
2:B:92:PRO:O	2:B:94:VAL:CA	2.54	0.55
2:B:1:MSE:HE3	2:B:13:LEU:HD21	1.89	0.53
2:D:98:ASN:O	2:D:102:ILE:HG12	2.09	0.51
1:A:322:ASP:HB3	2:B:194:LYS:NZ	2.28	0.49
2:B:159:LYS:H	2:B:159:LYS:HD3	1.77	0.48
2:B:54:LEU:HG	2:B:86:ILE:HD11	1.95	0.48
2:B:81:ASN:CA	2:B:92:PRO:CG	2.91	0.48
1:A:304:LEU:HB3	1:A:312:GLU:HG3	1.95	0.47
1:A:320:GLY:HA3	1:A:325:ARG:O	2.15	0.47
2:B:152:GLU:HA	2:B:156:PHE:HB3	1.96	0.47
2:B:159:LYS:HA	2:B:160:SER:HA	1.71	0.46
2:D:125:GLY:O	2:D:129:ILE:HG13	2.14	0.46
2:B:2:LYS:HB2	2:B:14:ASN:HB2	1.98	0.46
2:D:150:GLU:N	2:D:151:GLU:HA	2.30	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:83:ARG:C	2:B:91:ASN:O	2.55	0.45
1:C:334:LYS:HA	1:C:335:VAL:HA	1.64	0.44
2:B:98:ASN:HA	2:B:101:ILE:HD12	1.98	0.44
1:C:304:LEU:CB	1:C:305:ASP:HA	2.48	0.44
2:B:18:ARG:HB3	2:B:19:ASN:H	1.49	0.43
2:B:92:PRO:C	2:B:94:VAL:H	2.16	0.43
2:D:144:TYR:CZ	2:D:168:ALA:HA	2.53	0.43
1:A:329:LEU:HD23	1:A:340:ARG:HB3	2.00	0.43
2:B:92:PRO:O	2:B:95:ARG:N	2.50	0.42
1:A:316:ARG:HD3	1:A:322:ASP:OD2	2.19	0.42
1:C:323:ASP:OD2	1:C:327:ARG:NH1	2.53	0.42
2:D:114:THR:HA	2:D:115:GLU:HA	1.80	0.42
2:D:43:TYR:HD1	2:D:58:ARG:HA	1.84	0.42
1:A:310:ARG:HD2	2:B:41:ASN:HA	2.01	0.42
2:B:126:PHE:HA	2:B:129:ILE:HG22	2.02	0.41
2:B:41:ASN:CG	2:B:188:ASN:HD21	2.23	0.41
2:D:110:LEU:HA	2:D:111:PRO:HD3	1.96	0.41
1:A:327:ARG:HD3	1:A:327:ARG:HA	1.84	0.41
2:B:40:SER:O	2:B:41:ASN:HB2	2.21	0.41
2:D:58:ARG:HD3	2:D:72:ASN:OD1	2.21	0.40
2:D:170:LYS:HA	2:D:171:GLU:HA	1.75	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	59/62 (95%)	56 (95%)	3 (5%)	0	100 100
1	C	50/62 (81%)	43 (86%)	5 (10%)	2 (4%)	3 17
2	B	195/198 (98%)	179 (92%)	9 (5%)	7 (4%)	3 19
2	D	167/198 (84%)	147 (88%)	14 (8%)	6 (4%)	3 19

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	471/520 (91%)	425 (90%)	31 (7%)	15 (3%)	4 22

All (15) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	91	ASN
2	B	92	PRO
2	B	93	ALA
1	C	310	ARG
2	D	67	TYR
2	D	92	PRO
2	B	8	LYS
2	B	51	GLY
1	C	321	LEU
2	D	113	ILE
2	D	148	LYS
2	B	6	LEU
2	B	152	GLU
2	D	69	LEU
2	D	196	ILE

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	42/55 (76%)	42 (100%)	0	100 100
1	C	9/55 (16%)	9 (100%)	0	100 100
2	B	154/179 (86%)	147 (96%)	7 (4%)	27 64
2	D	55/179 (31%)	55 (100%)	0	100 100
All	All	260/468 (56%)	253 (97%)	7 (3%)	44 77

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	45	TYR
2	B	49	ARG
2	B	90	TYR
2	B	134	GLN
2	B	159	LYS
2	B	166	ASN
2	B	180	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	188	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	60/62 (96%)	-0.03	0 [100] [100]	68, 103, 183, 203	0
1	C	54/62 (87%)	0.02	3 (5%) 24 8	136, 193, 239, 271	0
2	B	193/198 (97%)	-0.22	1 (0%) 91 75	52, 81, 189, 221	0
2	D	173/198 (87%)	0.15	13 (7%) 14 4	85, 162, 222, 265	0
All	All	480/520 (92%)	-0.04	17 (3%) 44 18	52, 119, 221, 271	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	84	CYS	7.7
2	D	73	ASN	5.7
1	C	322	ASP	4.6
2	D	80	ASN	4.2
2	D	74	GLN	4.1
1	C	340	ARG	4.0
2	D	24	THR	3.5
2	D	87	GLY	3.4
2	D	37	TYR	3.3
2	D	71	ASP	3.2
2	D	7	ASP	3.2
2	D	89	TRP	2.5
2	D	79	ASP	2.4
2	B	90	TYR	2.4
2	D	160	SER	2.1
2	D	161	GLU	2.1
1	C	339	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.