



# Full wwPDB X-ray Structure Validation Report ⓘ

May 24, 2020 – 10:23 pm BST

PDB ID : 6GB5  
Title : Structure of H-2Db with truncated SEV peptide and GL  
Authors : Hafstrand, I.; Sandalova, T.; Achour, A.  
Deposited on : 2018-04-13  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	338	
1	C	338	
2	B	99	
2	D	99	
3	E	7	
3	F	7	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	G	2	 50% 50%
4	H	2	 50% 50%

## 2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 6445 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H-2 class I histocompatibility antigen, D-B alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	274	Total	C	N	O	S	0	2	0
			2234	1414	394	417	9			
1	C	275	Total	C	N	O	S	0	2	0
			2242	1419	396	418	9			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	99	Total	C	N	O	S	0	0	0
			813	518	136	152	7			
2	D	99	Total	C	N	O	S	0	0	0
			821	524	138	152	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	85	ASP	ALA	variant	UNP P01887
D	85	ASP	ALA	variant	UNP P01887

- Molecule 3 is a protein called PHE-ALA-PRO-GLY-ASN-TYR-PRO.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	E	7	Total	C	N	O	0	0	0
			55	37	8	10			
3	F	7	Total	C	N	O	0	0	0
			55	37	8	10			

- Molecule 4 is a protein called GLY-LEU.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	G	2	Total	C	N	O	0	0	0
			13	8	2	3			

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	H	2	13	8	2	3	0	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
5	A	1	5	4	1	0	0
5	A	1	5	4	1	0	0
5	C	1	5	4	1	0	0
5	C	1	5	4	1	0	0
5	C	1	5	4	1	0	0
5	C	1	5	4	1	0	0

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 6 3 3	0	0
6	C	1	Total C O 6 3 3	0	0
6	D	1	Total C O 6 3 3	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	45	Total O 45 45	0	0
7	B	29	Total O 29 29	0	0
7	C	52	Total O 52 52	0	0
7	D	21	Total O 21 21	0	0
7	E	2	Total O 2 2	0	0
7	F	2	Total O 2 2	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: H-2 class I histocompatibility antigen, D-B alpha chain



- Molecule 1: H-2 class I histocompatibility antigen, D-B alpha chain




- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin





- Molecule 3: PHE-ALA-PRO-GLY-ASN-TYR-PRO

Chain E:  100%

There are no outlier residues recorded for this chain.

- Molecule 3: PHE-ALA-PRO-GLY-ASN-TYR-PRO

Chain F:  100%

There are no outlier residues recorded for this chain.

- Molecule 4: GLY-LEU

Chain G:  50% 50%



- Molecule 4: GLY-LEU

Chain H:  50% 50%





## 4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	91.93Å 123.36Å 99.21Å 90.00° 103.73° 90.00°	Depositor
Resolution (Å)	75.94 – 2.30 58.90 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.3 (75.94-2.30) 99.3 (58.90-2.30)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.81 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.198 , 0.240 0.205 , 0.240	Depositor DCC
$R_{free}$ test set	2185 reflections (4.61%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.8	Xtrriage
Anisotropy	0.042	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 36.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6445	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	59.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.72% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.79	0/2303	0.76	0/3132
1	C	0.83	0/2314	0.78	0/3144
2	B	0.81	0/839	0.81	0/1140
2	D	0.85	0/847	0.82	0/1148
3	E	0.85	0/58	0.68	0/78
3	F	0.79	0/58	0.68	0/78
4	G	1.28	0/12	0.89	0/13
4	H	0.73	0/12	0.69	0/13
All	All	0.82	0/6443	0.78	0/8746

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2234	0	2081	24	0
1	C	2242	0	2101	28	0
2	B	813	0	774	3	0
2	D	821	0	796	8	0
3	E	55	0	48	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	55	0	48	0	0
4	G	13	0	16	1	0
4	H	13	0	16	1	0
5	A	10	0	0	0	0
5	C	20	0	0	0	0
6	A	6	0	8	0	0
6	C	6	0	8	0	0
6	D	6	0	8	1	0
7	A	45	0	0	1	0
7	B	29	0	0	0	0
7	C	52	0	0	0	0
7	D	21	0	0	0	0
7	E	2	0	0	0	0
7	F	2	0	0	0	0
All	All	6445	0	5904	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:CYS:SG	1:A:164:CYS:CB	2.13	1.36
1:A:194:ARG:HG2	1:A:195:SER:H	1.28	0.97
1:A:101:CYS:CB	1:A:164:CYS:SG	2.52	0.97
1:C:191:HIS:NE2	1:C:199:VAL:HG11	1.79	0.97
1:C:226:GLN:C	1:C:228:MET:N	2.23	0.92
1:A:194:ARG:HG2	1:A:195:SER:N	2.01	0.76
1:A:101:CYS:SG	1:A:164:CYS:SG	0.69	0.69
1:C:191:HIS:HE2	1:C:199:VAL:HG11	1.58	0.68
1:A:255:GLN:NE2	1:A:274:TRP:O	2.27	0.67
1:C:195:SER:O	1:C:197:GLY:N	2.28	0.65
1:A:101:CYS:CB	1:A:164:CYS:HG	2.06	0.64
1:C:251:LEU:HD12	1:C:252:GLY:H	1.64	0.63
1:C:191:HIS:CD2	1:C:199:VAL:HG11	2.34	0.61
1:A:194:ARG:CG	1:A:195:SER:H	2.10	0.61
1:C:81:LEU:HD11	4:G:2:LEU:HD12	1.85	0.59
2:D:9:VAL:H	6:D:101:GOL:H2	1.68	0.59
1:C:192:HIS:O	1:C:199:VAL:HG13	2.02	0.58
1:A:250:PRO:HG2	1:A:253:LYS:CB	2.33	0.58
1:A:217:TRP:CD1	1:A:247:VAL:HG23	2.41	0.56

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:MET:HB2	1:A:168:LEU:HD13	1.87	0.55
1:C:201:LEU:O	1:C:246:SER:HA	2.07	0.55
1:C:191:HIS:HD2	1:C:201:LEU:HD21	1.72	0.54
1:C:198:GLU:O	1:C:199:VAL:HG23	2.09	0.53
1:A:251:LEU:HD23	1:A:252:GLY:N	2.24	0.52
2:D:98:ASP:C	2:D:99:MET:HG2	2.30	0.52
1:C:145:ARG:O	1:C:149:GLN:HG2	2.10	0.52
1:C:199:VAL:CG1	1:C:200:THR:N	2.74	0.51
2:D:21:ASN:HB3	2:D:70:PHE:CE1	2.45	0.51
1:C:251:LEU:HD12	1:C:252:GLY:N	2.25	0.51
2:D:98:ASP:O	2:D:99:MET:HG2	2.10	0.51
1:C:252:GLY:N	1:C:254:GLU:OE2	2.44	0.50
1:A:55:GLU:OE1	1:A:170[A]:ARG:NH2	2.45	0.50
1:A:14:ARG:HD3	7:A:515:HOH:O	2.09	0.49
1:A:81:LEU:HD13	1:A:118:TYR:CD1	2.47	0.49
1:C:219:LEU:HB2	1:C:257:TYR:CE2	2.48	0.48
1:A:101:CYS:SG	1:A:164:CYS:HB3	2.38	0.47
1:C:199:VAL:HG12	1:C:200:THR:N	2.29	0.47
1:A:191:HIS:NE2	1:A:199:VAL:HG21	2.29	0.47
2:D:48:LYS:HG3	2:D:48:LYS:O	2.15	0.47
1:C:199:VAL:O	1:C:248:VAL:HA	2.15	0.47
1:C:50:PRO:O	1:C:53:GLU:HG3	2.15	0.46
1:A:55:GLU:HG2	1:A:59:TYR:CG	2.51	0.45
1:C:196:LYS:O	1:C:197:GLY:C	2.55	0.45
1:C:217:TRP:CD1	1:C:247:VAL:HG23	2.52	0.45
1:C:229:GLU:HG3	1:C:246:SER:OG	2.16	0.45
1:C:51:TRP:CZ3	1:C:52:MET:SD	3.10	0.45
1:C:117:ALA:HB2	2:D:60:TRP:CE2	2.53	0.44
1:C:271:THR:O	1:C:272:LEU:HD23	2.17	0.44
1:A:191:HIS:CE1	1:A:199:VAL:HG21	2.53	0.43
2:B:16:GLU:HG3	2:B:19:LYS:HB2	2.00	0.43
1:C:219:LEU:CD1	1:C:256:ASN:HB3	2.49	0.43
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.53	0.43
1:A:202:ARG:HG2	1:A:204:TRP:NE1	2.34	0.42
1:C:219:LEU:HD22	1:C:257:TYR:CZ	2.53	0.42
2:B:96:ASP:C	2:B:96:ASP:OD1	2.59	0.41
1:C:202:ARG:NH1	2:D:98:ASP:O	2.53	0.41
1:A:106:ASP:O	1:A:107:TRP:HB2	2.21	0.41
1:A:254:GLU:O	1:A:255:GLN:HB2	2.20	0.41
2:D:10:TYR:N	2:D:10:TYR:CD1	2.88	0.41
1:A:77:SER:HB3	4:H:2:LEU:HD12	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/338 (80%)	261 (96%)	9 (3%)	2 (1%)	22	26
1	C	273/338 (81%)	264 (97%)	8 (3%)	1 (0%)	34	42
2	B	97/99 (98%)	95 (98%)	2 (2%)	0	100	100
2	D	97/99 (98%)	95 (98%)	2 (2%)	0	100	100
3	E	5/7 (71%)	5 (100%)	0	0	100	100
3	F	5/7 (71%)	5 (100%)	0	0	100	100
All	All	749/888 (84%)	725 (97%)	21 (3%)	3 (0%)	34	42

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	196	LYS
1	A	225	THR
1	A	54	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/280 (81%)	225 (100%)	1 (0%)	91	96
1	C	228/280 (81%)	225 (99%)	3 (1%)	69	82

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	92/94 (98%)	91 (99%)	1 (1%)	73	86
2	D	94/94 (100%)	92 (98%)	2 (2%)	53	70
3	E	5/5 (100%)	5 (100%)	0	100	100
3	F	5/5 (100%)	5 (100%)	0	100	100
4	G	1/1 (100%)	1 (100%)	0	100	100
4	H	1/1 (100%)	1 (100%)	0	100	100
All	All	652/760 (86%)	645 (99%)	7 (1%)	78	86

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	101	CYS
2	B	70	PHE
1	C	144[A]	ARG
1	C	144[B]	ARG
1	C	194	ARG
2	D	58	LYS
2	D	70	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	191	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	SO4	C	402	-	4,4,4	0.37	0	6,6,6	0.16	0
5	SO4	A	402	-	4,4,4	0.41	0	6,6,6	0.21	0
6	GOL	D	101	-	5,5,5	0.13	0	5,5,5	0.45	0
5	SO4	C	403	-	4,4,4	0.40	0	6,6,6	0.39	0
6	GOL	C	405	-	5,5,5	0.52	0	5,5,5	0.57	0
5	SO4	C	404	-	4,4,4	0.54	0	6,6,6	0.64	0
5	SO4	C	401	-	4,4,4	0.48	0	6,6,6	0.48	0
6	GOL	A	403	-	5,5,5	0.30	0	5,5,5	0.43	0
5	SO4	A	401	-	4,4,4	0.35	0	6,6,6	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	D	101	-	-	2/4/4/4	-
6	GOL	A	403	-	-	0/4/4/4	-
6	GOL	C	405	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	C	405	GOL	O1-C1-C2-C3
6	C	405	GOL	C1-C2-C3-O3
6	C	405	GOL	O2-C2-C3-O3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
6	D	101	GOL	O1-C1-C2-C3
6	D	101	GOL	O1-C1-C2-O2
6	C	405	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	D	101	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	274/338 (81%)	0.39	20 (7%) 15 20	30, 59, 117, 134	0
1	C	275/338 (81%)	0.36	25 (9%) 9 12	29, 54, 116, 141	0
2	B	99/99 (100%)	-0.06	0 100 100	34, 48, 77, 83	0
2	D	99/99 (100%)	0.05	0 100 100	31, 53, 76, 91	0
3	E	7/7 (100%)	0.12	0 100 100	45, 52, 62, 64	0
3	F	7/7 (100%)	0.13	0 100 100	39, 42, 52, 65	0
4	G	2/2 (100%)	0.82	0 100 100	49, 49, 49, 66	0
4	H	2/2 (100%)	0.17	0 100 100	54, 54, 54, 61	0
All	All	765/892 (85%)	0.27	45 (5%) 22 28	29, 54, 111, 141	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	17	LEU	7.8
1	C	197	GLY	5.4
1	A	251	LEU	5.4
1	A	224	LEU	5.4
1	A	225	THR	5.1
1	A	274	TRP	5.1
1	A	250	PRO	4.6
1	C	16	GLY	4.6
1	C	220	ASN	4.6
1	C	249	VAL	4.4
1	A	249	VAL	4.4
1	A	179	LEU	3.7
1	C	276	PRO	3.6
1	C	257	TYR	3.6
1	A	257	TYR	3.5
1	C	274	TRP	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	195	SER	3.4
1	C	222	GLU	3.4
1	C	219	LEU	3.3
1	C	224	LEU	3.2
1	A	219	LEU	3.0
1	C	248	VAL	3.0
1	C	255	GLN	2.9
1	A	252	GLY	2.8
1	A	222	GLU	2.8
1	C	217	TRP	2.7
1	A	227	ASP	2.6
1	A	199	VAL	2.6
1	C	228	MET	2.5
1	C	196	LYS	2.5
1	C	199	VAL	2.4
1	C	247	VAL	2.3
1	A	192	HIS	2.3
1	C	253	LYS	2.3
1	A	178	THR	2.3
1	A	191	HIS	2.3
1	A	89	ALA	2.2
1	C	256	ASN	2.2
1	A	217	TRP	2.1
1	C	252	GLY	2.1
1	C	230	LEU	2.1
1	C	178	THR	2.0
1	A	54	GLN	2.0
1	A	248	VAL	2.0
1	C	201	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	SO4	C	404	5/5	0.75	0.23	100,107,124,128	0
6	GOL	D	101	6/6	0.78	0.21	63,75,79,79	0
5	SO4	C	403	5/5	0.84	0.23	119,119,123,127	0
5	SO4	A	401	5/5	0.85	0.21	123,125,134,134	0
6	GOL	A	403	6/6	0.86	0.18	56,59,63,69	0
5	SO4	A	402	5/5	0.88	0.31	117,124,129,133	0
5	SO4	C	402	5/5	0.88	0.19	115,124,127,129	0
6	GOL	C	405	6/6	0.90	0.20	47,50,52,54	0
5	SO4	C	401	5/5	0.96	0.10	84,91,99,100	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.