

# wwPDB X-ray Structure Validation Summary Report (i)

May 13, 2020 – 02:20 am BST

PDB ID : 6GBO

Title : Crystal Structure of the oligomerization domain of Vp35 from Ebola virus Authors : Zinzula, L.; Nagy, I.; Orsini, M.; Weyher-Stingl, E.; Baumeister, W.; Bracher,

Α.

Deposited on : 2018-04-16

Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467 Xtriage (Phenix) : 1.13

EDS : 2.11

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar  resolution} \\ (\#{\rm Entries,  resolution  range(\AA)}) \end{array}$
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1			7%		
1	A	73	97%		•
	_		16%		
1	В	73	96%		•
			16%		
1	С	73	85%	7%	8%
	-		10%		
1	D	73	89%	5%	5%
	_		12%		
1	E	73	97%		•
			21%		
1	F	73	95%		5%

Continued on next page...



Continued from previous page...

Mol	Chain	Length	Quality of chain	
			21%	
1	G	73	92%	7% •
			15%	
1	Н	73	93%	5% •
			12%	
1	I	73	85%	5% 10%
			11%	
1	J	73	86%	10% •
			14%	
1	K	73	92%	7% •
			23%	
1	$ box{L}$	73	86%	12% •



# 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 6842 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Polymerase cofactor VP35.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	73	Total	С	N	О	S	0	0	0
1	A	(3	578	360	101	113	4	0	U	0
1	В	73	Total	С	N	О	S	0	0	0
1	Б	(3	578	360	101	113	4	U	0	0
1	С	67	Total	С	N	О	S	0	0	0
1		07	514	322	83	106	3	0	U	0
1	D	69	Total	С	N	О	S	0	0	0
1	D	09	534	334	89	108	3	0	U	0
1	Е	73	Total	С	N	О	S	0	0	0
1	E	(3	574	358	101	111	4	0	0	
1	F	73	Total	С	N	О	S	0	0	0
1	$\Gamma$	(3	575	358	101	113	3	U	0	U
1	G	72	Total	С	N	О	S	0	0	0
1	G	12	562	351	99	109	3	0	U	0
1	Н	72	Total	С	N	О	S	0	0	0
1	11	12	566	353	100	110	3	0	0	0
1	I	66	Total	С	N	О	S	0	0	0
1	1	00	501	315	81	102	3	0	0	0
1	J	70	Total	С	N	О	S	0	0	0
1	J	10	547	342	92	109	4	0	0	0
1	K	72	Total	С	N	О	S	0	0	0
1	IX	12	570	355	100	112	3	U	U	0
1	L	72	Total	С	N	О	S	0	0	0
1	ь	12	570	355	100	112	3	U	U	U

There are 108 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	81	MET	_	initiating methionine	UNP Q05127
A	146	LEU	-	expression tag	UNP Q05127
A	147	GLU	-	expression tag	UNP Q05127
A	148	HIS	_	expression tag	UNP Q05127
A	149	HIS	-	expression tag	UNP Q05127



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	150	HIS	_	expression tag	UNP Q05127
A	151	HIS	-	expression tag	UNP Q05127
A	152	HIS	-	expression tag	UNP Q05127
A	153	HIS	-	expression tag	UNP Q05127
В	81	MET	_	initiating methionine	UNP Q05127
В	146	LEU	_	expression tag	UNP Q05127
В	147	GLU	-	expression tag	UNP Q05127
В	148	HIS	_	expression tag	UNP Q05127
В	149	HIS	_	expression tag	UNP Q05127
В	150	HIS	_	expression tag	UNP Q05127
В	151	HIS	_	expression tag	UNP Q05127
В	152	HIS	_	expression tag	UNP Q05127
В	153	HIS	_	expression tag	UNP Q05127
С	81	MET	_	initiating methionine	UNP Q05127
С	146	LEU	_	expression tag	UNP Q05127
С	147	GLU	_	expression tag	UNP Q05127
С	148	HIS	_	expression tag	UNP Q05127
С	149	HIS	_	expression tag	UNP Q05127
С	150	HIS	_	expression tag	UNP Q05127
С	151	HIS	-	expression tag	UNP Q05127
С	152	HIS	-	expression tag	UNP Q05127
С	153	HIS	_	expression tag	UNP Q05127
D	81	MET	_	initiating methionine	UNP Q05127
D	146	LEU	_	expression tag	UNP Q05127
D	147	GLU	_	expression tag	UNP Q05127
D	148	HIS	_	expression tag	UNP Q05127
D	149	HIS	_	expression tag	UNP Q05127
D	150	HIS	_	expression tag	UNP Q05127
D	151	HIS	_	expression tag	UNP Q05127
D	152	HIS	_	expression tag	UNP Q05127
D	153	HIS	_	expression tag	UNP Q05127
Е	81	MET	_	initiating methionine	UNP Q05127
Е	146	LEU	-	expression tag	UNP Q05127
Е	147	GLU	-	expression tag	UNP Q05127
Е	148	HIS	-	expression tag	UNP Q05127
Е	149	HIS	-	expression tag	UNP Q05127
Е	150	HIS	-	expression tag	UNP Q05127
Е	151	HIS	-	expression tag	UNP Q05127
Е	152	HIS	-	expression tag	UNP Q05127
Е	153	HIS	-	expression tag	UNP Q05127
F	81	MET	-	initiating methionine	UNP Q05127
F	146	LEU	-	expression tag	UNP Q05127



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	147	GLU	_	expression tag	UNP Q05127
F	148	HIS	_	expression tag	UNP Q05127
F	149	HIS	_	expression tag	UNP Q05127
F	150	HIS	-	expression tag	UNP Q05127
F	151	HIS	_	expression tag	UNP Q05127
F	152	HIS	_	expression tag	UNP Q05127
F	153	HIS	-	expression tag	UNP Q05127
G	81	MET	_	initiating methionine	UNP Q05127
G	146	LEU	_	expression tag	UNP Q05127
G	147	GLU	_	expression tag	UNP Q05127
G	148	HIS	_	expression tag	UNP Q05127
G	149	HIS	_	expression tag	UNP Q05127
G	150	HIS	_	expression tag	UNP Q05127
G	151	HIS	_	expression tag	UNP Q05127
G	152	HIS	_	expression tag	UNP Q05127
G	153	HIS	_	expression tag	UNP Q05127
Н	81	MET	_	initiating methionine	UNP Q05127
Н	146	LEU	_	expression tag	UNP Q05127
Н	147	GLU	_	expression tag	UNP Q05127
Н	148	HIS	-	expression tag	UNP Q05127
Н	149	HIS	-	expression tag	UNP Q05127
Н	150	HIS	_	expression tag	UNP Q05127
Н	151	HIS	_	expression tag	UNP Q05127
Н	152	HIS	_	expression tag	UNP Q05127
Н	153	HIS	_	expression tag	UNP Q05127
I	81	MET	_	initiating methionine	UNP Q05127
I	146	LEU	_	expression tag	UNP Q05127
I	147	GLU	_	expression tag	UNP Q05127
I	148	HIS	_	expression tag	UNP Q05127
I	149	HIS	-	expression tag	UNP Q05127
I	150	HIS	-	expression tag	UNP Q05127
I	151	HIS	-	expression tag	UNP Q05127
I	152	HIS	-	expression tag	UNP Q05127
I	153	HIS	-	expression tag	UNP Q05127
J	81	MET	-	initiating methionine	UNP Q05127
J	146	LEU	-	expression tag	UNP Q05127
J	147	GLU	-	expression tag	UNP Q05127
J	148	HIS	-	expression tag	UNP Q05127
J	149	HIS	-	expression tag	UNP Q05127
J	150	HIS	-	expression tag	UNP Q05127
J	151	HIS	-	expression tag	UNP Q05127
J	152	HIS	-	expression tag	UNP Q05127



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
J	153	HIS	-	expression tag	UNP Q05127
K	81	MET	-	initiating methionine	UNP Q05127
K	146	LEU	-	expression tag	UNP Q05127
K	147	GLU	-	expression tag	UNP Q05127
K	148	HIS	-	expression tag	UNP Q05127
K	149	HIS	-	expression tag	UNP Q05127
K	150	HIS	-	expression tag	UNP Q05127
K	151	HIS	-	expression tag	UNP Q05127
K	152	HIS	-	expression tag	UNP Q05127
K	153	HIS	-	expression tag	UNP Q05127
L	81	MET	-	initiating methionine	UNP Q05127
L	146	LEU	-	expression tag	UNP Q05127
L	147	GLU	-	expression tag	UNP Q05127
L	148	HIS	-	expression tag	UNP Q05127
L	149	HIS	-	expression tag	UNP Q05127
L	150	HIS	-	expression tag	UNP Q05127
L	151	HIS	-	expression tag	UNP Q05127
L	152	HIS	-	expression tag	UNP Q05127
L	153	HIS	-	expression tag	UNP Q05127

#### • Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	16	Total O 16 16	0	0
2	В	8	Total O 8 8	0	0
2	С	14	Total O 14 14	0	0
2	D	24	Total O 24 24	0	0
2	Е	11	Total O 11 11	0	0
2	F	6	Total O 6 6	0	0
2	G	14	Total O 14 14	0	0
2	Н	13	Total O 13 13	0	0
2	I	10	Total O 10 10	0	0
2	J	20	Total O 20 20	0	0



 $Continued\ from\ previous\ page...$ 


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	К	20	Total O 20 20	0	0
2	L	17	Total O 17 17	0	0



# 3 Residue-property plots (i)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Polymerase cofactor VP35



• Molecule 1: Polymerase cofactor VP35







# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	63.78Å 98.91Å 185.07Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 - 2.10	Depositor
resolution (A)	29.95 - 2.10	EDS
% Data completeness	99.8 (30.00-2.10)	Depositor
(in resolution range)	99.9 (29.95-2.10)	EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.84 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
P. P.	0.211 , 0.242	Depositor
$R, R_{free}$	0.216 , $0.245$	DCC
$R_{free}$ test set	3501  reflections  (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.6	Xtriage
Anisotropy	0.275	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.36, 47.5	EDS
L-test for twinning <sup>2</sup>	$ < L > = 0.49, < L^2> = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6842	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.87% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $< L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z >5	RMSZ	# Z  > 5	
1	A	0.43	0/587	0.53	0/793	
1	В	0.39	0/587	0.51	0/793	
1	С	0.42	0/517	0.56	0/700	
1	D	0.45	0/539	0.57	0/730	
1	E	0.44	0/583	0.50	0/788	
1	F	0.46	0/584	0.56	0/790	
1	G	0.49	0/571	0.53	0/773	
1	Н	0.41	0/575	0.51	0/778	
1	I	0.41	0/504	0.52	0/683	
1	J	0.50	0/553	0.60	0/748	
1	K	0.46	0/579	0.56	0/783	
1	L	0.47	0/579	0.58	0/783	
All	All	0.45	0/6758	0.54	0/9142	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

$\mathbf{Mol}$	Chain	Non-H	$\mathbf{H}(\mathbf{model})$	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes	
1	A	578	0	572	1	0	
1	В	578	0	572	3	0	
1	С	514	0	523	4	0	
1	D	534	0	537	3	0	



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	574	0	568	2	0
1	F	575	0	565	4	0
1	G	562	0	553	4	0
1	Н	566	0	559	5	0
1	I	501	0	511	4	0
1	J	547	0	551	12	0
1	K	570	0	563	4	0
1	L	570	0	563	11	0
2	A	16	0	0	0	0
2	В	8	0	0	0	0
2	С	14	0	0	1	0
2	D	24	0	0	0	0
2	Ε	11	0	0	0	0
2	F	6	0	0	0	0
2	G	14	0	0	0	0
2	Η	13	0	0	0	0
2	I	10	0	0	0	0
2	J	20	0	0	0	0
2	K	20	0	0	0	0
2	L	17	0	0	1	0
All	All	6842	0	6637	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 34 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} &  ext{Interatomic} \ &  ext{distance} \ &  ext{(Å)} \end{aligned}$	$egin{array}{c}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{array}$
1:J:93:LEU:C	1:J:93:LEU:HD23	2.19	0.63
1:B:111:ILE:HG12	1:C:111:ILE:HD11	1.82	0.61
1:A:90:LEU:HD21	1:B:89:THR:HB	1.85	0.59
1:J:111:ILE:HD11	1:L:111:ILE:HG12	1.85	0.58
1:J:90:LEU:HD21	1:K:89:THR:CG2	2.35	0.56

There are no symmetry-related clashes.



## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	71/73 (97%)	71 (100%)	0	0	100	100
1	В	71/73~(97%)	71 (100%)	0	0	100	100
1	С	65/73~(89%)	65 (100%)	0	0	100	100
1	D	$67/73 \ (92\%)$	67 (100%)	0	0	100	100
1	E	71/73 (97%)	71 (100%)	0	0	100	100
1	F	71/73 (97%)	71 (100%)	0	0	100	100
1	G	70/73~(96%)	70 (100%)	0	0	100	100
1	Н	70/73~(96%)	70 (100%)	0	0	100	100
1	I	64/73 (88%)	63 (98%)	1 (2%)	0	100	100
1	J	68/73 (93%)	67 (98%)	1 (2%)	0	100	100
1	K	70/73~(96%)	68 (97%)	2 (3%)	0	100	100
1	L	70/73 (96%)	70 (100%)	0	0	100	100
All	All	828/876 (94%)	824 (100%)	4 (0%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	66/66 (100%)	65 (98%)	1 (2%)	65 71
1	В	66/66 (100%)	66 (100%)	0	100 100



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentile	$\overline{\mathbf{s}}$
1	С	59/66 (89%)	59 (100%)	0	100 100	
1	D	61/66 (92%)	61 (100%)	0	100 100	
1	E	65/66 (98%)	65 (100%)	0	100 100	
1	F	65/66~(98%)	64 (98%)	1 (2%)	65 71	
1	G	63/66 (96%)	62 (98%)	1 (2%)	62 69	
1	Н	$64/66 \; (97\%)$	63 (98%)	1 (2%)	62 69	
1	I	57/66 (86%)	55 (96%)	2 (4%)	36 38	
1	J	63/66 (96%)	63 (100%)	0	100 100	
1	K	65/66 (98%)	64 (98%)	1 (2%)	65 71	
1	L	65/66~(98%)	64 (98%)	1 (2%)	65 71	
All	All	759/792 (96%)	751 (99%)	8 (1%)	73 79	

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Н	82	SER
1	L	84	GLU
1	I	146	LEU
1	G	129	SER
1	I	143	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	Е	132	ASN
1	F	100	GLN
1	K	98	GLN
1	E	109	GLN
1	K	132	ASN

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



# 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry (i)

There are no ligands in this entry.

### 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	73/73~(100%)	0.55	5 (6%) 17 21	31, 51, 80, 85	0
1	В	73/73 (100%)	0.84	12 (16%) 1 2	33, 53, 111, 116	0
1	С	67/73 (91%)	0.75	12 (17%) 1 1	36, 56, 96, 108	0
1	D	$69/73 \; (94\%)$	0.46	7 (10%) 7 9	31, 42, 101, 125	0
1	E	73/73 (100%)	0.66	9 (12%) 4 5	32, 47, 103, 118	0
1	F	73/73 (100%)	1.14	15 (20%) 1 1	31, 45, 105, 120	0
1	G	72/73~(98%)	1.00	15 (20%) 1 0	28, 44, 124, 133	0
1	Н	72/73~(98%)	0.85	11 (15%) 2 3	29, 47, 129, 146	0
1	I	$66/73 \; (90\%)$	0.81	9 (13%) 3 4	32, 51, 115, 123	0
1	J	70/73~(95%)	0.81	8 (11%) 5 6	29, 42, 109, 116	0
1	K	72/73~(98%)	0.63	10 (13%) 2 3	30, 49, 93, 115	0
1	L	72/73 (98%)	1.41	17 (23%) 0 0	28, 43, 101, 114	0
All	All	852/876 (97%)	0.83	130 (15%) 2 3	28, 48, 110, 146	0

The worst 5 of 130 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	87	VAL	9.3
1	L	91	ALA	9.2
1	E	87	VAL	8.1
1	L	82	SER	7.3
1	I	83	PHE	7.2

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



# 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

# 6.4 Ligands (i)

There are no ligands in this entry.

# 6.5 Other polymers (i)

There are no such residues in this entry.

