



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 20, 2023 – 12:04 AM EDT

PDB ID : 2GEZ  
Title : Crystal structure of potassium-independent plant asparaginase  
Authors : Michalska, K.; Bujacz, G.; Jaskolski, M.  
Deposited on : 2006-03-21  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	195	 3% 63% 17% 19%
1	C	195	 0% 72% 13% 15%
1	E	195	 0% 67% 16% 15%
1	G	195	 4% 71% 8% 19%
2	B	133	 2% 80% 17%
2	D	133	 2% 74% 24%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	133	 2% 83% 14%
2	H	133	 2% 83% 15%

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8815 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-asparaginase alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	158	1190	745	210	229	6	0	0	0
1	C	166	1259	787	223	243	6	0	0	0
1	E	165	1250	782	221	241	6	0	0	0
1	G	158	1190	745	210	229	6	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	cloning artifact	UNP Q9ZSD6
A	-1	SER	-	cloning artifact	UNP Q9ZSD6
A	0	HIS	-	cloning artifact	UNP Q9ZSD6
C	-2	GLY	-	cloning artifact	UNP Q9ZSD6
C	-1	SER	-	cloning artifact	UNP Q9ZSD6
C	0	HIS	-	cloning artifact	UNP Q9ZSD6
E	-2	GLY	-	cloning artifact	UNP Q9ZSD6
E	-1	SER	-	cloning artifact	UNP Q9ZSD6
E	0	HIS	-	cloning artifact	UNP Q9ZSD6
G	-2	GLY	-	cloning artifact	UNP Q9ZSD6
G	-1	SER	-	cloning artifact	UNP Q9ZSD6
G	0	HIS	-	cloning artifact	UNP Q9ZSD6

- Molecule 2 is a protein called L-asparaginase beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	133	951	593	162	189	7	0	0	0
2	D	133	951	593	162	189	7	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	133	Total	C	N	O	S	0	0	0
			951	593	162	189	7			
2	H	133	Total	C	N	O	S	0	0	0
			951	593	162	189	7			

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		
3	C	1	Total	Na	0	0
			1	1		
3	E	1	Total	Na	0	0
			1	1		
3	G	1	Total	Na	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		
4	D	2	Total	Cl	0	0
			2	2		
4	F	2	Total	Cl	0	0
			2	2		
4	H	1	Total	Cl	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	19	Total	O	0	0
			19	19		
5	B	15	Total	O	0	0
			15	15		
5	C	15	Total	O	0	0
			15	15		
5	D	13	Total	O	0	0
			13	13		
5	E	17	Total	O	0	0
			17	17		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
5	F	9	Total O 9 9	0	0
5	G	9	Total O 9 9	0	0
5	H	15	Total O 15 15	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: L-asparaginase alpha subunit




- Molecule 1: L-asparaginase alpha subunit



- Molecule 1: L-asparaginase alpha subunit





- Molecule 1: L-asparaginase alpha subunit



SER	GLN	TYR	ASN	PRO	GLU	PRO	VAL	LYS	ASP	ASP	ALA	GLU	LYS	GLU	LEU	PRO	LEU	THR	ASN	GLY	ASP	SER	GLN	ILE	GLY
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----


• Molecule 2: L-asparaginase beta subunit





• Molecule 2: L-asparaginase beta subunit



• Molecule 2: L-asparaginase beta subunit



• Molecule 2: L-asparaginase beta subunit





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.68Å 60.20Å 114.63Å 100.60° 92.90° 113.40°	Depositor
Resolution (Å)	25.00 – 2.60 24.94 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.9 (25.00-2.60) 97.4 (24.94-2.60)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.41 (at 2.60Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.189 , 0.254 0.202 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.9	Xtrriage
Anisotropy	0.843	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 46.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.054 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	8815	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.73	1/1209 (0.1%)	0.77	0/1637
1	C	0.73	0/1279	0.76	0/1732
1	E	0.79	1/1270 (0.1%)	0.84	1/1720 (0.1%)
1	G	0.72	1/1209 (0.1%)	0.77	0/1637
2	B	0.73	0/964	0.78	1/1307 (0.1%)
2	D	0.77	0/964	0.81	1/1307 (0.1%)
2	F	0.78	0/964	0.83	2/1307 (0.2%)
2	H	0.74	1/964 (0.1%)	0.81	1/1307 (0.1%)
All	All	0.75	4/8823 (0.0%)	0.79	6/11954 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	1
2	F	0	1
All	All	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	93	CYS	CB-SG	-7.79	1.69	1.82
1	E	40	GLU	CG-CD	5.87	1.60	1.51
1	G	93	CYS	CB-SG	-5.70	1.72	1.81
2	H	238	CYS	CB-SG	-5.09	1.73	1.81

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	32	ARG	NE-CZ-NH1	7.26	123.93	120.30
2	D	256	ARG	NE-CZ-NH1	6.42	123.51	120.30
2	F	200	ASP	CB-CG-OD2	6.00	123.70	118.30
2	H	256	ARG	NE-CZ-NH2	5.98	123.29	120.30
2	F	256	ARG	NE-CZ-NH1	5.55	123.08	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	F	324	THR	Peptide
1	G	139	ASP	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1190	0	1199	20	0
1	C	1259	0	1264	18	0
1	E	1250	0	1256	19	0
1	G	1190	0	1199	12	0
2	B	951	0	951	17	0
2	D	951	0	951	27	0
2	F	951	0	951	11	0
2	H	951	0	951	11	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0
3	E	1	0	0	0	0
3	G	1	0	0	0	0
4	B	1	0	0	0	0
4	D	2	0	0	1	0
4	F	2	0	0	0	0
4	H	1	0	0	0	0
5	A	19	0	0	0	0
5	B	15	0	0	0	0
5	C	15	0	0	0	0
5	D	13	0	0	0	0
5	E	17	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	F	9	0	0	0	0
5	G	9	0	0	0	0
5	H	15	0	0	0	0
All	All	8815	0	8722	110	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 110 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:116:THR:HG22	1:E:118:HIS:H	1.34	0.92
1:C:116:THR:HG22	1:C:118:HIS:H	1.41	0.84
1:A:116:THR:HG22	1:A:118:HIS:H	1.49	0.77
2:D:319:ILE:H	2:D:319:ILE:HD13	1.51	0.75
1:E:112:VAL:O	1:E:116:THR:HB	1.88	0.73

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	156/195 (80%)	149 (96%)	7 (4%)	0	100	100
1	C	164/195 (84%)	155 (94%)	9 (6%)	0	100	100
1	E	163/195 (84%)	156 (96%)	7 (4%)	0	100	100
1	G	156/195 (80%)	151 (97%)	5 (3%)	0	100	100
2	B	131/133 (98%)	125 (95%)	6 (5%)	0	100	100
2	D	131/133 (98%)	125 (95%)	6 (5%)	0	100	100
2	F	131/133 (98%)	125 (95%)	6 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	H	131/133 (98%)	127 (97%)	4 (3%)	0	100	100
All	All	1163/1312 (89%)	1113 (96%)	50 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	129/162 (80%)	112 (87%)	17 (13%)	4	7
1	C	137/162 (85%)	131 (96%)	6 (4%)	28	53
1	E	136/162 (84%)	124 (91%)	12 (9%)	10	19
1	G	129/162 (80%)	122 (95%)	7 (5%)	22	44
2	B	97/97 (100%)	91 (94%)	6 (6%)	18	37
2	D	97/97 (100%)	89 (92%)	8 (8%)	11	22
2	F	97/97 (100%)	88 (91%)	9 (9%)	9	17
2	H	97/97 (100%)	91 (94%)	6 (6%)	18	37
All	All	919/1036 (89%)	848 (92%)	71 (8%)	13	25

5 of 71 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	319	ILE
1	G	82	GLU
2	H	214	LEU
1	C	32	ARG
1	C	29	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	89	ASN
1	G	132	GLN
2	H	235	ASN
2	B	235	ASN
1	C	167	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	158/195 (81%)	0.11	6 (3%) 40 33	21, 26, 32, 36	0
1	C	166/195 (85%)	-0.23	2 (1%) 79 76	18, 25, 33, 37	0
1	E	165/195 (84%)	-0.23	2 (1%) 79 76	18, 25, 33, 36	0
1	G	158/195 (81%)	0.02	8 (5%) 28 22	21, 26, 32, 34	0
2	B	133/133 (100%)	-0.15	2 (1%) 73 70	22, 25, 30, 36	1 (0%)
2	D	133/133 (100%)	-0.14	2 (1%) 73 70	20, 26, 32, 38	0
2	F	133/133 (100%)	-0.09	3 (2%) 60 54	21, 26, 32, 39	1 (0%)
2	H	133/133 (100%)	-0.04	3 (2%) 60 54	21, 26, 31, 36	0
All	All	1179/1312 (89%)	-0.10	28 (2%) 59 53	18, 26, 32, 39	2 (0%)

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	325	THR	8.1
1	A	21	PRO	5.6
2	H	325	THR	5.0
2	D	325	THR	4.7
1	C	2	GLY	3.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NA	G	403	1/1	0.91	0.10	36,36,36,36	0
3	NA	A	401	1/1	0.95	0.11	28,28,28,28	0
4	CL	D	405	1/1	0.95	0.14	28,28,28,28	0
4	CL	F	404	1/1	0.96	0.14	27,27,27,27	0
4	CL	F	408	1/1	0.97	0.06	29,29,29,29	0
3	NA	E	402	1/1	0.98	0.02	16,16,16,16	0
4	CL	D	407	1/1	0.98	0.14	26,26,26,26	0
3	NA	C	400	1/1	0.98	0.07	16,16,16,16	0
4	CL	B	406	1/1	0.98	0.12	39,39,39,39	0
4	CL	H	409	1/1	0.98	0.15	30,30,30,30	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.