

wwPDB X-ray Structure Validation Summary Report (i)

Mar 4, 2024 – 10:59 PM EST

PDB ID : 8GQF

Title : Crystal structure of Thiolase Authors : Hong, J.; Son, H.F.; Kim, K.J.

Deposited on : 2022-08-30

Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.36

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	407	86%	7%	7%
1	В	407	85%	8%	7%
1	С	407	82%	10%	7%
1	D	407	83%	10%	7%



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 11805 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Thiolase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Λ	379	Total	С	N	О	S	0	0	0
1	A	319	2743	1713	492	519	19	U	0	
1	В	379	Total	С	N	О	S	0	0	0
1		319	2743	1713	492	519	19	U	0	
1	С	377	Total	С	N	О	S	0	0	0
1		311	2729	1705	490	516	18	U		
1	D	D 378	Total	С	N	О	S	0	0	0
1			2735	1708	491	518	18	U	U	

There are 112 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP A0A0F7QQ39
A	-18	GLY	-	expression tag	UNP A0A0F7QQ39
A	-17	SER	-	expression tag	UNP A0A0F7QQ39
A	-16	SER	-	expression tag	UNP A0A0F7QQ39
A	-15	HIS	-	expression tag	UNP A0A0F7QQ39
A	-14	HIS	-	expression tag	UNP A0A0F7QQ39
A	-13	HIS	-	expression tag	UNP A0A0F7QQ39
A	-12	HIS	-	expression tag	UNP A0A0F7QQ39
A	-11	HIS	-	expression tag	UNP A0A0F7QQ39
A	-10	HIS	-	expression tag	UNP A0A0F7QQ39
A	-9	SER	-	expression tag	UNP A0A0F7QQ39
A	-8	SER	-	expression tag	UNP A0A0F7QQ39
A	-7	GLY	-	expression tag	UNP A0A0F7QQ39
A	-6	LEU	-	expression tag	UNP A0A0F7QQ39
A	-5	VAL	-	expression tag	UNP A0A0F7QQ39
A	-4	PRO	-	expression tag	UNP A0A0F7QQ39
A	-3	ARG	-	expression tag	UNP A0A0F7QQ39
A	-2	GLY	-	expression tag	UNP A0A0F7QQ39
A	-1	SER	-	expression tag	UNP A0A0F7QQ39
A	0	HIS	-	expression tag	UNP A0A0F7QQ39
A	380	LEU	-	expression tag	UNP A0A0F7QQ39



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	381	GLU	_	expression tag	UNP A0A0F7QQ39
A	382	HIS	_	expression tag	UNP A0A0F7QQ39
A	383	HIS	-	expression tag	UNP A0A0F7QQ39
A	384	HIS	-	expression tag	UNP A0A0F7QQ39
A	385	HIS	-	expression tag	UNP A0A0F7QQ39
A	386	HIS	-	expression tag	UNP A0A0F7QQ39
A	387	HIS	-	expression tag	UNP A0A0F7QQ39
В	-19	MET	-	initiating methionine	UNP A0A0F7QQ39
В	-18	GLY	-	expression tag	UNP A0A0F7QQ39
В	-17	SER	-	expression tag	UNP A0A0F7QQ39
В	-16	SER	-	expression tag	UNP A0A0F7QQ39
В	-15	HIS	-	expression tag	UNP A0A0F7QQ39
В	-14	HIS	-	expression tag	UNP A0A0F7QQ39
В	-13	HIS	-	expression tag	UNP A0A0F7QQ39
В	-12	HIS	-	expression tag	UNP A0A0F7QQ39
В	-11	HIS	-	expression tag	UNP A0A0F7QQ39
В	-10	HIS	_	expression tag	UNP A0A0F7QQ39
В	-9	SER	-	expression tag	UNP A0A0F7QQ39
В	-8	SER	-	expression tag	UNP A0A0F7QQ39
В	-7	GLY	-	expression tag	UNP A0A0F7QQ39
В	-6	LEU	-	expression tag	UNP A0A0F7QQ39
В	-5	VAL	-	expression tag	UNP A0A0F7QQ39
В	-4	PRO	-	expression tag	UNP A0A0F7QQ39
В	-3	ARG	-	expression tag	UNP A0A0F7QQ39
В	-2	GLY	-	expression tag	UNP A0A0F7QQ39
В	-1	SER	_	expression tag	UNP A0A0F7QQ39
В	0	HIS	-	expression tag	UNP A0A0F7QQ39
В	380	LEU	-	expression tag	UNP A0A0F7QQ39
В	381	GLU	-	expression tag	UNP A0A0F7QQ39
В	382	HIS	-	expression tag	UNP A0A0F7QQ39
В	383	HIS	-	expression tag	UNP A0A0F7QQ39
В	384	HIS	-	expression tag	UNP A0A0F7QQ39
В	385	HIS	-	expression tag	UNP A0A0F7QQ39
В	386	HIS	-	expression tag	UNP A0A0F7QQ39
В	387	HIS	-	expression tag	UNP A0A0F7QQ39
С	-19	MET	-	initiating methionine	UNP A0A0F7QQ39
С	-18	GLY	-	expression tag	UNP A0A0F7QQ39
С	-17	SER	-	expression tag	UNP A0A0F7QQ39
С	-16	SER	-	expression tag	UNP A0A0F7QQ39
С	-15	HIS	-	expression tag	UNP A0A0F7QQ39
С	-14	HIS	-	expression tag	UNP A0A0F7QQ39
С	-13	HIS	-	expression tag	UNP A0A0F7QQ39



Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
С	-12	HIS	-	expression tag	UNP A0A0F7QQ39
С	-11	HIS	_	expression tag	UNP A0A0F7QQ39
С	-10	HIS	-	expression tag	UNP A0A0F7QQ39
С	-9	SER	_	expression tag	UNP A0A0F7QQ39
С	-8	SER	-	expression tag	UNP A0A0F7QQ39
С	-7	GLY	-	expression tag	UNP A0A0F7QQ39
С	-6	LEU	-	expression tag	UNP A0A0F7QQ39
С	-5	VAL	-	expression tag	UNP A0A0F7QQ39
С	-4	PRO	-	expression tag	UNP A0A0F7QQ39
С	-3	ARG	-	expression tag	UNP A0A0F7QQ39
С	-2	GLY	-	expression tag	UNP A0A0F7QQ39
С	-1	SER	-	expression tag	UNP A0A0F7QQ39
С	0	HIS	-	expression tag	UNP A0A0F7QQ39
С	380	LEU	-	expression tag	UNP A0A0F7QQ39
С	381	GLU	-	expression tag	UNP A0A0F7QQ39
С	382	HIS	-	expression tag	UNP A0A0F7QQ39
С	383	HIS	-	expression tag	UNP A0A0F7QQ39
С	384	HIS	-	expression tag	UNP A0A0F7QQ39
С	385	HIS	-	expression tag	UNP A0A0F7QQ39
С	386	HIS	-	expression tag	UNP A0A0F7QQ39
С	387	HIS	-	expression tag	UNP A0A0F7QQ39
D	-19	MET	-	initiating methionine	UNP A0A0F7QQ39
D	-18	GLY	-	expression tag	UNP A0A0F7QQ39
D	-17	SER	-	expression tag	UNP A0A0F7QQ39
D	-16	SER	_	expression tag	UNP A0A0F7QQ39
D	-15	HIS	-	expression tag	UNP A0A0F7QQ39
D	-14	HIS	_	expression tag	UNP A0A0F7QQ39
D	-13	HIS	-	expression tag	UNP A0A0F7QQ39
D	-12	HIS	-	expression tag	UNP A0A0F7QQ39
D	-11	HIS	-	expression tag	UNP A0A0F7QQ39
D	-10	HIS	-	expression tag	UNP A0A0F7QQ39
D	-9	SER	-	expression tag	UNP A0A0F7QQ39
D	-8	SER	-	expression tag	UNP A0A0F7QQ39
D	-7	GLY	-	expression tag	UNP A0A0F7QQ39
D	-6	LEU	-	expression tag	UNP A0A0F7QQ39
D	-5	VAL	-	expression tag	UNP A0A0F7QQ39
D	-4	PRO	-	expression tag	UNP A0A0F7QQ39
D	-3	ARG	-	expression tag	UNP A0A0F7QQ39
D	-2	GLY	-	expression tag	UNP A0A0F7QQ39
D	-1	SER	-	expression tag	UNP A0A0F7QQ39
D	0	HIS	-	expression tag	UNP A0A0F7QQ39
D	380	LEU	-	expression tag	tinued on ment mage



Continued	trom	mmoninonic	maaa
COHABABACA		DIEUIUU	DUIUE
0 0 1000100000			

Chain	Residue	Modelled	Actual	Comment	Reference
D	381	GLU	-	expression tag	UNP A0A0F7QQ39
D	382	HIS	-	expression tag	UNP A0A0F7QQ39
D	383	HIS	-	expression tag	UNP A0A0F7QQ39
D	384	HIS	-	expression tag	UNP A0A0F7QQ39
D	385	HIS	-	expression tag	UNP A0A0F7QQ39
D	386	HIS	-	expression tag	UNP A0A0F7QQ39
D	387	HIS	=	expression tag	UNP A0A0F7QQ39

 \bullet Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $\mathrm{C_3H_8O_3}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	В	1	Total C O 6 3 3	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	229	Total O 229 229	0	0
3	В	236	Total O 236 236	0	0
3	С	183	Total O 183 183	0	0



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	189	Total O 189 189	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Thiolase









4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	50.56Å 82.27Å 97.74Å	Depositor
a, b, c, α , β , γ	90.01° 75.00° 89.95°	Depositor
Resolution (Å)	31.49 - 1.80	Depositor
Resolution (A)	31.47 - 1.80	EDS
% Data completeness	95.8 (31.49-1.80)	Depositor
(in resolution range)	95.8 (31.47-1.80)	EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	4.29 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
P. P.	0.181 , 0.220	Depositor
R, R_{free}	0.194 , 0.229	DCC
R_{free} test set	6597 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	19.2	Xtriage
Anisotropy	0.062	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.38, 33.4	EDS
L-test for twinning ²	$< L >=0.49, < L^2>=0.32$	Xtriage
	0.021 for h,-k,h-l	
Estimated twinning fraction	0.438 for -h,k,-l	Xtriage
	$0.019 ext{ for } -h,-k,-h+l$	
F_o, F_c correlation	0.95	EDS
Total number of atoms	11805	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.82% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5	
1	A	0.80	0/2781	0.93	1/3767 (0.0%)	
1	В	0.80	0/2781	0.90	0/3767	
1	С	0.76	0/2767	0.89	1/3748 (0.0%)	
1	D	0.77	0/2773	0.88	1/3757 (0.0%)	
All	All	0.78	0/11102	0.90	3/15039 (0.0%)	

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	D	76	ARG	NE-CZ-NH2	-5.47	117.56	120.30
1	A	76	ARG	NE-CZ-NH2	-5.42	117.59	120.30
1	С	188	ARG	NE-CZ-NH1	5.18	122.89	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2743	0	2788	23	0
1	В	2743	0	2788	26	0
1	С	2729	0	2772	31	0
1	D	2735	0	2776	38	0



I 'omtamalod	trom	mmonia	maaa
Continued	11 0116	DICUIUUS	Daue
	.,	10	1

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	12	0	16	0	0
2	В	6	0	8	0	0
3	A	229	0	0	1	0
3	В	236	0	0	2	0
3	С	183	0	0	2	0
3	D	189	0	0	2	0
All	All	11805	0	11148	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 100 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:88:LEU:HD23	1:C:88:LEU:HD13	1.39	1.05
1:A:88:LEU:HD13	1:D:88:LEU:HD13	1.47	0.93
1:A:175:HIS:HD2	1:A:208:ARG:H	1.30	0.79
1:D:149:MET:SD	1:D:274:MET:CE	2.72	0.78
1:C:173:ARG:NH2	1:C:321:ASP:OD1	2.17	0.77

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	377/407 (93%)	366 (97%)	11 (3%)	0	100	100
1	В	377/407 (93%)	368 (98%)	9 (2%)	0	100	100
1	С	375/407 (92%)	360 (96%)	15 (4%)	0	100	100
1	D	376/407 (92%)	363 (96%)	13 (4%)	0	100	100
All	All	$1505/1628 \; (92\%)$	1457 (97%)	48 (3%)	0	100	100



There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Out		Percentiles
1	A	273/298 (92%)	270 (99%)	3 (1%)	73 68
1	В	273/298 (92%)	269 (98%)	4 (2%)	65 56
1	С	271/298 (91%)	268 (99%)	3 (1%)	73 68
1	D	272/298 (91%)	268 (98%)	4 (2%)	65 56
All	All	1089/1192 (91%)	1075 (99%)	14 (1%)	69 62

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	142	TYR
1	С	215	SER
1	D	293	ILE
1	D	142	TYR
1	D	274	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	\mathbf{Type}
1	D	135	ASN
1	D	175	HIS
1	D	369	GLN
1	В	175	HIS
1	В	135	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	ype Chain Res Link Bond lengths		Chain Res Link		В	ond ang	gles		
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
2	GOL	A	401	-	5,5,5	0.09	0	5,5,5	0.27	0
2	GOL	A	402	-	5,5,5	0.14	0	5,5,5	0.35	0
2	GOL	В	401	-	5,5,5	0.06	0	5,5,5	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	401	-	-	0/4/4/4	-
2	GOL	A	402	-	-	0/4/4/4	-
2	GOL	В	401	_	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	В	401	GOL	C1-C2-C3-O3
2	В	401	GOL	O1-C1-C2-C3
2	В	401	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	379/407 (93%)	-0.23	2 (0%) 91 89	11, 18, 33, 56	0
1	В	379/407 (93%)	-0.21	3 (0%) 86 84	12, 18, 32, 55	0
1	С	377/407 (92%)	-0.09	3 (0%) 86 84	13, 22, 39, 56	0
1	D	378/407 (92%)	-0.09	3 (0%) 86 84	13, 22, 39, 69	0
All	All	1513/1628 (92%)	-0.15	11 (0%) 87 86	11, 20, 37, 69	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	2	SER	4.3
1	С	3	PRO	3.2
1	В	140	GLN	3.0
1	В	4	ILE	2.9
1	С	293	ILE	2.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	GOL	A	401	6/6	0.81	0.12	30,31,34,35	0
2	GOL	В	401	6/6	0.86	0.12	28,32,34,35	0
2	GOL	A	402	6/6	0.88	0.16	28,33,39,40	0

6.5 Other polymers (i)

There are no such residues in this entry.

