



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 20, 2023 – 01:59 AM EDT

PDB ID : 2GSY
Title : The 2.6A structure of Infectious Bursal Virus Derived T=1 Particles
Authors : Garriga, D.; Querol-Audi, J.; Abaitua, F.; Saugar, I.; Pous, J.; Verdaguer, N.; Caston, J.R.; Rodriguez, J.F.
Deposited on : 2006-04-27
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	456	 11% 76% 15% • 6%
1	B	456	 12% 77% 13% • 6%
1	C	456	 13% 77% 14% • 6%
1	D	456	 14% 78% 12% • 6%
1	E	456	 7% 77% 14% • 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	456	
1	G	456	
1	H	456	
1	I	456	
1	J	456	
1	K	456	
1	L	456	
1	M	456	
1	N	456	
1	O	456	
1	P	456	
1	Q	456	
1	R	456	
1	S	456	
1	T	456	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 64447 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	428	3209	2036	533	630	10	0	0	0
1	B	428	3209	2036	533	630	10	0	0	0
1	C	428	3209	2036	533	630	10	0	0	0
1	D	428	3209	2036	533	630	10	0	0	0
1	E	433	3241	2057	538	636	10	0	0	0
1	F	433	3241	2057	538	636	10	0	0	0
1	G	433	3241	2057	538	636	10	0	0	0
1	H	428	3209	2036	533	630	10	0	0	0
1	I	428	3209	2036	533	630	10	0	0	0
1	J	428	3209	2036	533	630	10	0	0	0
1	K	428	3209	2036	533	630	10	0	0	0
1	L	428	3209	2036	533	630	10	0	0	0
1	M	428	3209	2036	533	630	10	0	0	0
1	N	428	3209	2036	533	630	10	0	0	0
1	O	428	3209	2036	533	630	10	0	0	0
1	P	430	3218	2041	535	632	10	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Q	429	Total	C	N	O	S	0	0	0
			3213	2038	534	631	10			
1	R	433	Total	C	N	O	S	0	0	0
			3241	2057	538	636	10			
1	S	428	Total	C	N	O	S	0	0	0
			3209	2036	533	630	10			
1	T	428	Total	C	N	O	S	0	0	0
			3209	2036	533	630	10			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		
2	B	1	Total	Ca	0	0
			1	1		
2	C	1	Total	Ca	0	0
			1	1		
2	D	1	Total	Ca	0	0
			1	1		
2	F	1	Total	Ca	0	0
			1	1		
2	G	1	Total	Ca	0	0
			1	1		
2	L	1	Total	Ca	0	0
			1	1		
2	M	1	Total	Ca	0	0
			1	1		
2	O	1	Total	Ca	0	0
			1	1		
2	S	1	Total	Ca	0	0
			1	1		
2	T	1	Total	Ca	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	O	0	0
			2	2		
3	B	4	Total	O	0	0
			4	4		



Continued on next page...

Continued from previous page...



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	2	Total O 2 2	0	0
3	D	2	Total O 2 2	0	0
3	E	8	Total O 8 8	0	0
3	F	4	Total O 4 4	0	0
3	G	6	Total O 6 6	0	0
3	H	7	Total O 7 7	0	0
3	I	6	Total O 6 6	0	0
3	J	1	Total O 1 1	0	0
3	K	7	Total O 7 7	0	0
3	L	6	Total O 6 6	0	0
3	M	11	Total O 11 11	0	0
3	N	5	Total O 5 5	0	0
3	O	4	Total O 4 4	0	0
3	P	5	Total O 5 5	0	0
3	Q	5	Total O 5 5	0	0
3	R	5	Total O 5 5	0	0
3	S	7	Total O 7 7	0	0
3	T	18	Total O 18 18	0	0



• Molecule 1: polyprotein




• Molecule 1: polyprotein




• Molecule 1: polyprotein


• Molecule 1: polyprotein




• Molecule 1: polyprotein





• Molecule 1: polyprotein









• Molecule 1: polyprotein




• Molecule 1: polyprotein





• Molecule 1: polyprotein









• Molecule 1: polyprotein




• Molecule 1: polyprotein





• Molecule 1: polyprotein









• Molecule 1: polyprotein




• Molecule 1: polyprotein



• Molecule 1: polyprotein





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	326.30Å 326.30Å 326.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.97 – 2.60 19.93 – 2.60	Depositor EDS
% Data completeness (in resolution range)	88.8 (19.97-2.60) 88.8 (19.93-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.19	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.27 (at 2.59Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.254 , 0.267 0.274 , 0.281	Depositor DCC
R_{free} test set	15643 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	40.6	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 12.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.011 for l,-k,h	Xtrriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	64447	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/3272	0.61	0/4471
1	B	0.39	0/3272	0.61	0/4471
1	C	0.39	0/3272	0.61	0/4471
1	D	0.38	0/3272	0.61	0/4471
1	E	0.38	0/3306	0.59	0/4519
1	F	0.39	0/3306	0.61	0/4519
1	G	0.39	0/3306	0.62	1/4519 (0.0%)
1	H	0.39	0/3272	0.61	0/4471
1	I	0.39	0/3272	0.61	0/4471
1	J	0.39	0/3272	0.61	0/4471
1	K	0.39	0/3272	0.61	0/4471
1	L	0.39	0/3272	0.61	0/4471
1	M	0.38	0/3272	0.60	0/4471
1	N	0.39	0/3272	0.61	0/4471
1	O	0.39	0/3272	0.61	0/4471
1	P	0.39	0/3281	0.61	0/4483
1	Q	0.39	0/3276	0.61	0/4476
1	R	0.38	0/3306	0.60	0/4519
1	S	0.38	0/3272	0.61	0/4471
1	T	0.40	0/3272	0.61	0/4471
All	All	0.39	0/65589	0.61	1/89629 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	3
1	C	0	3

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	3
1	E	0	3
1	F	0	3
1	G	0	3
1	H	0	3
1	I	0	3
1	J	0	3
1	K	0	3
1	L	0	3
1	M	0	3
1	N	0	3
1	O	0	3
1	P	0	3
1	Q	0	3
1	R	0	3
1	S	0	4
1	T	0	3
All	All	0	61

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	G	113	LEU	CA-CB-CG	5.69	128.39	115.30

There are no chirality outliers.

5 of 61 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	10	GLN	Peptide
1	A	430	ALA	Peptide
1	A	435	PRO	Peptide
1	B	10	GLN	Peptide
1	B	430	ALA	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3209	0	3191	48	0
1	B	3209	0	3191	46	0
1	C	3209	0	3191	50	0
1	D	3209	0	3191	44	1
1	E	3241	0	3221	54	1
1	F	3241	0	3221	55	1
1	G	3241	0	3221	53	0
1	H	3209	0	3191	49	0
1	I	3209	0	3191	55	0
1	J	3209	0	3191	49	0
1	K	3209	0	3191	55	0
1	L	3209	0	3191	48	1
1	M	3209	0	3191	55	0
1	N	3209	0	3191	55	0
1	O	3209	0	3191	54	0
1	P	3218	0	3199	63	0
1	Q	3213	0	3194	57	0
1	R	3241	0	3221	63	0
1	S	3209	0	3191	49	0
1	T	3209	0	3191	51	2
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	L	1	0	0	0	0
2	M	1	0	0	0	0
2	O	1	0	0	0	0
2	S	1	0	0	0	0
2	T	1	0	0	0	0
3	A	2	0	0	0	0
3	B	4	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
3	E	8	0	0	0	0
3	F	4	0	0	0	0
3	G	6	0	0	0	0
3	H	7	0	0	0	0
3	I	6	0	0	0	0
3	J	1	0	0	0	0
3	K	7	0	0	0	0
3	L	6	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	M	11	0	0	0	0
3	N	5	0	0	0	0
3	O	4	0	0	0	0
3	P	5	0	0	1	0
3	Q	5	0	0	0	0
3	R	5	0	0	0	0
3	S	7	0	0	0	0
3	T	18	0	0	1	0
All	All	64447	0	63951	990	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 990 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:R:110:SER:HB3	1:R:160:VAL:HG12	1.43	0.97
1:M:286:THR:HG21	1:R:279:ASN:HB2	1.44	0.96
1:R:204:ARG:HB3	1:R:337:ILE:HD13	1.56	0.88
1:G:204:ARG:HD2	1:G:337:ILE:HD11	1.56	0.87
1:P:150:ASN:HD21	1:Q:385:THR:H	1.20	0.87

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:320:GLN:OE1	1:F:217:SER:OG[3_545]	1.77	0.43
1:E:284:THR:CG2	1:T:217:SER:CB[11_456]	2.06	0.14
1:L:317:SER:O	1:T:284:THR:CG2[7_564]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	424/456 (93%)	396 (93%)	17 (4%)	11 (3%)	5	9
1	B	424/456 (93%)	394 (93%)	19 (4%)	11 (3%)	5	9
1	C	424/456 (93%)	395 (93%)	18 (4%)	11 (3%)	5	9
1	D	424/456 (93%)	396 (93%)	17 (4%)	11 (3%)	5	9
1	E	431/456 (94%)	401 (93%)	20 (5%)	10 (2%)	6	11
1	F	431/456 (94%)	401 (93%)	18 (4%)	12 (3%)	5	7
1	G	431/456 (94%)	403 (94%)	17 (4%)	11 (3%)	5	9
1	H	424/456 (93%)	395 (93%)	18 (4%)	11 (3%)	5	9
1	I	424/456 (93%)	396 (93%)	16 (4%)	12 (3%)	5	7
1	J	424/456 (93%)	396 (93%)	17 (4%)	11 (3%)	5	9
1	K	424/456 (93%)	395 (93%)	18 (4%)	11 (3%)	5	9
1	L	424/456 (93%)	396 (93%)	17 (4%)	11 (3%)	5	9
1	M	424/456 (93%)	396 (93%)	18 (4%)	10 (2%)	6	10
1	N	424/456 (93%)	395 (93%)	17 (4%)	12 (3%)	5	7
1	O	424/456 (93%)	396 (93%)	17 (4%)	11 (3%)	5	9
1	P	426/456 (93%)	397 (93%)	17 (4%)	12 (3%)	5	7
1	Q	425/456 (93%)	396 (93%)	18 (4%)	11 (3%)	5	9
1	R	431/456 (94%)	401 (93%)	23 (5%)	7 (2%)	9	19
1	S	424/456 (93%)	394 (93%)	19 (4%)	11 (3%)	5	9
1	T	424/456 (93%)	395 (93%)	18 (4%)	11 (3%)	5	9
All	All	8511/9120 (93%)	7934 (93%)	359 (4%)	218 (3%)	5	9

5 of 218 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	11	ILE
1	A	279	ASN
1	A	435	PRO
1	B	11	ILE
1	B	279	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	354/380 (93%)	328 (93%)	26 (7%)	14	28
1	B	354/380 (93%)	328 (93%)	26 (7%)	14	28
1	C	354/380 (93%)	328 (93%)	26 (7%)	14	28
1	D	354/380 (93%)	328 (93%)	26 (7%)	14	28
1	E	356/380 (94%)	332 (93%)	24 (7%)	16	33
1	F	356/380 (94%)	329 (92%)	27 (8%)	13	26
1	G	356/380 (94%)	330 (93%)	26 (7%)	14	28
1	H	354/380 (93%)	329 (93%)	25 (7%)	14	29
1	I	354/380 (93%)	328 (93%)	26 (7%)	14	28
1	J	354/380 (93%)	326 (92%)	28 (8%)	12	24
1	K	354/380 (93%)	327 (92%)	27 (8%)	13	26
1	L	354/380 (93%)	329 (93%)	25 (7%)	14	29
1	M	354/380 (93%)	328 (93%)	26 (7%)	14	28
1	N	354/380 (93%)	329 (93%)	25 (7%)	14	29
1	O	354/380 (93%)	326 (92%)	28 (8%)	12	24
1	P	354/380 (93%)	329 (93%)	25 (7%)	14	29
1	Q	354/380 (93%)	329 (93%)	25 (7%)	14	29
1	R	356/380 (94%)	332 (93%)	24 (7%)	16	33
1	S	354/380 (93%)	328 (93%)	26 (7%)	14	28
1	T	354/380 (93%)	329 (93%)	25 (7%)	14	29
All	All	7088/7600 (93%)	6572 (93%)	516 (7%)	14	28

5 of 516 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	R	407	ARG
1	S	201	ASP
1	R	396	ASN
1	H	273	THR
1	H	193	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 161 such sidechains are listed below:

Mol	Chain	Res	Type
1	O	150	ASN
1	R	288	ASN
1	P	75	GLN
1	Q	90	GLN
1	S	150	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 11 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	428/456 (93%)	0.56	52 (12%) 4 2	8, 12, 17, 24	0
1	B	428/456 (93%)	0.74	53 (12%) 4 2	8, 12, 17, 24	0
1	C	428/456 (93%)	0.70	60 (14%) 2 1	8, 12, 18, 24	0
1	D	428/456 (93%)	0.73	66 (15%) 2 1	8, 12, 16, 24	0
1	E	433/456 (94%)	0.46	31 (7%) 15 11	5, 11, 16, 24	0
1	F	433/456 (94%)	0.76	57 (13%) 3 2	8, 12, 16, 24	0
1	G	433/456 (94%)	0.60	40 (9%) 9 6	6, 12, 16, 24	0
1	H	428/456 (93%)	0.65	43 (10%) 7 4	8, 12, 18, 24	0
1	I	428/456 (93%)	0.68	50 (11%) 4 3	8, 12, 18, 24	0
1	J	428/456 (93%)	0.68	60 (14%) 2 1	8, 12, 17, 24	0
1	K	428/456 (93%)	0.63	48 (11%) 5 3	8, 12, 18, 24	0
1	L	428/456 (93%)	0.56	44 (10%) 6 4	8, 12, 18, 24	0
1	M	428/456 (93%)	0.66	42 (9%) 7 5	8, 12, 18, 24	0
1	N	428/456 (93%)	0.63	46 (10%) 6 3	8, 12, 18, 24	0
1	O	428/456 (93%)	0.66	60 (14%) 2 1	8, 12, 18, 24	0
1	P	430/456 (94%)	0.63	48 (11%) 5 3	8, 12, 16, 24	0
1	Q	429/456 (94%)	0.66	55 (12%) 3 2	8, 12, 18, 24	0
1	R	433/456 (94%)	0.54	27 (6%) 20 15	2, 11, 17, 24	0
1	S	428/456 (93%)	0.61	30 (7%) 16 12	4, 11, 18, 24	0
1	T	428/456 (93%)	0.69	33 (7%) 13 10	8, 12, 18, 24	0
All	All	8583/9120 (94%)	0.64	945 (11%) 5 3	2, 12, 18, 24	0

The worst 5 of 945 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	440	GLY	14.1
1	Q	439	ALA	12.9
1	R	439	ALA	12.5
1	N	439	ALA	11.4
1	K	439	ALA	11.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CA	A	501	1/1	0.95	0.12	10,10,10,10	1
2	CA	L	507	1/1	0.95	0.12	2,2,2,2	1
2	CA	O	509	1/1	0.95	0.19	2,2,2,2	0
2	CA	M	508	1/1	0.97	0.09	2,2,2,2	1
2	CA	D	504	1/1	0.97	0.17	2,2,2,2	1
2	CA	S	510	1/1	0.97	0.09	5,5,5,5	1
2	CA	B	502	1/1	0.98	0.19	12,12,12,12	1
2	CA	G	506	1/1	0.98	0.12	2,2,2,2	0
2	CA	F	505	1/1	0.99	0.11	2,2,2,2	0
2	CA	C	503	1/1	0.99	0.17	2,2,2,2	1
2	CA	T	511	1/1	0.99	0.13	4,4,4,4	1

6.5 Other polymers [i](#)

There are no such residues in this entry.