



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 30, 2021 – 04:29 PM EST

PDB ID : 3GV0  
Title : Crystal structure of LacI family transcription regulator from *Agrobacterium tumefaciens*  
Authors : Bagaria, A.; Burley, S.K.; Swaminathan, S.; New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : 2009-03-30  
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.16  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.16


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	288	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2185 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulator, LacI family.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	269	2097	1326	367	396	1	7	0	0	0

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	59	MSE	-	expression tag	UNP Q7CZ24
A	60	SER	-	expression tag	UNP Q7CZ24
A	61	LEU	-	expression tag	UNP Q7CZ24
A	339	GLU	-	expression tag	UNP Q7CZ24
A	340	GLY	-	expression tag	UNP Q7CZ24
A	341	HIS	-	expression tag	UNP Q7CZ24
A	342	HIS	-	expression tag	UNP Q7CZ24
A	343	HIS	-	expression tag	UNP Q7CZ24
A	344	HIS	-	expression tag	UNP Q7CZ24
A	345	HIS	-	expression tag	UNP Q7CZ24
A	346	HIS	-	expression tag	UNP Q7CZ24

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	88	Total	O	0	0
			88	88		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.71Å 86.71Å 111.22Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.36 – 2.35 43.36 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.7 (43.36-2.35) 99.7 (43.36-2.35)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.38 (at 2.34Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.198 , 0.255 0.215 , 0.256	Depositor DCC
$R_{free}$ test set	1056 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.9	Xtrriage
Anisotropy	0.169	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 44.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.025 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2185	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.45	0/2132	0.67	0/2875

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2097	0	2085	72	0
2	A	88	0	0	8	0
All	All	2185	0	2085	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:259:ILE:HD11	1:A:286:PHE:CZ	1.14	1.66
1:A:259:ILE:CD1	1:A:286:PHE:HZ	1.04	1.63
1:A:80:MSE:HE2	1:A:286:PHE:CE2	1.09	1.57
1:A:80:MSE:CE	1:A:286:PHE:HE2	1.28	1.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:286:PHE:CE1	1:A:290:ILE:HD11	1.70	1.26
1:A:80:MSE:CE	1:A:286:PHE:CE2	2.05	1.26
1:A:286:PHE:CE1	1:A:290:ILE:CD1	2.37	1.08
1:A:286:PHE:CD1	1:A:290:ILE:HD11	1.91	1.06
1:A:259:ILE:CD1	1:A:286:PHE:CZ	1.96	1.01
1:A:286:PHE:HE1	1:A:290:ILE:HD11	1.27	0.96
1:A:259:ILE:HD13	1:A:286:PHE:HZ	1.30	0.96
1:A:238:GLN:NE2	1:A:267:ALA:O	2.00	0.94
1:A:286:PHE:O	1:A:290:ILE:HG13	1.71	0.91
1:A:253:ILE:O	1:A:253:ILE:HG22	1.76	0.85
1:A:94:LEU:O	1:A:97:THR:OG1	1.95	0.83
1:A:73:LEU:O	1:A:73:LEU:HD12	1.79	0.82
1:A:189:ALA:HB1	1:A:221:ILE:HD12	1.60	0.82
1:A:286:PHE:HE1	1:A:290:ILE:CD1	1.82	0.81
1:A:72:VAL:HG22	1:A:104:THR:OG1	1.80	0.81
1:A:253:ILE:O	1:A:253:ILE:CG2	2.30	0.79
1:A:80:MSE:HE2	1:A:286:PHE:CD2	2.10	0.78
1:A:238:GLN:HE21	1:A:267:ALA:C	1.87	0.78
1:A:259:ILE:HD11	1:A:286:PHE:CE2	2.13	0.77
1:A:106:HIS:HD2	1:A:108:HIS:O	1.69	0.76
1:A:72:VAL:O	2:A:471:HOH:O	2.07	0.73
1:A:80:MSE:HE2	1:A:286:PHE:CZ	2.10	0.73
1:A:259:ILE:HD13	1:A:286:PHE:CZ	2.12	0.70
1:A:234:ARG:HD3	2:A:415:HOH:O	1.93	0.68
1:A:286:PHE:CE1	1:A:290:ILE:HD12	2.31	0.66
1:A:81:GLY:HA2	2:A:426:HOH:O	1.94	0.66
1:A:179:ARG:HH11	1:A:295:HIS:HD2	1.44	0.65
1:A:238:GLN:NE2	1:A:267:ALA:CB	2.60	0.65
1:A:234:ARG:NH1	1:A:235:ASP:OD1	2.30	0.64
1:A:238:GLN:NE2	1:A:267:ALA:HB1	2.14	0.63
1:A:179:ARG:HH11	1:A:295:HIS:CD2	2.18	0.62
1:A:319:GLU:OE1	2:A:414:HOH:O	2.16	0.61
1:A:75:VAL:O	1:A:75:VAL:HG12	2.00	0.61
1:A:238:GLN:HG3	1:A:268:ALA:HB2	1.84	0.59
1:A:75:VAL:HB	1:A:107:ILE:HD13	1.85	0.58
1:A:286:PHE:CD1	1:A:290:ILE:CD1	2.72	0.58
1:A:241:MSE:HG3	1:A:270:VAL:HG21	1.90	0.52
1:A:315:ARG:NH1	1:A:319:GLU:O	2.43	0.52
1:A:275:ASP:N	1:A:275:ASP:OD1	2.41	0.52
1:A:141:ARG:HH11	1:A:141:ARG:HG3	1.75	0.51
1:A:266:GLU:C	1:A:268:ALA:H	2.13	0.51

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:179:ARG:HD2	1:A:295:HIS:CD2	2.46	0.50
1:A:240:LEU:HG	1:A:241:MSE:CE	2.42	0.49
1:A:238:GLN:HE22	1:A:267:ALA:HB1	1.77	0.49
1:A:306:ARG:O	1:A:310:LYS:HG3	2.12	0.49
1:A:196:ARG:HD2	2:A:409:HOH:O	2.12	0.49
1:A:189:ALA:HB1	1:A:221:ILE:CD1	2.36	0.48
1:A:241:MSE:HE2	1:A:241:MSE:HA	1.97	0.47
1:A:106:HIS:CD2	1:A:108:HIS:O	2.59	0.47
1:A:85:GLN:NE2	2:A:403:HOH:O	2.34	0.47
1:A:234:ARG:NH1	2:A:415:HOH:O	2.41	0.47
1:A:262:VAL:HG21	1:A:294:ILE:HD11	1.98	0.45
1:A:164:PHE:CD1	1:A:164:PHE:C	2.90	0.45
1:A:119:ILE:O	1:A:123:GLY:N	2.49	0.44
1:A:287:LEU:HA	1:A:287:LEU:HD23	1.86	0.44
1:A:72:VAL:HG22	1:A:104:THR:HG1	1.78	0.44
1:A:221:ILE:O	1:A:221:ILE:HG22	2.19	0.43
1:A:118:TYR:O	1:A:122:THR:HB	2.19	0.43
1:A:170:GLU:OE2	1:A:209:ARG:NH1	2.50	0.43
1:A:141:ARG:NH1	1:A:141:ARG:HG3	2.33	0.42
1:A:80:MSE:SE	1:A:286:PHE:HE2	2.44	0.42
1:A:196:ARG:CD	2:A:409:HOH:O	2.67	0.42
1:A:122:THR:HG23	1:A:122:THR:O	2.20	0.42
1:A:195:SER:HA	1:A:200:HIS:CG	2.54	0.42
1:A:240:LEU:O	1:A:241:MSE:HE2	2.20	0.42
1:A:286:PHE:HD1	1:A:290:ILE:HD11	1.69	0.41
1:A:72:VAL:HG11	1:A:115:PRO:HB2	2.03	0.40
1:A:109:ALA:HB1	1:A:196:ARG:HH12	1.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	267/288 (93%)	261 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	228/235 (97%)	214 (94%)	14 (6%)	18	20

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	65	LYS
1	A	122	THR
1	A	132	LYS
1	A	231	GLU
1	A	234	ARG
1	A	272	ILE
1	A	274	GLU
1	A	275	ASP
1	A	283	SER
1	A	285	GLU
1	A	286	PHE
1	A	301	ILE
1	A	302	LYS
1	A	313	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	85	GLN
1	A	106	HIS
1	A	238	GLN
1	A	295	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	317	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	262/288 (90%)	0.08	14 (5%) 26 38	28, 42, 70, 80	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	286	PHE	5.5
1	A	223	ALA	3.6
1	A	290	ILE	3.5
1	A	244	SER	3.2
1	A	72	VAL	3.1
1	A	79	LEU	2.9
1	A	83	THR	2.6
1	A	124	SER	2.4
1	A	73	LEU	2.4
1	A	75	VAL	2.3
1	A	239	ARG	2.3
1	A	242	GLN	2.1
1	A	103	VAL	2.1
1	A	74	SER	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.