



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 8, 2024 – 01:47 pm GMT

PDB ID : 6HBD  
Title : Crystal structure of MSMEG\_1712 from Mycobacterium smegmatis in complex with Beta-D-Galactofuranose  
Authors : Li, M.; Mueller, C.; Einsle, O.; Jessen-Trefzer, C.  
Deposited on : 2018-08-10  
Resolution : 2.44 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1564 (2.46-2.42)
Clashscore	141614	1631 (2.46-2.42)
Ramachandran outliers	138981	1617 (2.46-2.42)
Sidechain outliers	138945	1617 (2.46-2.42)
RSRZ outliers	127900	1547 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	320	 6% 85% 10% • 5%
1	B	320	 11% 81% 13% • 5%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9099 atoms, of which 4457 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ABC transporter periplasmic-binding protein YtfQ.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	305	4519	1429	2232	399	453	6	12	0	0
1	B	305	4485	1428	2201	399	451	6	30	0	0

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP A0QT50
A	308	LYS	-	expression tag	UNP A0QT50
A	309	LEU	-	expression tag	UNP A0QT50
A	310	ALA	-	expression tag	UNP A0QT50
A	311	ALA	-	expression tag	UNP A0QT50
A	312	ALA	-	expression tag	UNP A0QT50
A	313	LEU	-	expression tag	UNP A0QT50
A	314	GLU	-	expression tag	UNP A0QT50
A	315	HIS	-	expression tag	UNP A0QT50
A	316	HIS	-	expression tag	UNP A0QT50
A	317	HIS	-	expression tag	UNP A0QT50
A	318	HIS	-	expression tag	UNP A0QT50
A	319	HIS	-	expression tag	UNP A0QT50
A	320	HIS	-	expression tag	UNP A0QT50
B	1	MET	-	initiating methionine	UNP A0QT50
B	308	LYS	-	expression tag	UNP A0QT50
B	309	LEU	-	expression tag	UNP A0QT50
B	310	ALA	-	expression tag	UNP A0QT50
B	311	ALA	-	expression tag	UNP A0QT50
B	312	ALA	-	expression tag	UNP A0QT50
B	313	LEU	-	expression tag	UNP A0QT50
B	314	GLU	-	expression tag	UNP A0QT50
B	315	HIS	-	expression tag	UNP A0QT50
B	316	HIS	-	expression tag	UNP A0QT50
B	317	HIS	-	expression tag	UNP A0QT50

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	318	HIS	-	expression tag	UNP A0QT50
B	319	HIS	-	expression tag	UNP A0QT50
B	320	HIS	-	expression tag	UNP A0QT50

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	9	Total Zn 9 9	0	0
2	B	6	Total Zn 6 6	0	0

- Molecule 3 is beta-D-galactofuranose (three-letter code: GZL) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C H O 24 6 12 6	0	0
3	B	1	Total C H O 24 6 12 6	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	20	Total O 20 20	0	0

Continued on next page...


*Continued from previous page...*


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	B	12	Total	O	0	0
			12	12		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ABC transporter periplasmic-binding protein YtfQ

Chain A: 



- Molecule 1: ABC transporter periplasmic-binding protein YtfQ

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	118.30Å 118.30Å 231.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	105.31 – 2.44 105.31 – 2.45	Depositor EDS
% Data completeness (in resolution range)	58.3 (105.31-2.44) 58.3 (105.31-2.45)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.51 (at 2.45Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, $R_{free}$	0.227 , 0.266 0.233 , 0.272	Depositor DCC
$R_{free}$ test set	872 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	70.8	Xtrriage
Anisotropy	0.069	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 69.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	9099	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	76.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GZL, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/2328	0.70	0/3157
1	B	0.50	0/2325	0.72	0/3153
All	All	0.51	0/4653	0.71	0/6310

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2287	2232	2237	7	0
1	B	2284	2201	2235	9	0
2	A	9	0	0	0	0
2	B	6	0	0	0	0
3	A	12	12	9	0	0
3	B	12	12	9	0	0
4	A	20	0	0	0	0
4	B	12	0	0	0	0
All	All	4642	4457	4490	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.



All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:70:ILE:HD11	1:B:94:THR:HG22	1.65	0.76
1:B:220:ALA:O	1:B:224:ILE:HG23	2.03	0.58
1:A:181:ALA:HB1	1:A:199:PHE:CE1	2.40	0.56
1:B:179:ILE:HG23	1:B:179:ILE:O	2.11	0.51
1:B:25:VAL:HG12	1:B:26:GLY:H	1.77	0.49
1:A:149:VAL:CG1	1:A:199:PHE:CD2	2.96	0.49
1:B:237:ILE:HG23	1:B:255:PHE:HA	1.95	0.48
1:A:36:THR:HG23	1:A:52:PHE:CZ	2.50	0.47
1:B:36:THR:HG23	1:B:52:PHE:CZ	2.50	0.47
1:B:74:VAL:HG23	1:B:99:ILE:HD11	1.98	0.45
1:A:90:VAL:O	1:A:94:THR:HG23	2.16	0.45
1:A:241:ASP:O	1:A:242:ALA:HB3	2.16	0.44
1:A:136:ALA:HA	1:A:177:LEU:HD11	1.99	0.44
1:B:136:ALA:HA	1:B:177:LEU:HD11	2.00	0.44
1:A:250:LEU:HD22	1:A:258:ILE:HG13	2.02	0.42
1:B:250:LEU:HD22	1:B:258:ILE:HG13	2.02	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	303/320 (95%)	281 (93%)	18 (6%)	4 (1%)	12	11
1	B	303/320 (95%)	283 (93%)	12 (4%)	8 (3%)	5	2
All	All	606/640 (95%)	564 (93%)	30 (5%)	12 (2%)	7	5

All (12) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	241	ASP
1	A	303	PRO
1	B	56	ASN
1	B	241	ASP
1	B	303	PRO
1	B	204	PRO
1	B	110	THR
1	A	204	PRO
1	A	81	PRO
1	B	25	VAL
1	B	81	PRO
1	B	266	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233/242 (96%)	215 (92%)	18 (8%)	13	15
1	B	232/242 (96%)	208 (90%)	24 (10%)	7	7
All	All	465/484 (96%)	423 (91%)	42 (9%)	9	10

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	84	ARG
1	A	90	VAL
1	A	104	THR
1	A	115	VAL
1	A	120	ILE
1	A	126	GLU
1	A	171	ILE
1	A	183	GLN
1	A	208	VAL
1	A	209	VAL
1	A	213	ASN
1	A	215	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	229	LYS
1	A	238	VAL
1	A	247	MET
1	A	287	VAL
1	A	305	ARG
1	A	313	LEU
1	B	28	GLU
1	B	32	ARG
1	B	72	GLN
1	B	110	THR
1	B	116	TYR
1	B	143	THR
1	B	150	GLN
1	B	183	GLN
1	B	193	LYS
1	B	194	GLN
1	B	205	GLN
1	B	206	ILE
1	B	213	ASN
1	B	215	ASP
1	B	221	MET
1	B	224	ILE
1	B	225	GLU
1	B	230	LYS
1	B	235	ILE
1	B	237	ILE
1	B	276	VAL
1	B	285	ARG
1	B	305	ARG
1	B	313	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 15 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	GZL	B	407	-	12,12,12	0.21	0	16,17,17	0.85	1 (6%)
3	GZL	A	410	-	12,12,12	0.31	0	16,17,17	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GZL	B	407	-	-	1/6/22/22	0/1/1/1
3	GZL	A	410	-	-	0/6/22/22	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	407	GZL	C1-C2-C3	-2.37	99.33	102.30

There are no chirality outliers.


All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	407	GZL	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	305/320 (95%)	0.71	18 (5%) 22 18	40, 66, 90, 101	1 (0%)
1	B	305/320 (95%)	0.88	36 (11%) 4 3	43, 90, 116, 127	3 (0%)
All	All	610/640 (95%)	0.79	54 (8%) 9 7	40, 75, 111, 127	4 (0%)

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	107	ALA	5.0
1	B	103	LEU	4.1
1	B	148	ILE	4.1
1	B	151	LEU	4.0
1	B	77	ILE	3.8
1	B	66	ILE	3.7
1	A	120	ILE	3.7
1	B	91	LEU	3.5
1	A	115	VAL	3.5
1	A	116	TYR	3.4
1	B	63	ILE	3.3
1	B	102	ILE	3.2
1	A	90	VAL	3.1
1	B	226	ALA	3.1
1	B	225	GLU	3.0
1	A	39	ILE	3.0
1	B	147	ASN	3.0
1	B	115	VAL	2.9
1	B	146	VAL	2.9
1	A	119	PHE	2.9
1	B	36	THR	2.9
1	B	181	ALA	2.9
1	A	282	VAL	2.8
1	B	237	ILE	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	63	ILE	2.7
1	A	179	ILE	2.7
1	B	272	LEU	2.7
1	B	201	LYS	2.6
1	A	286	VAL	2.5
1	B	302	LEU	2.5
1	A	171	ILE	2.5
1	A	103	LEU	2.5
1	B	175	PRO	2.5
1	B	50	LEU	2.5
1	A	109	ASP	2.5
1	B	177	LEU	2.4
1	A	82	VAL	2.4
1	B	20	LEU	2.4
1	B	69	PHE	2.4
1	A	84	ARG	2.4
1	B	229	LYS	2.4
1	B	240	VAL	2.3
1	B	259	VAL	2.3
1	A	135	VAL	2.3
1	B	150	GLN	2.3
1	B	46	ALA	2.2
1	B	179	ILE	2.2
1	B	52	PHE	2.2
1	B	276	VAL	2.1
1	B	234	ASP	2.1
1	B	149	VAL	2.1
1	A	79	PHE	2.1
1	B	199	PHE	2.1
1	A	284	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.



## 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ZN	A	409	1/1	0.91	0.14	160,160,160,160	0
2	ZN	B	401	1/1	0.92	0.18	148,148,148,148	0
2	ZN	A	404	1/1	0.95	0.16	87,87,87,87	0
3	GZL	B	407	12/12	0.95	0.15	54,60,66,69	0
3	GZL	A	410	12/12	0.96	0.16	46,47,50,51	0
2	ZN	A	408	1/1	0.97	0.29	110,110,110,110	0
2	ZN	B	402	1/1	0.98	0.25	52,52,52,52	0
2	ZN	B	404	1/1	0.98	0.25	65,65,65,65	0
2	ZN	A	407	1/1	0.98	0.25	59,59,59,59	0
2	ZN	A	405	1/1	0.98	0.27	67,67,67,67	0
2	ZN	B	403	1/1	0.99	0.26	57,57,57,57	0
2	ZN	A	406	1/1	0.99	0.23	69,69,69,69	0
2	ZN	B	405	1/1	0.99	0.26	65,65,65,65	0
2	ZN	B	406	1/1	0.99	0.27	63,63,63,63	0
2	ZN	A	401	1/1	0.99	0.27	48,48,48,48	0
2	ZN	A	403	1/1	0.99	0.20	93,93,93,93	0
2	ZN	A	402	1/1	1.00	0.28	44,44,44,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around GZL B 407:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.