



# Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 01:14 pm BST

PDB ID : 4HJH  
Title : Iodide SAD phased crystal structure of a phosphoglucomutase from *Brucella melitensis* complexed with glucose-6-phosphate  
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)  
Deposited on : 2012-10-12  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	481	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 92%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 5%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">2%      92%      5% •</p>
1	B	481	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 91%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 5%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">3%      91%      5% •</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	IOD	A	516	-	-	X	-
2	IOD	A	534	-	-	X	-
2	IOD	A	536	-	-	X	-
2	IOD	B	527	-	-	X	-
3	G6Q	B	533	-	X	-	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7469 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphomannomutase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	467	Total 3422	C 2161	N 594	O 657	P 1	S 9	0	7	0
1	B	464	Total 3302	C 2083	N 570	O 638	P 1	S 10	0	1	0


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	EXPRESSION TAG	UNP D0B6G5
A	-2	PRO	-	EXPRESSION TAG	UNP D0B6G5
B	-3	GLY	-	EXPRESSION TAG	UNP D0B6G5
B	-2	PRO	-	EXPRESSION TAG	UNP D0B6G5

- Molecule 2 is IODIDE ION (three-letter code: IOD) (formula: I).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	32	Total 32	I 32	0	0
2	A	43	Total 43	I 43	0	0

- Molecule 3 is GLUCOSE-6-PHOSPHATE (three-letter code: G6Q) (formula: C<sub>6</sub>H<sub>13</sub>O<sub>9</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
3	A	1	Total	C	O	P	0	0
			16	6	9	1		
3	B	1	Total	C	O	P	0	0
			16	6	9	1		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Mg 1 1	0	0
5	A	1	Total Mg 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	340	Total O 340 340	0	0
6	B	264	Total O 264 264	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphomannomutase



- Molecule 1: Phosphomannomutase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.95Å 55.98Å 131.67Å 90.00° 92.18° 90.00°	Depositor
Resolution (Å)	46.01 – 2.10 45.96 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.01-2.10) 99.8 (45.96-2.10)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.91 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.174 , 0.209 0.181 , 0.210	Depositor DCC
$R_{free}$ test set	2695 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.9	Xtrriage
Anisotropy	0.264	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 56.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.027 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7469	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IOD, MG, G6Q, EDO, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.55	0/3491	0.71	4/4756 (0.1%)
1	B	0.46	0/3351	0.67	1/4576 (0.0%)
All	All	0.51	0/6842	0.69	5/9332 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	382	ARG	NE-CZ-NH1	6.44	123.52	120.30
1	A	382[A]	ARG	NE-CZ-NH1	5.31	122.95	120.30
1	A	382[B]	ARG	NE-CZ-NH1	5.31	122.95	120.30
1	A	192	TYR	CA-CB-CG	5.07	123.03	113.40
1	A	371	ARG	NE-CZ-NH2	-5.06	117.77	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	146	HIS	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3422	0	3359	18	0
1	B	3302	0	3173	14	0
2	A	43	0	0	17	0
2	B	32	0	0	9	0
3	A	16	0	11	5	0
3	B	16	0	11	3	0
4	A	28	0	42	2	0
4	B	4	0	6	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	340	0	0	9	0
6	B	264	0	0	3	0
All	All	7469	0	6602	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:110:ARG:HD3	2:A:516:IOD:I	1.86	1.44
1:A:110:ARG:NH1	2:A:516:IOD:I	2.32	1.32
1:B:361:GLU:OE1	2:B:527:IOD:I	2.20	1.28
1:A:110:ARG:CD	2:A:516:IOD:I	2.74	1.05
4:A:548:EDO:H12	6:A:714:HOH:O	1.66	0.94
2:A:539:IOD:I	6:A:931:HOH:O	2.59	0.90
2:A:534:IOD:I	6:A:883:HOH:O	2.67	0.81
2:A:534:IOD:I	6:A:674:HOH:O	2.70	0.79
3:B:533:G6Q:H62	3:B:533:G6Q:O1	1.84	0.77
2:A:541:IOD:I	6:A:907:HOH:O	2.74	0.75
2:A:514:IOD:I	6:A:924:HOH:O	2.77	0.72
1:A:139:ALA:O	1:A:141:LEU:N	2.24	0.71
2:A:531:IOD:I	6:A:870:HOH:O	2.82	0.67
1:A:110:ARG:CZ	2:A:516:IOD:I	3.12	0.66
2:B:504:IOD:I	6:B:693:HOH:O	2.85	0.65

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:134:TYR:OH	2:B:519:IOD:I	2.80	0.64
3:A:544:G6Q:O1	3:A:544:G6Q:C5	2.46	0.63
3:A:544:G6Q:O1	3:A:544:G6Q:C6	2.48	0.61
1:A:294:LEU:HD21	2:A:536:IOD:I	2.72	0.60
1:A:455[B]:GLN:NE2	6:A:926:HOH:O	2.24	0.60
2:A:508:IOD:I	6:A:828:HOH:O	2.86	0.59
1:A:247:ARG:HD3	1:A:324:ASN:HB2	1.85	0.59
1:B:56:PRO:HG3	2:B:522:IOD:I	2.73	0.58
3:A:544:G6Q:H62	3:A:544:G6Q:O1	2.05	0.57
1:B:247:ARG:HD3	1:B:324:ASN:HB2	1.86	0.57
1:B:152:THR:HA	2:B:531:IOD:I	2.76	0.55
3:B:533:G6Q:C6	3:B:533:G6Q:O1	2.54	0.54
1:A:15:GLY:HA2	2:A:516:IOD:I	2.78	0.54
1:B:294:LEU:HD21	2:B:512:IOD:I	2.79	0.53
1:B:322:GLU:OE2	3:B:533:G6Q:O4	2.27	0.52
1:A:428:GLN:HG2	2:A:542:IOD:I	2.80	0.52
1:A:322:GLU:OE2	3:A:544:G6Q:O4	2.30	0.49
1:A:7:LYS:HE3	2:A:515:IOD:I	2.83	0.49
3:A:544:G6Q:O1	3:A:544:G6Q:O5	2.27	0.48
1:B:152:THR:HG23	1:B:154:ALA:HB3	1.95	0.48
1:B:92:MET:HE1	1:B:343:LEU:HD12	1.97	0.46
1:A:334:GLU:HB3	4:A:549:EDO:H22	1.98	0.46
1:A:92:MET:SD	1:A:120:GLY:HA2	2.55	0.46
1:A:159:TYR:CE2	1:A:349[A]:LEU:HD22	2.51	0.45
1:B:361:GLU:HB2	2:B:527:IOD:I	2.86	0.45
1:A:294:LEU:CD2	2:A:536:IOD:I	3.35	0.45
1:B:84:THR:N	1:B:85:PRO:HD2	2.32	0.44
1:B:382:ARG:NH2	6:B:773:HOH:O	2.49	0.44
1:A:84:THR:N	1:A:85:PRO:HD2	2.33	0.43
2:B:513:IOD:I	6:B:819:HOH:O	2.92	0.42
1:A:118:ARG:HD3	2:A:518:IOD:I	2.89	0.42
1:B:180:TYR:HB3	1:B:240:SER:HB2	2.02	0.42
1:B:22:GLY:HA2	2:B:524:IOD:I	2.90	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	467/481 (97%)	461 (99%)	5 (1%)	1 (0%)	47	49
1	B	458/481 (95%)	450 (98%)	5 (1%)	3 (1%)	22	18
All	All	925/962 (96%)	911 (98%)	10 (1%)	4 (0%)	34	32

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	146	HIS
1	B	147	VAL
1	A	148	GLY
1	B	182	HIS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	338/371 (91%)	332 (98%)	6 (2%)	59	65
1	B	315/371 (85%)	309 (98%)	6 (2%)	57	63
All	All	653/742 (88%)	641 (98%)	12 (2%)	62	65

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	286	LEU
1	A	331	SER

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type
1	A	382[A]	ARG
1	A	382[B]	ARG
1	A	406[A]	SER
1	A	406[B]	SER
1	B	42	LEU
1	B	82	LEU
1	B	286	LEU
1	B	331	SER
1	B	382	ARG
1	B	442	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	385	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	SEP	B	104	1,5	8,9,10	0.83	0	8,12,14	2.15	1 (12%)
1	SEP	A	104	1,5	8,9,10	0.90	0	8,12,14	2.02	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	B	104	1,5	-	1/5/8/10	-
1	SEP	A	104	1,5	-	1/5/8/10	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	104	SEP	OG-CB-CA	5.46	113.46	108.14
1	A	104	SEP	OG-CB-CA	4.96	112.98	108.14

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	104	SEP	N-CA-CB-OG
1	A	104	SEP	N-CA-CB-OG

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 87 ligands modelled in this entry, 77 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	EDO	A	545	-	3,3,3	0.59	0	2,2,2	0.07	0
4	EDO	A	549	-	3,3,3	0.44	0	2,2,2	0.40	0
3	G6Q	A	544	-	14,15,15	0.78	0	20,21,21	2.93	10 (50%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	G6Q	B	533	-	14,15,15	0.80	0	20,21,21	2.34	10 (50%)
4	EDO	A	550	-	3,3,3	0.36	0	2,2,2	0.54	0
4	EDO	A	551	-	3,3,3	0.49	0	2,2,2	0.09	0
4	EDO	A	548	-	3,3,3	0.51	0	2,2,2	0.20	0
4	EDO	A	547	-	3,3,3	0.38	0	2,2,2	0.71	0
4	EDO	A	546	-	3,3,3	0.64	0	2,2,2	0.23	0
4	EDO	B	534	-	3,3,3	0.50	0	2,2,2	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	545	-	-	1/1/1/1	-
4	EDO	A	549	-	-	1/1/1/1	-
3	G6Q	A	544	-	-	12/18/20/20	-
3	G6Q	B	533	-	-	14/18/20/20	-
4	EDO	A	550	-	-	0/1/1/1	-
4	EDO	A	551	-	-	1/1/1/1	-
4	EDO	A	548	-	-	1/1/1/1	-
4	EDO	A	547	-	-	1/1/1/1	-
4	EDO	A	546	-	-	1/1/1/1	-
4	EDO	B	534	-	-	1/1/1/1	-

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	544	G6Q	C4-C3-C2	-7.32	100.75	113.54
3	B	533	G6Q	C4-C3-C2	-5.13	104.59	113.54
3	A	544	G6Q	C5-C4-C3	4.22	119.07	112.47
3	A	544	G6Q	O2-C2-C3	4.11	119.22	109.46
3	B	533	G6Q	O2-C2-C3	3.85	118.60	109.46
3	A	544	G6Q	O4-C4-C5	-3.80	99.62	108.81
3	B	533	G6Q	C5-C4-C3	3.57	118.06	112.47
3	A	544	G6Q	O2-C2-C1	3.54	118.44	110.08
3	A	544	G6Q	O2P-P-O6	-3.15	98.35	106.73
3	B	533	G6Q	O6-C6-C5	3.11	117.68	109.36
3	A	544	G6Q	O6-C6-C5	3.09	117.61	109.36
3	A	544	G6Q	C6-C5-C4	3.04	118.07	112.20

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	533	G6Q	C6-C5-C4	2.70	117.42	112.20
3	B	533	G6Q	O2P-P-O6	-2.66	99.64	106.73
3	B	533	G6Q	C3-C2-C1	-2.31	103.84	111.10
3	B	533	G6Q	O4-C4-C5	-2.24	103.40	108.81
3	A	544	G6Q	C3-C2-C1	-2.21	104.15	111.10
3	A	544	G6Q	O5-C5-C4	2.12	114.26	109.10
3	B	533	G6Q	O3-C3-C2	2.06	112.95	109.17
3	B	533	G6Q	O3-C3-C4	2.00	114.14	109.47

There are no chirality outliers.

All (33) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	533	G6Q	C1-C2-C3-C4
3	B	533	G6Q	C1-C2-C3-O3
3	B	533	G6Q	O2-C2-C3-C4
3	B	533	G6Q	O2-C2-C3-O3
3	B	533	G6Q	O3-C3-C4-C5
3	B	533	G6Q	C3-C4-C5-C6
3	B	533	G6Q	C4-C5-C6-O6
3	A	544	G6Q	C1-C2-C3-C4
3	A	544	G6Q	O2-C2-C3-C4
3	A	544	G6Q	O2-C2-C3-O3
3	A	544	G6Q	C3-C4-C5-C6
3	A	544	G6Q	C3-C4-C5-O5
3	A	544	G6Q	O4-C4-C5-C6
3	A	544	G6Q	C4-C5-C6-O6
3	A	544	G6Q	O5-C5-C6-O6
3	A	544	G6Q	O4-C4-C5-O5
3	B	533	G6Q	C3-C4-C5-O5
3	B	533	G6Q	O4-C4-C5-C6
3	B	533	G6Q	C2-C3-C4-C5
4	A	551	EDO	O1-C1-C2-O2
3	A	544	G6Q	C1-C2-C3-O3
3	A	544	G6Q	O3-C3-C4-C5
3	B	533	G6Q	O4-C4-C5-O5
4	B	534	EDO	O1-C1-C2-O2
4	A	547	EDO	O1-C1-C2-O2
4	A	548	EDO	O1-C1-C2-O2
3	A	544	G6Q	C2-C3-C4-C5
3	B	533	G6Q	C2-C3-C4-O4
4	A	549	EDO	O1-C1-C2-O2

*Continued on next page...*



*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
4	A	546	EDO	O1-C1-C2-O2
4	A	545	EDO	O1-C1-C2-O2
3	B	533	G6Q	O3-C3-C4-O4
3	B	533	G6Q	O5-C5-C6-O6


There are no ring outliers.

4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	549	EDO	1	0
3	A	544	G6Q	5	0
3	B	533	G6Q	3	0
4	A	548	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	466/481 (96%)	-0.16	8 (1%) 70 74	5, 14, 37, 51	0
1	B	463/481 (96%)	-0.04	13 (2%) 53 59	11, 22, 44, 62	0
All	All	929/962 (96%)	-0.10	21 (2%) 60 65	5, 18, 42, 62	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	139	ALA	4.6
1	A	147	VAL	4.5
1	A	139	ALA	4.5
1	B	39	LYS	3.9
1	B	474	ALA	3.8
1	A	148	GLY	3.7
1	B	147	VAL	3.6
1	B	134	TYR	3.6
1	B	38	ALA	3.3
1	B	18	VAL	3.2
1	B	41	GLN	3.1
1	A	40	GLY	2.8
1	B	475	THR	2.7
1	A	311	SER	2.4
1	B	40	GLY	2.4
1	B	473	ASP	2.3
1	A	150	THR	2.3
1	B	130	ILE	2.2
1	B	37	ALA	2.2
1	A	141	LEU	2.1
1	A	146	HIS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	SEP	B	104	10/11	0.97	0.09	19,20,21,21	1
1	SEP	A	104	10/11	0.97	0.09	11,12,12,13	1

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	IOD	B	520	1/1	0.51	0.28	73,73,73,73	1
2	IOD	B	531	1/1	0.57	0.15	66,66,66,66	1
2	IOD	A	538	1/1	0.80	0.18	50,50,50,50	1
4	EDO	A	546	4/4	0.82	0.16	24,27,29,33	0
2	IOD	A	536	1/1	0.84	0.12	50,50,50,50	1
2	IOD	A	537	1/1	0.84	0.21	36,36,36,36	1
4	EDO	A	549	4/4	0.84	0.26	24,27,28,32	0
2	IOD	A	517	1/1	0.85	0.10	54,54,54,54	1
2	IOD	A	511	1/1	0.87	0.10	56,56,56,56	0
2	IOD	A	525	1/1	0.88	0.14	48,48,48,48	1
2	IOD	B	506	1/1	0.88	0.09	53,53,53,53	0
4	EDO	B	534	4/4	0.89	0.20	35,36,38,39	0
2	IOD	A	503	1/1	0.89	0.09	47,47,47,47	0
2	IOD	B	512	1/1	0.89	0.09	68,68,68,68	1
2	IOD	A	543	1/1	0.90	0.08	57,57,57,57	1
2	IOD	B	532	1/1	0.90	0.09	50,50,50,50	1
2	IOD	B	515	1/1	0.90	0.14	40,40,40,40	1
4	EDO	A	551	4/4	0.90	0.17	30,31,32,32	0
2	IOD	A	540	1/1	0.90	0.17	38,38,38,38	1
2	IOD	A	539	1/1	0.91	0.15	51,51,51,51	1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	IOD	A	519	1/1	0.91	0.07	39,39,39,39	1
3	G6Q	B	533	16/16	0.91	0.22	26,32,37,39	0
2	IOD	B	527	1/1	0.91	0.15	47,47,47,47	1
2	IOD	A	530	1/1	0.91	0.09	54,54,54,54	1
2	IOD	A	532	1/1	0.92	0.24	33,33,33,33	1
4	EDO	A	547	4/4	0.92	0.16	21,21,21,21	0
2	IOD	A	534	1/1	0.92	0.20	25,25,25,25	1
2	IOD	A	527	1/1	0.92	0.15	37,37,37,37	1
2	IOD	A	518	1/1	0.92	0.11	43,43,43,43	1
2	IOD	B	525	1/1	0.93	0.11	59,59,59,59	1
2	IOD	A	521	1/1	0.93	0.23	39,39,39,39	1
2	IOD	B	509	1/1	0.93	0.13	41,41,41,41	1
4	EDO	A	545	4/4	0.94	0.13	17,19,20,20	0
2	IOD	A	516	1/1	0.94	0.16	36,36,36,36	1
2	IOD	A	529	1/1	0.94	0.11	40,40,40,40	1
2	IOD	B	517	1/1	0.94	0.07	39,39,39,39	1
3	G6Q	A	544	16/16	0.94	0.14	16,21,23,24	0
4	EDO	A	548	4/4	0.94	0.15	16,18,19,22	0
2	IOD	A	523	1/1	0.94	0.10	43,43,43,43	1
2	IOD	A	535	1/1	0.95	0.16	66,66,66,66	1
2	IOD	B	502	1/1	0.95	0.06	44,44,44,44	1
2	IOD	A	528	1/1	0.95	0.15	30,30,30,30	1
2	IOD	A	504	1/1	0.95	0.14	35,35,35,35	1
2	IOD	B	526	1/1	0.95	0.10	47,47,47,47	1
2	IOD	A	526	1/1	0.95	0.14	45,45,45,45	1
2	IOD	B	524	1/1	0.95	0.07	48,48,48,48	1
2	IOD	A	531	1/1	0.95	0.13	45,45,45,45	1
2	IOD	A	524	1/1	0.96	0.13	57,57,57,57	1
2	IOD	A	541	1/1	0.96	0.09	35,35,35,35	1
2	IOD	B	529	1/1	0.96	0.11	31,31,31,31	1
2	IOD	B	519	1/1	0.96	0.10	38,38,38,38	1
2	IOD	B	530	1/1	0.96	0.06	49,49,49,49	1
2	IOD	B	522	1/1	0.96	0.19	33,33,33,33	1
2	IOD	A	522	1/1	0.97	0.16	34,34,34,34	1
2	IOD	A	520	1/1	0.97	0.17	29,29,29,29	1
2	IOD	A	515	1/1	0.97	0.07	47,47,47,47	1
2	IOD	B	518	1/1	0.97	0.08	32,32,32,32	1
2	IOD	B	508	1/1	0.97	0.10	46,46,46,46	1
2	IOD	B	516	1/1	0.97	0.05	34,34,34,34	1
2	IOD	B	528	1/1	0.97	0.07	47,47,47,47	1
2	IOD	B	507	1/1	0.97	0.05	33,33,33,33	0
2	IOD	B	521	1/1	0.97	0.10	50,50,50,50	1

*Continued on next page...*




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	IOD	B	510	1/1	0.97	0.06	37,37,37,37	0
2	IOD	B	504	1/1	0.98	0.04	41,41,41,41	1
2	IOD	B	523	1/1	0.98	0.05	37,37,37,37	1
2	IOD	B	513	1/1	0.98	0.05	36,36,36,36	1
2	IOD	A	510	1/1	0.98	0.07	32,32,32,32	1
2	IOD	A	512	1/1	0.98	0.08	28,28,28,28	1
2	IOD	A	514	1/1	0.98	0.11	35,35,35,35	1
2	IOD	B	505	1/1	0.98	0.05	47,47,47,47	1
2	IOD	A	513	1/1	0.98	0.10	29,29,29,29	1
4	EDO	A	550	4/4	0.98	0.10	12,13,14,16	0
2	IOD	A	501	1/1	0.99	0.05	31,31,31,31	0
2	IOD	A	505	1/1	0.99	0.08	24,24,24,24	1
2	IOD	A	509	1/1	0.99	0.06	32,32,32,32	1
5	MG	A	552	1/1	0.99	0.05	6,6,6,6	0
5	MG	B	535	1/1	0.99	0.06	14,14,14,14	0
2	IOD	A	542	1/1	0.99	0.06	23,23,23,23	0
2	IOD	B	514	1/1	0.99	0.04	31,31,31,31	0
2	IOD	A	506	1/1	0.99	0.08	30,30,30,30	1
2	IOD	B	503	1/1	0.99	0.02	36,36,36,36	1
2	IOD	A	507	1/1	0.99	0.07	34,34,34,34	1
2	IOD	B	501	1/1	0.99	0.04	31,31,31,31	0
2	IOD	A	508	1/1	0.99	0.07	25,25,25,25	0
2	IOD	A	533	1/1	0.99	0.14	36,36,36,36	1
2	IOD	A	502	1/1	1.00	0.03	19,19,19,19	0
2	IOD	B	511	1/1	1.00	0.05	30,30,30,30	1


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around G6Q B 533:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.