



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 19, 2023 – 08:32 PM EDT

PDB ID : 5HP1  
Title : STRUCTURE OF HIV-1 REVERSE TRANSCRIPTASE In COMPLEX  
WITH A DNA aptamer and FOSCARNET, a Pyrophosphate analog  
Authors : Das, K.; Arnold, E.  
Deposited on : 2016-01-19  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	555	
1	C	555	
2	B	444	
2	D	444	
3	E	38	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	38	 61% 29% 8%
4	G	2	 50% 50%
4	H	2	 50% 50%

## 2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 17386 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HIV-1 REVERSE TRANSCRIPTASE P66 SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	553	Total	C	N	O	S	0	0	0
			4494	2909	747	831	7			
1	C	553	Total	C	N	O	S	0	0	0
			4495	2910	748	830	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	280	SER	CYS	engineered mutation	UNP P03366
C	280	SER	CYS	engineered mutation	UNP P03366

- Molecule 2 is a protein called HIV-1 REVERSE TRANSCRIPTASE P51 SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	416	Total	C	N	O	S	0	0	0
			3430	2234	567	622	7			
2	D	410	Total	C	N	O	S	0	1	0
			3391	2212	559	614	6			

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-15	MET	-	initiating methionine	UNP P03366
B	-14	ALA	-	expression tag	UNP P03366
B	-13	HIS	-	expression tag	UNP P03366
B	-12	HIS	-	expression tag	UNP P03366
B	-11	HIS	-	expression tag	UNP P03366
B	-10	HIS	-	expression tag	UNP P03366
B	-9	HIS	-	expression tag	UNP P03366
B	-8	HIS	-	expression tag	UNP P03366
B	-7	ALA	-	expression tag	UNP P03366
B	-6	LEU	-	expression tag	UNP P03366

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-5	GLU	-	expression tag	UNP P03366
B	-4	VAL	-	expression tag	UNP P03366
B	-3	LEU	-	expression tag	UNP P03366
B	-2	PHE	-	expression tag	UNP P03366
B	-1	GLN	-	expression tag	UNP P03366
B	0	GLY	-	expression tag	UNP P03366
B	280	SER	CYS	engineered mutation	UNP P03366
D	-15	MET	-	initiating methionine	UNP P03366
D	-14	ALA	-	expression tag	UNP P03366
D	-13	HIS	-	expression tag	UNP P03366
D	-12	HIS	-	expression tag	UNP P03366
D	-11	HIS	-	expression tag	UNP P03366
D	-10	HIS	-	expression tag	UNP P03366
D	-9	HIS	-	expression tag	UNP P03366
D	-8	HIS	-	expression tag	UNP P03366
D	-7	ALA	-	expression tag	UNP P03366
D	-6	LEU	-	expression tag	UNP P03366
D	-5	GLU	-	expression tag	UNP P03366
D	-4	VAL	-	expression tag	UNP P03366
D	-3	LEU	-	expression tag	UNP P03366
D	-2	PHE	-	expression tag	UNP P03366
D	-1	GLN	-	expression tag	UNP P03366
D	0	GLY	-	expression tag	UNP P03366
D	280	SER	CYS	engineered mutation	UNP P03366

- Molecule 3 is a DNA chain called DNA (38-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	F	35	720	340	130	215	35	0	0	0
3	E	35	720	340	130	215	35	0	0	0

- Molecule 4 is an oligosaccharide called beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
4	G	2	23	12	11	0	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
4	H	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	4	Total	Mg	0	0
			4	4		
5	C	4	Total	Mg	0	0
			4	4		

- Molecule 6 is PHOSPHONOFORMIC ACID (three-letter code: PPF) (formula: CH<sub>3</sub>O<sub>5</sub>P).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	O	P	0	0
			7	1	5	1		
6	C	1	Total	C	O	P	0	0
			7	1	5	1		

- Molecule 7 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total O S 5 4 1	0	0
7	C	1	Total O S 5 4 1	0	0

- Molecule 8 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	B	1	Total C O 6 3 3	0	0
8	B	1	Total C O 6 3 3	0	0

- Molecule 9 is 3'-AZIDO-3'-DEOXYTHYMIDINE-5'-MONOPHOSPHATE (three-letter code: ATM) (formula: C<sub>10</sub>H<sub>14</sub>N<sub>5</sub>O<sub>7</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
9	F	1	22	10	5	6	1	0	0
9	E	1	22	10	5	6	1	0	0

- Molecule 10 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
10	A	1	1	1	0	0
10	C	1	1	1	0	0




### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: HIV-1 REVERSE TRANSCRIPTASE P66 SUBUNIT





#### • Molecule 1: HIV-1 REVERSE TRANSCRIPTASE P66 SUBUNIT







• Molecule 2: HIV-1 REVERSE TRANSCRIPTASE P51 SUBUNIT



• Molecule 2: HIV-1 REVERSE TRANSCRIPTASE P51 SUBUNIT



• Molecule 3: DNA (38-MER)



• Molecule 3: DNA (38-MER)





- Molecule 4: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose

Chain G: 50% 50%



- Molecule 4: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose

Chain H: 50% 50%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.33Å 129.07Å 131.54Å 90.00° 101.00° 90.00°	Depositor
Resolution (Å)	45.64 – 2.90 45.64 – 2.90	Depositor EDS
% Data completeness (in resolution range)	96.7 (45.64-2.90) 96.5 (45.64-2.90)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.87 (at 2.91Å)	Xtrriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, $R_{free}$	0.246 , 0.273 0.247 , 0.272	Depositor DCC
$R_{free}$ test set	2560 reflections (4.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	70.7	Xtrriage
Anisotropy	0.413	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 65.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	17386	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	104.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, MG, FRU, GLC, ATM, PPF, GOL, OMC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.28	0/4612	0.50	0/6268
1	C	0.27	0/4613	0.54	1/6268 (0.0%)
2	B	0.28	0/3528	0.49	0/4791
2	D	0.29	0/3493	0.55	2/4747 (0.0%)
3	E	0.58	0/759	1.03	3/1170 (0.3%)
3	F	0.58	0/759	1.03	3/1170 (0.3%)
All	All	0.31	0/17764	0.59	9/24414 (0.0%)

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	0	DA	O4'-C4'-C3'	-6.29	101.98	104.50
2	D	245	VAL	C-N-CA	-6.16	106.31	121.70
3	E	33	DG	O4'-C4'-C3'	-6.09	102.06	104.50
2	D	308	GLU	CA-CB-CG	5.96	126.50	113.40
3	F	31	DG	C4'-C3'-C2'	-5.54	98.11	103.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4494	0	4538	90	0
1	C	4495	0	4549	78	1
2	B	3430	0	3460	37	0
2	D	3391	0	3411	48	1
3	E	720	0	396	6	0
3	F	720	0	396	8	0
4	G	23	0	21	3	0
4	H	23	0	21	1	0
5	A	4	0	0	0	0
5	C	4	0	0	0	0
6	A	7	0	0	1	0
6	C	7	0	0	1	0
7	A	5	0	0	0	0
7	C	5	0	0	0	0
8	B	12	0	16	2	0
9	E	22	0	12	1	0
9	F	22	0	12	0	0
10	A	1	0	0	0	0
10	C	1	0	0	0	0
All	All	17386	0	16832	253	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 253 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:206:ARG:NH2	2:B:230:MET:O	2.02	0.94
2:B:70:LYS:HG3	2:B:226:PRO:HD2	1.53	0.90
1:C:134:SER:HG	1:C:139:THR:HG1	1.14	0.86
1:C:122:GLU:HA	1:C:125:ARG:HE	1.41	0.83
1:C:330:GLN:HE22	1:C:340:GLN:HE22	1.24	0.83

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:197:GLN:NE2	2:D:91:GLN:OE1[2_657]	2.06	0.14

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	551/555 (99%)	526 (96%)	24 (4%)	1 (0%)	47	78
1	C	551/555 (99%)	526 (96%)	23 (4%)	2 (0%)	34	66
2	B	412/444 (93%)	399 (97%)	13 (3%)	0	100	100
2	D	407/444 (92%)	396 (97%)	11 (3%)	0	100	100
All	All	1921/1998 (96%)	1847 (96%)	71 (4%)	3 (0%)	47	78

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	313	PRO
1	C	313	PRO
1	C	135	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	492/495 (99%)	469 (95%)	23 (5%)	26	59
1	C	493/495 (100%)	477 (97%)	16 (3%)	39	73
2	B	376/403 (93%)	373 (99%)	3 (1%)	81	94
2	D	371/403 (92%)	368 (99%)	3 (1%)	81	94
All	All	1732/1796 (96%)	1687 (97%)	45 (3%)	46	77

5 of 45 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	36	GLU
1	C	123	ASP
1	C	39	THR
1	C	73	LYS
1	C	136	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	428	GLN
1	C	509	GLN
1	C	545	ASN
1	C	151	GLN
1	A	545	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	OMC	F	4	3	19,22,23	2.70	6 (31%)	26,31,34	0.68	0
3	OMC	E	4	3	19,22,23	2.66	6 (31%)	26,31,34	0.70	0
3	OMC	E	2	3	19,22,23	2.70	6 (31%)	26,31,34	0.78	0
3	OMC	F	2	3	19,22,23	2.69	6 (31%)	26,31,34	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.



'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OMC	F	4	3	-	1/9/27/28	0/2/2/2
3	OMC	E	4	3	-	1/9/27/28	0/2/2/2
3	OMC	E	2	3	-	0/9/27/28	0/2/2/2
3	OMC	F	2	3	-	1/9/27/28	0/2/2/2

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	4	OMC	C6-C5	6.00	1.49	1.35
3	F	2	OMC	C6-C5	5.94	1.48	1.35
3	E	2	OMC	C6-C5	5.93	1.48	1.35
3	E	4	OMC	C6-C5	5.91	1.48	1.35
3	F	4	OMC	C2-N3	5.73	1.48	1.36

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	2	OMC	C1'-C2'-O2'-CM2
3	F	4	OMC	C1'-C2'-O2'-CM2
3	E	4	OMC	C1'-C2'-O2'-CM2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	4	OMC	1	0

## 5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GLC	G	1	4	11,11,12	0.54	0	15,15,17	0.86	1 (6%)
4	FRU	G	2	4	11,12,12	0.53	0	10,18,18	0.68	0
4	GLC	H	1	4	11,11,12	0.58	0	15,15,17	0.74	0
4	FRU	H	2	4	11,12,12	0.51	0	10,18,18	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GLC	G	1	4	-	2/2/19/22	0/1/1/1
4	FRU	G	2	4	-	3/5/24/24	0/1/1/1
4	GLC	H	1	4	-	0/2/19/22	0/1/1/1
4	FRU	H	2	4	-	2/5/24/24	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	1	GLC	C1-O5-C5	2.05	114.97	112.19

There are no chirality outliers.

5 of 7 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	H	2	FRU	O5-C5-C6-O6
4	H	2	FRU	C4-C5-C6-O6
4	G	2	FRU	C4-C5-C6-O6
4	G	2	FRU	O5-C5-C6-O6
4	G	2	FRU	O1-C1-C2-C3


There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	1	GLC	1	0
4	G	2	FRU	2	0
4	G	1	GLC	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
8	GOL	B	2003	-	5,5,5	0.40	0	5,5,5	0.23	0
9	ATM	F	101	3,5	19,23,24	1.29	2 (10%)	27,32,35	2.45	9 (33%)
8	GOL	B	2002	-	5,5,5	0.37	0	5,5,5	0.17	0
6	PPF	A	602	5	4,6,6	1.15	0	3,9,9	0.80	0
7	SO4	C	605	-	4,4,4	0.14	0	6,6,6	0.05	0
6	PPF	C	601	5	4,6,6	1.11	0	3,9,9	0.93	0
9	ATM	E	101	3,5	19,23,24	1.19	2 (10%)	27,32,35	2.36	9 (33%)
7	SO4	A	606	-	4,4,4	0.13	0	6,6,6	0.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	GOL	B	2003	-	-	4/4/4/4	-
9	ATM	F	101	3,5	-	0/10/24/25	0/2/2/2
8	GOL	B	2002	-	-	2/4/4/4	-
6	PPF	A	602	5	-	0/0/6/6	-
6	PPF	C	601	5	-	0/0/6/6	-
9	ATM	E	101	3,5	-	0/10/24/25	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	F	101	ATM	C6-C5	3.20	1.39	1.34
9	F	101	ATM	C4-C5	-3.10	1.39	1.44
9	E	101	ATM	C4-C5	-2.94	1.39	1.44
9	E	101	ATM	C6-C5	2.91	1.39	1.34

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	F	101	ATM	C4-N3-C2	-5.99	119.60	127.35
9	E	101	ATM	C4-N3-C2	-5.97	119.63	127.35
9	E	101	ATM	C5-C4-N3	5.79	120.25	115.31
9	F	101	ATM	C5-C4-N3	5.74	120.21	115.31
9	F	101	ATM	O4-C4-C5	-4.15	120.10	124.90

There are no chirality outliers.

5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
8	B	2002	GOL	O1-C1-C2-C3
8	B	2003	GOL	O1-C1-C2-C3
8	B	2003	GOL	C1-C2-C3-O3
8	B	2003	GOL	O1-C1-C2-O2
8	B	2002	GOL	O1-C1-C2-O2


There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	B	2003	GOL	2	0
6	A	602	PPF	1	0
6	C	601	PPF	1	0
9	E	101	ATM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	553/555 (99%)	0.87	89 (16%) <b>1</b> <b>1</b>	35, 103, 171, 223	0
1	C	553/555 (99%)	1.34	139 (25%) <b>0</b> <b>0</b>	35, 121, 215, 263	0
2	B	416/444 (93%)	0.48	33 (7%) <b>12</b> <b>10</b>	42, 82, 149, 195	0
2	D	410/444 (92%)	0.74	42 (10%) <b>6</b> <b>5</b>	37, 98, 167, 238	0
3	E	33/38 (86%)	0.37	0 <b>100</b> <b>100</b>	80, 118, 143, 210	0
3	F	33/38 (86%)	-0.07	0 <b>100</b> <b>100</b>	71, 103, 134, 175	0
All	All	1998/2074 (96%)	0.87	303 (15%) <b>2</b> <b>1</b>	35, 99, 191, 263	0

The worst 5 of 303 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	358	ARG	15.0
1	A	73	LYS	13.1
1	C	131	THR	11.2
1	C	26	LEU	11.1
1	A	71	TRP	9.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	OMC	E	2	21/22	0.87	0.26	104,117,125,132	0
3	OMC	E	4	21/22	0.89	0.29	86,101,112,115	0
3	OMC	F	4	21/22	0.94	0.20	64,73,78,83	0
3	OMC	F	2	21/22	0.95	0.18	88,97,105,106	0



### 6.3 Carbohydrates




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GLC	H	1	11/12	0.89	0.19	81,90,98,100	0
4	FRU	H	2	12/12	0.90	0.21	87,93,97,101	0
4	FRU	G	2	12/12	0.93	0.15	69,85,92,94	0
4	GLC	G	1	11/12	0.94	0.18	64,74,82,83	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain G:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	MG	A	604	1/1	0.49	0.18	130,130,130,130	0
5	MG	C	604	1/1	0.51	0.09	166,166,166,166	0

*Continued on next page...*




*Continued from previous page...*


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	MG	C	602	1/1	0.63	0.17	62,62,62,62	0
6	PPF	C	601	7/7	0.70	0.16	150,162,172,189	0
6	PPF	A	602	7/7	0.73	0.19	124,142,159,169	0
5	MG	A	601	1/1	0.76	0.13	80,80,80,80	0
9	ATM	E	101	22/23	0.82	0.30	121,134,156,168	0
8	GOL	B	2002	6/6	0.84	0.27	71,79,80,81	0
5	MG	C	606	1/1	0.85	0.13	63,63,63,63	0
5	MG	A	605	1/1	0.87	0.16	80,80,80,80	0
7	SO4	A	606	5/5	0.87	0.15	95,108,116,127	0
9	ATM	F	101	22/23	0.88	0.30	141,148,164,171	0
7	SO4	C	605	5/5	0.90	0.11	102,108,113,123	0
5	MG	A	603	1/1	0.90	0.14	133,133,133,133	1
5	MG	C	603	1/1	0.92	0.12	156,156,156,156	0
8	GOL	B	2003	6/6	0.92	0.26	60,64,68,73	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around ATM E 101:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.