



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 10:27 am BST

PDB ID : 3HUE
Title : Structure of the *S. pombe* Nbs1 FHA-BRCT1-BRCT2 domains
Authors : Williams, R.S.; Guenther, G.; Tainer, J.A.
Deposited on : 2009-06-13
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	341	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2497 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA repair and telomere maintenance protein nbs1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	307	2429	1544	400	470	7	8	0	0	0

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	331	ALA	-	EXPRESSION TAG	UNP O43070
A	332	ALA	-	EXPRESSION TAG	UNP O43070
A	333	ALA	-	EXPRESSION TAG	UNP O43070
A	334	LEU	-	EXPRESSION TAG	UNP O43070
A	335	GLU	-	EXPRESSION TAG	UNP O43070
A	336	HIS	-	EXPRESSION TAG	UNP O43070
A	337	HIS	-	EXPRESSION TAG	UNP O43070
A	338	HIS	-	EXPRESSION TAG	UNP O43070
A	339	HIS	-	EXPRESSION TAG	UNP O43070
A	340	HIS	-	EXPRESSION TAG	UNP O43070
A	341	HIS	-	EXPRESSION TAG	UNP O43070


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	68	Total	O	0	0
			68	68		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA repair and telomere maintenance protein nbs1



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	54.67Å 204.54Å 62.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80 42.65 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-2.80) 99.7 (42.65-2.70)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 2.69Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.236 , 0.285 0.234 , 0.284	Depositor DCC
R_{free} test set	502 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	39.6	Xtrriage
Anisotropy	0.574	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 17.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	2497	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/2464	0.83	0/3310

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	121	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2429	0	2434	24	0
2	A	68	0	0	2	0
All	All	2497	0	2434	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:236:THR:HG21	2:A:359:HOH:O	1.80	0.79
1:A:166:MSE:HE2	1:A:243:LEU:HD21	1.80	0.63
1:A:212:ASN:HD22	1:A:214:GLN:H	1.53	0.57
1:A:178:ASP:N	1:A:178:ASP:OD1	2.39	0.54
1:A:218:SER:HB2	1:A:220:GLU:OE1	2.09	0.53
1:A:166:MSE:HE1	1:A:294:ILE:HG22	1.92	0.51
1:A:244:LEU:HA	1:A:247:LEU:HD12	1.93	0.50
1:A:233:MSE:HB3	1:A:272:VAL:HB	1.93	0.49
1:A:244:LEU:HB2	1:A:251:ILE:HD11	1.93	0.49
1:A:166:MSE:HE3	1:A:298:TRP:CE3	2.50	0.47
1:A:260:ILE:HA	1:A:264:LEU:HD12	1.97	0.47
1:A:98:LEU:HB2	1:A:111:ALA:HB3	1.96	0.47
1:A:1:MSE:HE2	1:A:1:MSE:HB3	1.62	0.46
1:A:29:VAL:HG13	1:A:35:HIS:HB3	1.97	0.46
1:A:126:LEU:HD22	1:A:142:LEU:HD23	1.99	0.44
1:A:122:ASN:ND2	1:A:124:GLU:OE2	2.52	0.43
1:A:51:THR:HB	1:A:69:GLU:HB3	1.99	0.43
1:A:105:CYS:HA	1:A:106:PRO:HD3	1.89	0.43
1:A:81:VAL:HG22	1:A:100:ILE:HG12	2.00	0.43
1:A:8:GLY:HA2	2:A:380:HOH:O	2.20	0.41
1:A:151:PHE:HD2	1:A:175:ILE:HG23	1.86	0.41
1:A:70:VAL:HG21	1:A:81:VAL:HG21	2.01	0.41
1:A:152:VAL:HG21	1:A:181:LEU:HD11	2.01	0.41
1:A:55:PRO:HD3	1:A:137:GLY:HA2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	301/341 (88%)	283 (94%)	17 (6%)	1 (0%)	41 72

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	237	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	278/299 (93%)	246 (88%)	32 (12%)	5 17

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ILE
1	A	29	VAL
1	A	52	ILE
1	A	54	THR
1	A	74	ASP
1	A	89	ASN
1	A	91	ASP
1	A	114	ARG
1	A	116	MSE
1	A	117	CYS
1	A	119	GLN
1	A	122	ASN
1	A	140	THR
1	A	142	LEU
1	A	176	ILE
1	A	178	ASP
1	A	179	SER
1	A	184	LEU
1	A	186	THR
1	A	202	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	212	ASN
1	A	217	SER
1	A	231	CYS
1	A	234	LEU
1	A	237	SER
1	A	243	LEU
1	A	257	LEU
1	A	289	ILE
1	A	294	ILE
1	A	297	LEU
1	A	305	ASN
1	A	311	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	GLN
1	A	132	ASN
1	A	212	ASN
1	A	224	ASN
1	A	305	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	299/341 (87%)	-0.06	5 (1%) 70 63	22, 33, 42, 52	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	289	ILE	3.3
1	A	236	THR	3.2
1	A	88	GLN	2.5
1	A	314	ILE	2.4
1	A	321	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.