



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 20, 2023 – 10:30 AM EDT

PDB ID : 2HVR
Title : Structure of T4 RNA Ligase 2 with Nicked 5'-Adenylated nucleic acid duplex containing a 3'-deoxyribonucleotide at the nick
Authors : Nandakumar, J.; Lima, C.D.
Deposited on : 2006-07-30
Resolution : 2.45 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	24	 8% 25% 71%
1	F	24	 8% 25% 71%
2	D	12	 8% 17% 58% 25%
2	G	12	 25% 58% 17%
3	E	13	 23% 77%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	H	13	
4	A	335	
4	B	335	

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 7219 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*AP*TP*TP*CP*CP*GP*AP*TP*AP*GP*TP*GP*GP*GP*GP*TP*CP*GP*CP*AP*AP*TP*TP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	24	Total	C	N	O	P	0	0	0
			494	236	91	144	23			
1	F	24	Total	C	N	O	P	0	0	0
			494	236	91	144	23			

- Molecule 2 is DNA/RNA hybrid called 5'-D(*CP*AP*AP*TP*TP*GP*CP*GP*AP*C)-R(P*(OMC)P*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	12	Total	C	N	O	P	0	0	0
			241	116	44	70	11			
2	G	12	Total	C	N	O	P	0	0	0
			241	116	44	70	11			

- Molecule 3 is DNA/RNA hybrid called 5'-R(P*A)-D(P*CP*AP*CP*TP*AP*TP*CP*GP*GP*AP*AP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	13	Total	C	N	O	P	0	0	0
			268	127	50	78	13			
3	H	13	Total	C	N	O	P	0	0	0
			268	127	50	78	13			


- Molecule 4 is a protein called T4 RNA ligase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	A	319	Total	C	N	O	S	0	0	0
			2521	1617	409	482	13			
4	B	319	Total	C	N	O	S	0	0	0
			2521	1617	409	482	13			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	cloning artifact	UNP P32277
A	112	GLY	CYS	SEE REMARK 999	UNP P32277
B	0	SER	-	cloning artifact	UNP P32277
B	112	GLY	CYS	SEE REMARK 999	UNP P32277

- Molecule 5 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (three-letter code: BTB) (formula: C₈H₁₉NO₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	C	1	Total 14	C 8	N 1	O 5	0	0
5	F	1	Total 14	C 8	N 1	O 5	0	0
5	A	1	Total 14	C 8	N 1	O 5	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	7	Total 7	O 7	0	0
6	F	2	Total 2	O 2	0	0
6	G	2	Total 2	O 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	H	2	Total O 2 2	0	0
6	A	76	Total O 76 76	0	0
6	B	40	Total O 40 40	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 5'-D(*AP*TP*TP*CP*CP*GP*AP*TP*AP*GP*TP*GP*GP*GP*GP*TP*CP*GP*CP*AP*AP*TP*TP*G)-3'



- Molecule 1: 5'-D(*AP*TP*TP*CP*CP*GP*AP*TP*AP*GP*TP*GP*GP*GP*GP*TP*CP*GP*CP*AP*AP*TP*TP*G)-3'





- Molecule 2: 5'-D(*CP*AP*AP*TP*TP*GP*CP*GP*AP*C)-R(P*(OMC)P*C)-3'




- Molecule 2: 5'-D(*CP*AP*AP*TP*TP*GP*CP*GP*AP*C)-R(P*(OMC)P*C)-3'




- Molecule 3: 5'-R(P*A)-D(P*CP*AP*CP*TP*AP*TP*CP*GP*GP*AP*AP*T)-3'




- Molecule 3: 5'-R(P*A)-D(P*CP*AP*CP*TP*AP*TP*CP*GP*GP*AP*AP*T)-3'



- Molecule 4: T4 RNA ligase 2



- Molecule 4: T4 RNA ligase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	83.57Å 106.16Å 125.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.92 – 2.45 19.92 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.0 (19.92-2.45) 99.0 (19.92-2.45)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.82 (at 2.44Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.236 , 0.295 0.224 , 0.282	Depositor DCC
R_{free} test set	2056 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	48.9	Xtrriage
Anisotropy	0.296	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 54.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7219	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: O2C, BTB, OMC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.33	0/554	0.74	0/855
1	F	0.35	0/554	0.73	0/855
2	D	0.30	0/225	0.72	0/345
2	G	0.32	0/225	0.75	0/345
3	E	1.14	4/299 (1.3%)	1.76	8/455 (1.8%)
3	H	1.20	5/299 (1.7%)	1.75	8/455 (1.8%)
4	A	0.42	0/2570	0.64	1/3472 (0.0%)
4	B	0.40	0/2570	0.64	1/3471 (0.0%)
All	All	0.50	9/7296 (0.1%)	0.82	18/10253 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	F	0	2
2	D	0	2
2	G	0	2
All	All	0	7

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	236	A	N3-C4	10.31	1.41	1.34
3	E	236	A	N3-C4	9.10	1.40	1.34
3	H	236	A	C2-N3	7.27	1.40	1.33
3	E	236	A	C2-N3	6.97	1.39	1.33
3	H	236	A	N1-C2	6.86	1.40	1.34

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	E	236	A	C2-N3-C4	19.72	120.46	110.60
3	H	236	A	C2-N3-C4	19.56	120.38	110.60
3	E	236	A	N1-C2-N3	-16.95	120.83	129.30
3	H	236	A	N1-C2-N3	-16.68	120.96	129.30
3	E	236	A	N3-C4-C5	-11.19	118.97	126.80

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	14	DG	Sidechain
2	D	133	DA	Sidechain
2	D	134	DC	Sidechain
1	F	14	DG	Sidechain
1	F	15	DG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	494	0	273	25	0
1	F	494	0	273	27	0
2	D	241	0	138	12	0
2	G	241	0	138	10	0
3	E	268	0	148	12	0
3	H	268	0	148	16	0
4	A	2521	0	2511	92	0
4	B	2521	0	2517	111	0
5	A	14	0	19	0	0
5	C	14	0	19	4	0
5	F	14	0	19	4	0
6	A	76	0	0	1	0
6	B	40	0	0	1	0
6	C	7	0	0	0	0
6	F	2	0	0	0	0
6	G	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	H	2	0	0	1	0
All	All	7219	0	6203	296	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 296 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:9:DA:H2''	1:C:10:DG:H5''	1.28	1.14
2:G:133:DA:H2''	2:G:134:DC:H5''	1.21	1.13
4:A:251:ASP:HB2	4:A:308:ASP:HB3	1.33	1.10
2:D:133:DA:H2''	2:D:134:DC:H5''	1.16	1.08
3:H:241:DA:H2''	3:H:242:DT:H5'	1.42	1.01

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	A	315/335 (94%)	277 (88%)	29 (9%)	9 (3%)	4	2
4	B	315/335 (94%)	272 (86%)	31 (10%)	12 (4%)	3	1
All	All	630/670 (94%)	549 (87%)	60 (10%)	21 (3%)	4	1

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	A	309	ASN
4	B	250	ALA
4	B	275	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	A	13	ASN
4	A	112	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	A	277/293 (94%)	257 (93%)	20 (7%)	14	17
4	B	278/293 (95%)	261 (94%)	17 (6%)	18	24
All	All	555/586 (95%)	518 (93%)	37 (7%)	16	20

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	B	168	LEU
4	B	288	LEU
4	B	169	ASP
4	B	218	ASN
4	A	253	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
4	B	173	GLN
4	B	193	ASN
4	B	322	GLN
4	B	228	ASN
4	B	268	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	O2C	D	136	2	16,20,21	0.48	0	20,28,31	0.41	0
2	OMC	G	135	1,2	19,22,23	0.73	0	26,31,34	0.73	0
2	O2C	G	136	2	16,20,21	0.45	0	20,28,31	0.47	0
2	OMC	D	135	1,2	19,22,23	0.73	0	26,31,34	0.94	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	O2C	D	136	2	-	0/7/21/22	0/2/2/2
2	OMC	G	135	1,2	-	0/9/27/28	0/2/2/2
2	O2C	G	136	2	-	0/7/21/22	0/2/2/2
2	OMC	D	135	1,2	-	1/9/27/28	0/2/2/2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	135	OMC	CM2-O2'-C2'	-3.09	106.43	114.52

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	135	OMC	C3'-C2'-O2'-CM2

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	135	OMC	4	0
2	D	135	OMC	4	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	BTB	A	335	-	13,13,13	2.36	2 (15%)	7,16,16	1.82	3 (42%)
5	BTB	F	202	-	13,13,13	2.26	3 (23%)	7,16,16	1.69	2 (28%)
5	BTB	C	201	-	13,13,13	2.32	3 (23%)	7,16,16	1.72	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BTB	A	335	-	-	6/21/21/21	-
5	BTB	F	202	-	-	5/21/21/21	-
5	BTB	C	201	-	-	5/21/21/21	-

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	201	BTB	C5-N	5.69	1.56	1.48
5	A	335	BTB	C5-N	5.69	1.56	1.48
5	F	202	BTB	C5-N	5.64	1.56	1.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	335	BTB	C2-N	5.45	1.59	1.48
5	C	201	BTB	C2-N	5.16	1.58	1.48

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	201	BTB	C8-C7-N	2.87	122.81	111.59
5	F	202	BTB	C8-C7-N	2.84	122.69	111.59
5	A	335	BTB	C8-C7-N	2.75	122.31	111.59
5	A	335	BTB	O3-C3-C2	-2.42	104.83	111.44
5	A	335	BTB	C6-C5-N	2.26	120.41	111.59

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	201	BTB	C1-C2-N-C7
5	C	201	BTB	C3-C2-N-C7
5	C	201	BTB	C4-C2-N-C7
5	F	202	BTB	C1-C2-N-C7
5	F	202	BTB	C3-C2-N-C7

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	F	202	BTB	4	0
5	C	201	BTB	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	24/24 (100%)	0.55	2 (8%) 11 8	52, 91, 124, 130	0
1	F	24/24 (100%)	0.44	2 (8%) 11 8	48, 74, 150, 161	0
2	D	10/12 (83%)	0.73	1 (10%) 7 4	60, 101, 121, 123	0
2	G	10/12 (83%)	0.24	0 100 100	63, 74, 90, 93	0
3	E	13/13 (100%)	0.56	0 100 100	38, 91, 118, 126	0
3	H	13/13 (100%)	0.91	3 (23%) 0 0	38, 94, 141, 146	0
4	A	319/335 (95%)	-0.03	8 (2%) 57 53	18, 47, 103, 122	0
4	B	319/335 (95%)	0.14	15 (4%) 31 29	23, 52, 110, 127	0
All	All	732/768 (95%)	0.12	31 (4%) 36 33	18, 54, 114, 161	0

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	B	219	GLY	4.2
4	B	334	SER	3.5
4	B	290	VAL	3.3
4	B	277	ILE	3.3
4	B	21	TYR	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	OMC	G	135	21/22	0.96	0.13	24,40,48,55	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	OMC	D	135	21/22	0.97	0.16	32,41,46,49	0
2	O2C	D	136	19/20	0.97	0.12	23,36,46,48	0
2	O2C	G	136	19/20	0.98	0.10	34,38,46,47	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	BTB	F	202	14/14	0.80	0.31	60,81,84,90	0
5	BTB	C	201	14/14	0.83	0.23	74,81,92,96	0
5	BTB	A	335	14/14	0.86	0.30	110,114,117,119	0

6.5 Other polymers [i](#)

There are no such residues in this entry.