



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 15, 2024 – 11:28 am GMT

PDB ID : 6IAP  
Title : structure of human NKp46 in complex with antibody NKp46-1 and NKp46-4  
Authors : Roussel, A.; Amigues, B.  
Deposited on : 2018-11-27  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	182	 8% 80% 19%
2	D	211	 2% 79% 20%
3	E	217	 4% 71% 27%
4	H	220	 74% 25%
5	L	211	 8% 73% 25%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 8138 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Natural cytotoxicity triggering receptor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	182	1444	929	241	264	10	0	0	0

- Molecule 2 is a protein called Fab NKp46-4 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	211	1622	1017	274	326	5	0	0	0

- Molecule 3 is a protein called Fab NKp46-4 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	217	1625	1025	270	324	6	7	0	0

- Molecule 4 is a protein called Fab NKp46-1 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	H	220	1653	1042	275	329	7	0	0	0

- Molecule 5 is a protein called Fab NKp46-1 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	L	211	1635	1023	279	328	5	0	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	30	Total	O	0	0
			30	30		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
6	D	27	Total O 27 27	0	0
6	E	44	Total O 44 44	0	0
6	H	30	Total O 30 30	0	0
6	L	28	Total O 28 28	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Natural cytotoxicity triggering receptor 1



- Molecule 2: Fab NKp46-4 light chain




- Molecule 3: Fab NKp46-4 heavy chain





- Molecule 4: Fab NKp46-1 heavy chain





- Molecule 5: Fab NKp46-1 light chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.86Å 73.83Å 128.52Å 90.00° 90.57° 90.00°	Depositor
Resolution (Å)	48.47 – 2.90 48.47 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.3 (48.47-2.90) 99.3 (48.47-2.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.26 (at 2.91Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, $R_{free}$	0.187 , 0.263 0.196 , 0.191	Depositor DCC
$R_{free}$ test set	1439 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	77.4	Xtrriage
Anisotropy	0.327	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 86.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.026 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8138	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	91.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	0/1491	0.75	0/2031
2	D	0.50	0/1659	0.76	0/2255
3	E	0.56	0/1665	0.77	0/2269
4	H	0.50	0/1693	0.78	0/2305
5	L	0.47	0/1672	0.71	0/2270
All	All	0.51	0/8180	0.76	0/11130

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1444	0	1382	20	0
2	D	1622	0	1573	27	0
3	E	1625	0	1586	58	0
4	H	1653	0	1617	34	0
5	L	1635	0	1588	34	0
6	A	30	0	0	0	0
6	D	27	0	0	0	0
6	E	44	0	0	1	0
6	H	30	0	0	1	0
6	L	28	0	0	0	0
All	All	8138	0	7746	167	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 167 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:L:105:GLU:OE2	5:L:173:TYR:OH	1.60	1.19
3:E:51:ILE:HD11	3:E:70:ILE:HG22	1.25	1.10
3:E:51:ILE:HD11	3:E:70:ILE:C	1.74	1.07
3:E:91:THR:HG22	3:E:115:VAL:H	1.11	1.06
3:E:51:ILE:CD1	3:E:70:ILE:HG22	1.85	1.05

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	180/182 (99%)	160 (89%)	18 (10%)	2 (1%)	14	42
2	D	209/211 (99%)	199 (95%)	10 (5%)	0	100	100
3	E	215/217 (99%)	202 (94%)	12 (6%)	1 (0%)	29	61
4	H	218/220 (99%)	203 (93%)	14 (6%)	1 (0%)	29	61
5	L	209/211 (99%)	195 (93%)	14 (7%)	0	100	100
All	All	1031/1041 (99%)	959 (93%)	68 (7%)	4 (0%)	34	66

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	E	65	LYS
1	A	122	LEU
1	A	136	ARG
4	H	29	PHE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	158/161 (98%)	151 (96%)	7 (4%)	28	61
2	D	183/183 (100%)	173 (94%)	10 (6%)	21	53
3	E	184/185 (100%)	171 (93%)	13 (7%)	14	40
4	H	184/184 (100%)	175 (95%)	9 (5%)	25	57
5	L	187/187 (100%)	175 (94%)	12 (6%)	17	45
All	All	896/900 (100%)	845 (94%)	51 (6%)	20	51

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	E	169	THR
4	H	155	GLU
5	L	197	THR
3	E	187	THR
4	H	88	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	182/182 (100%)	0.58	15 (8%) 11 9	66, 94, 132, 154	0
2	D	211/211 (100%)	0.37	4 (1%) 66 65	62, 87, 111, 127	0
3	E	217/217 (100%)	0.37	8 (3%) 41 37	62, 88, 124, 140	1 (0%)
4	H	220/220 (100%)	0.20	1 (0%) 91 91	68, 89, 115, 125	0
5	L	211/211 (100%)	0.67	17 (8%) 12 9	66, 93, 119, 152	0
All	All	1041/1041 (100%)	0.43	45 (4%) 35 31	62, 90, 122, 154	1 (0%)

The worst 5 of 45 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	L	73	PHE	5.3
5	L	20	THR	5.1
1	A	160	HIS	4.3
3	E	193	LEU	4.3
1	A	119	TYR	4.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.