



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 19, 2023 – 08:49 PM EDT

PDB ID : 5IBK
Title : Skp1-F-box in complex with a ubiquitin variant
Authors : Orlicky, S.; Sicheri, F.
Deposited on : 2016-02-22
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	151	 4% 79% 13% • 7%
1	D	151	 2% 79% 13% • 7%
2	B	63	 11% 60% 13% 27%
2	E	63	 3% 65% 11% 24%
3	C	78	 % 79% 18% •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	78	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a red segment on the left labeled '14%', a green segment in the middle labeled '76%', and a yellow segment on the right labeled '21%'. A small grey dot is visible at the far right end of the bar.</p>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4284 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called S-phase kinase-associated protein 1, S-phase kinase-associated protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	141	1110	705	179	220	6	0	0	0
1	D	140	1120	712	181	220	7	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP P63208
A	0	ALA	-	expression tag	UNP P63208
A	?	-	ASP	deletion	UNP P63208
A	?	-	GLU	deletion	UNP P63208
A	?	-	GLY	deletion	UNP P63208
A	?	-	ASP	deletion	UNP P63208
A	?	-	ASP	deletion	UNP P63208
A	?	-	ASP	deletion	UNP P63208
A	70	GLY	-	linker	UNP P63208
A	79	GLY	-	linker	UNP P63208
A	80	SER	-	linker	UNP P63208
A	81	GLY	-	linker	UNP P63208
D	-1	GLY	-	expression tag	UNP P63208
D	0	ALA	-	expression tag	UNP P63208
D	?	-	ASP	deletion	UNP P63208
D	?	-	GLU	deletion	UNP P63208
D	?	-	GLY	deletion	UNP P63208
D	?	-	ASP	deletion	UNP P63208
D	?	-	ASP	deletion	UNP P63208
D	?	-	ASP	deletion	UNP P63208
D	78	GLY	-	linker	UNP P63208
D	79	GLY	-	linker	UNP P63208
D	80	SER	-	linker	UNP P63208
D	81	GLY	-	linker	UNP P63208

- Molecule 2 is a protein called F-box/WD repeat-containing protein 7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	46	392	258	66	67	1	0	0	0
2	E	48	412	272	69	70	1	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	261	GLY	-	expression tag	UNP Q969H0
B	262	ALA	-	expression tag	UNP Q969H0
E	261	GLY	-	expression tag	UNP Q969H0
E	262	ALA	-	expression tag	UNP Q969H0

- Molecule 3 is a protein called Polyubiquitin-B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	F	75	604	382	106	115	1	0	0	0
3	C	76	608	384	107	116	1	1	0	0

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	-3	GLY	-	expression tag	UNP P0CG47
F	-2	SER	-	expression tag	UNP P0CG47
F	0	SER	GLY	engineered mutation	UNP P0CG47
F	8	GLY	LEU	engineered mutation	UNP P0CG47
F	9	ALA	THR	engineered mutation	UNP P0CG47
F	10	ARG	GLY	engineered mutation	UNP P0CG47
F	11	THR	LYS	engineered mutation	UNP P0CG47
F	12	ALA	THR	engineered mutation	UNP P0CG47
F	42	ILE	ARG	engineered mutation	UNP P0CG47
F	46	SER	ALA	engineered mutation	UNP P0CG47
F	47	ARG	GLY	engineered mutation	UNP P0CG47
F	49	LEU	GLN	engineered mutation	UNP P0CG47
F	62	HIS	GLN	engineered mutation	UNP P0CG47
F	63	ARG	LYS	engineered mutation	UNP P0CG47
F	68	ARG	HIS	engineered mutation	UNP P0CG47
F	72	ILE	ARG	engineered mutation	UNP P0CG47
F	73	PHE	LEU	engineered mutation	UNP P0CG47

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	GLY	-	expression tag	UNP P0CG47
C	-2	SER	-	expression tag	UNP P0CG47
C	0	SER	GLY	engineered mutation	UNP P0CG47
C	8	GLY	LEU	engineered mutation	UNP P0CG47
C	9	ALA	THR	engineered mutation	UNP P0CG47
C	10	ARG	GLY	engineered mutation	UNP P0CG47
C	11	THR	LYS	engineered mutation	UNP P0CG47
C	12	ALA	THR	engineered mutation	UNP P0CG47
C	42	ILE	ARG	engineered mutation	UNP P0CG47
C	46	SER	ALA	engineered mutation	UNP P0CG47
C	47	ARG	GLY	engineered mutation	UNP P0CG47
C	49	LEU	GLN	engineered mutation	UNP P0CG47
C	62	HIS	GLN	engineered mutation	UNP P0CG47
C	63	ARG	LYS	engineered mutation	UNP P0CG47
C	68	ARG	HIS	engineered mutation	UNP P0CG47
C	72	ILE	ARG	engineered mutation	UNP P0CG47
C	73	PHE	LEU	engineered mutation	UNP P0CG47


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	24	Total O 24 24	0	0
4	B	1	Total O 1 1	0	0
4	F	2	Total O 2 2	0	0
4	C	5	Total O 5 5	0	0
4	D	6	Total O 6 6	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: S-phase kinase-associated protein 1,S-phase kinase-associated protein 1




- Molecule 1: S-phase kinase-associated protein 1,S-phase kinase-associated protein 1





- Molecule 2: F-box/WD repeat-containing protein 7




- Molecule 2: F-box/WD repeat-containing protein 7



- Molecule 3: Polyubiquitin-B



- Molecule 3: Polyubiquitin-B

Chain C:  %



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.37Å 98.02Å 107.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.71 – 2.50 47.71 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.2 (47.71-2.50) 99.3 (47.71-2.50)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 2.51Å)	Xtrriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.200 , 0.240 0.202 , 0.195	Depositor DCC
R_{free} test set	1180 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	64.9	Xtrriage
Anisotropy	0.358	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 52.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4284	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/1127	0.46	0/1524
1	D	0.26	0/1138	0.46	0/1539
2	B	0.26	0/401	0.54	0/544
2	E	0.27	0/422	0.48	0/572
3	C	0.23	0/615	0.46	0/828
3	F	0.25	0/611	0.50	0/823
All	All	0.26	0/4314	0.48	0/5830

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1110	0	1120	15	0
1	D	1120	0	1124	15	0
2	B	392	0	401	6	0
2	E	412	0	418	5	0
3	C	608	0	635	9	0
3	F	604	0	632	11	0
4	A	24	0	0	0	0
4	B	1	0	0	0	0
4	C	5	0	0	0	0
4	D	6	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	F	2	0	0	0	0
All	All	4284	0	4330	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 50 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:296:LEU:O	2:B:322:ARG:NH1	2.05	0.90
3:F:68:ARG:NH1	3:F:69:LEU:O	2.21	0.72
3:C:63:ARG:HG2	3:C:64:GLU:HG2	1.76	0.68
1:A:140:ASN:HD22	1:D:13:ILE:HG13	1.60	0.66
3:C:68:ARG:NH1	3:C:69:LEU:O	2.32	0.63

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	137/151 (91%)	132 (96%)	5 (4%)	0	100	100
1	D	136/151 (90%)	132 (97%)	4 (3%)	0	100	100
2	B	44/63 (70%)	43 (98%)	1 (2%)	0	100	100
2	E	46/63 (73%)	45 (98%)	1 (2%)	0	100	100
3	C	74/78 (95%)	72 (97%)	2 (3%)	0	100	100
3	F	73/78 (94%)	72 (99%)	1 (1%)	0	100	100
All	All	510/584 (87%)	496 (97%)	14 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/134 (94%)	121 (96%)	5 (4%)	31	56
1	D	128/134 (96%)	126 (98%)	2 (2%)	62	84
2	B	42/57 (74%)	41 (98%)	1 (2%)	49	74
2	E	44/57 (77%)	44 (100%)	0	100	100
3	C	69/70 (99%)	69 (100%)	0	100	100
3	F	69/70 (99%)	69 (100%)	0	100	100
All	All	478/522 (92%)	470 (98%)	8 (2%)	60	82

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	84	ASP
1	D	25	VAL
1	A	146	THR
1	A	142	LYS
2	B	286	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	140	ASN
1	A	152	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	141/151 (93%)	0.21	6 (4%) 35 38	30, 49, 140, 153	0
1	D	140/151 (92%)	0.12	3 (2%) 63 66	40, 80, 135, 175	0
2	B	46/63 (73%)	0.76	7 (15%) 2 1	43, 73, 108, 122	0
2	E	48/63 (76%)	0.34	2 (4%) 36 39	48, 81, 122, 132	0
3	C	76/78 (97%)	0.33	1 (1%) 77 79	44, 81, 120, 156	0
3	F	75/78 (96%)	0.82	11 (14%) 2 2	52, 94, 145, 184	0
All	All	526/584 (90%)	0.35	30 (5%) 23 25	30, 77, 138, 184	0

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	153	VAL	7.7
1	A	82	THR	4.9
1	D	157	ASN	4.6
1	A	-1	GLY	4.4
3	F	18	GLU	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.