



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 6, 2023 – 04:24 PM EDT

PDB ID : 1IHD
Title : Crystal Structure of Trigonal Form of D90E Mutant of Escherichia coli Asparaginase II
Authors : Borek, D.; Jaskolski, M.
Deposited on : 2001-04-19
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	326	74% (green), 18% (yellow), 6% (red), 2% (orange), 0% (grey)
1	C	326	75% (green), 17% (yellow), 6% (red), 2% (orange), 0% (grey)

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4674 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-asparaginase II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	307	2306	1443	394	461	8	0	1	0
1	C	308	2314	1447	396	463	8	0	1	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	90	GLU	ASP	engineered mutation	UNP P00805
C	90	GLU	ASP	engineered mutation	UNP P00805


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	26	Total	O	0	0
			26	26		
2	C	28	Total	O	0	0
			28	28		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: L-asparaginase II



- Molecule 1: L-asparaginase II



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	123.25Å 123.25Å 83.87Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 2.65 19.95 – 2.65	Depositor EDS
% Data completeness (in resolution range)	100.0 (10.00-2.65) 97.2 (19.95-2.65)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 2.67Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.166 , 0.209 0.213 , 0.243	Depositor DCC
R_{free} test set	1080 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	45.4	Xtrriage
Anisotropy	0.017	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 23.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.034 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4674	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.66	1/2347 (0.0%)	0.91	11/3196 (0.3%)
1	C	0.68	2/2355 (0.1%)	0.95	13/3207 (0.4%)
All	All	0.67	3/4702 (0.1%)	0.93	24/6403 (0.4%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	93	GLU	CD-OE1	-6.16	1.18	1.25
1	A	93	GLU	CD-OE1	-5.36	1.19	1.25
1	C	93	GLU	CD-OE2	-5.33	1.19	1.25

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	60	ASP	CB-CG-OD2	7.65	125.18	118.30
1	C	93	GLU	OE1-CD-OE2	-7.63	114.14	123.30
1	C	286	ASP	CB-CG-OD2	7.05	124.65	118.30
1	A	93	GLU	OE1-CD-OE2	-6.83	115.10	123.30
1	C	285	ASP	CB-CG-OD2	6.65	124.29	118.30
1	A	233	ASP	CB-CG-OD2	6.60	124.24	118.30
1	C	106	ASP	CB-CG-OD2	6.24	123.92	118.30
1	A	78	ASP	CB-CG-OD2	6.18	123.87	118.30
1	A	285	ASP	CB-CG-OD2	6.18	123.86	118.30
1	A	106	ASP	CB-CG-OD2	6.15	123.84	118.30
1	A	60	ASP	CB-CG-OD2	5.97	123.68	118.30
1	A	138	ASP	CB-CG-OD2	5.96	123.67	118.30
1	C	188	ASP	CB-CG-OD2	5.96	123.66	118.30
1	C	159	ASP	CB-CG-OD2	5.89	123.61	118.30
1	A	188	ASP	CB-CG-OD2	5.75	123.47	118.30
1	C	138	ASP	CB-CG-OD2	5.66	123.39	118.30
1	A	159	ASP	CB-CG-OD2	5.53	123.28	118.30
1	C	233	ASP	CB-CG-OD2	5.53	123.28	118.30
1	C	315	ASP	CB-CG-OD2	5.34	123.11	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	225	ASP	CB-CG-OD2	5.23	123.01	118.30
1	A	255	ASP	CB-CG-OD2	5.20	122.98	118.30
1	C	156	ASP	CB-CG-OD2	5.17	122.95	118.30
1	C	255	ASP	CB-CG-OD2	5.10	122.89	118.30
1	A	286	ASP	CB-CG-OD2	5.02	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2306	0	2303	32	0
1	C	2314	0	2309	36	0
2	A	26	0	0	0	0
2	C	28	0	0	0	0
All	All	4674	0	4612	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:163:THR:HB	1:A:170:THR:O	1.77	0.83
1:C:163:THR:HB	1:C:170:THR:O	1.79	0.83
1:C:267:VAL:HG21	1:C:291:PHE:CD1	2.18	0.78
1:C:175:ASN:HD21	1:C:276:GLY:H	1.33	0.76
1:C:164:ASN:HD22	1:C:167:ASP:H	1.33	0.75
1:A:267:VAL:CG2	1:A:291:PHE:CD1	2.69	0.75
1:C:267:VAL:CG2	1:C:291:PHE:CD1	2.70	0.75
1:A:267:VAL:HG21	1:A:291:PHE:CD1	2.23	0.73
1:C:175:ASN:ND2	1:C:276:GLY:H	1.89	0.70
1:A:300:GLN:HE21	1:C:272:ARG:CZ	2.05	0.69
1:C:73:ILE:HD11	1:C:85:ILE:HD11	1.74	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:117:PRO:HG2	1:C:120:SER:HB3	1.75	0.69
1:A:225:ASP:OD2	1:A:229:LYS:HE3	1.93	0.68
1:A:117:PRO:HG2	1:A:120:SER:HB3	1.75	0.68
1:A:73:ILE:HD11	1:A:85:ILE:HD11	1.77	0.67
1:A:267:VAL:HG21	1:A:291:PHE:CE1	2.33	0.63
1:C:267:VAL:HG21	1:C:291:PHE:CE1	2.33	0.63
1:A:106:ASP:OD1	1:A:106:ASP:N	2.23	0.61
1:A:53:VAL:HG23	1:A:54:VAL:HG13	1.83	0.61
1:A:198:THR:O	1:A:201:THR:HG23	2.01	0.60
1:C:53:VAL:HG23	1:C:54:VAL:HG13	1.85	0.59
1:C:106:ASP:OD1	1:C:106:ASP:N	2.27	0.58
1:A:76:ASP:O	1:A:79:LYS:HG3	2.03	0.58
1:C:267:VAL:HG23	1:C:267:VAL:O	2.03	0.58
1:A:300:GLN:NE2	1:C:272:ARG:HD2	2.20	0.57
1:C:225:ASP:OD2	1:C:229:LYS:HE3	2.06	0.56
1:C:198:THR:O	1:C:201:THR:HG23	2.06	0.55
1:C:268:VAL:HG23	1:C:306:LEU:HD22	1.90	0.54
1:C:298:ASN:HB2	1:C:299:PRO:CD	2.37	0.54
1:A:300:GLN:HE22	1:C:272:ARG:HD2	1.73	0.53
1:A:267:VAL:O	1:A:267:VAL:HG23	2.09	0.52
1:A:300:GLN:HE21	1:C:272:ARG:NE	2.11	0.48
1:A:298:ASN:HB2	1:A:299:PRO:CD	2.44	0.48
1:A:225:ASP:HB3	1:A:252:SER:HB3	1.95	0.48
1:C:112:VAL:HG22	1:C:113:GLY:N	2.30	0.47
1:C:261:ALA:HA	1:C:265:THR:O	2.15	0.47
1:C:69:LEU:HD23	1:C:69:LEU:HA	1.79	0.45
1:C:220:TYR:CD1	1:C:220:TYR:N	2.85	0.45
1:C:225:ASP:HB3	1:C:252:SER:HB3	1.99	0.45
1:C:125:GLY:N	1:C:126:PRO:CD	2.80	0.45
1:C:76:ASP:O	1:C:79:LYS:HG3	2.17	0.44
1:A:182:ILE:HG12	1:A:187:ILE:HG12	1.99	0.43
1:A:315:ASP:HA	1:A:316:PRO:HD3	1.90	0.43
1:A:226:LEU:HB2	1:A:227:PRO:HD3	2.00	0.43
1:A:107:LYS:H	1:A:107:LYS:HG2	1.51	0.43
1:C:225:ASP:CG	1:C:229:LYS:HE3	2.40	0.43
1:C:268:VAL:CG2	1:C:306:LEU:HD22	2.48	0.42
1:C:250:TYR:O	1:C:251:LYS:C	2.57	0.42
1:C:298:ASN:HB2	1:C:299:PRO:HD3	2.02	0.42
1:A:69:LEU:HD23	1:A:69:LEU:HA	1.69	0.42
1:A:261:ALA:HA	1:A:265:THR:O	2.19	0.42
1:A:2:PRO:HG2	1:A:137:ALA:HB1	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:248:ASN:OD1	1:A:249:LEU:N	2.50	0.41
1:C:315:ASP:HA	1:C:316:PRO:HD3	1.93	0.41
1:A:250:TYR:O	1:A:251:LYS:C	2.58	0.41
1:A:39:VAL:HG22	1:A:39:VAL:O	2.20	0.41
1:A:272:ARG:HD2	1:C:300:GLN:NE2	2.36	0.41
1:C:56:ILE:O	1:C:56:ILE:HG13	2.18	0.41
1:C:260:ALA:HB1	1:C:265:THR:HB	2.03	0.41
1:A:220:TYR:CE2	1:A:223:ALA:HA	2.56	0.40
1:A:158:ARG:CZ	1:A:297:LEU:HD21	2.52	0.40
1:A:272:ARG:NH1	1:C:300:GLN:HE21	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	304/326 (93%)	293 (96%)	10 (3%)	1 (0%)	41	56
1	C	305/326 (94%)	297 (97%)	7 (2%)	1 (0%)	41	56
All	All	609/652 (93%)	590 (97%)	17 (3%)	2 (0%)	41	56

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	251	LYS
1	C	251	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/266 (95%)	234 (92%)	19 (8%)	13	21
1	C	254/266 (96%)	235 (92%)	19 (8%)	13	21
All	All	507/532 (95%)	469 (92%)	38 (8%)	13	21

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	VAL
1	A	45	ILE
1	A	51	GLU
1	A	68	THR
1	A	79	LYS
1	A	86	THR
1	A	106	ASP
1	A	163	THR
1	A	179	LEU
1	A	201	THR
1	A	207	LYS
1	A	208	LEU
1	A	211	LEU
1	A	222	ASN
1	A	251	LYS
1	A	254	PHE
1	A	300	GLN
1	A	308	LEU
1	A	318	GLN
1	C	39	VAL
1	C	45	ILE
1	C	68	THR
1	C	79	LYS
1	C	86	THR
1	C	106	ASP
1	C	163	THR
1	C	179	LEU
1	C	186	LYS
1	C	201	THR
1	C	207	LYS
1	C	208	LEU
1	C	211	LEU
1	C	222	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	251	LYS
1	C	254	PHE
1	C	300	GLN
1	C	308	LEU
1	C	318	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	GLN
1	A	64	ASN
1	A	131	ASN
1	A	190	GLN
1	C	52	GLN
1	C	131	ASN
1	C	164	ASN
1	C	175	ASN
1	C	190	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.