

wwPDB X-ray Structure Validation Summary Report (i)

Mar 4, 2024 – 04:22 PM JST

PDB ID : 8IOX

> Title : Escherichia coli OpgD mutant-D388N

Authors Motouchi, S.; Nakajima, M.

2023-03-13 Deposited on

2.95 Å(reported) Resolution

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity 4.02b-467

> 1.8.5 (274361), CSD as541be (2020) Mogul

Xtriage (Phenix) 1.13

EDS 2.36

20191225.v01 (using entries in the PDB archive December 25th 2019) Percentile statistics

> Refmac 5.8.0158

CCP4 7.0.044 (Gargrove)

Ideal geometry (proteins) Engh & Huber (2001) Ideal geometry (DNA, RNA) Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.95 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	130704	3104 (3.00-2.92)
Clashscore	141614	3462 (3.00-2.92)
Ramachandran outliers	138981	3340 (3.00-2.92)
Sidechain outliers	138945	3343 (3.00-2.92)
RSRZ outliers	127900	2986 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	559	77%	15%	• 6%
1	В	559	79%	14%	• 6%
1	С	559	76%	16%	• 6%
1	D	559	77%	16%	6%
1	Е	559	78%	14%	• 6%
1	F	559	75%	18%	• 6%





Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	559	78% 14% • 6%
1	Н	559	77% 16% • 6%
1	I	559	76% 16% • 6%
1	J	559	79% 14% 6%
1	K	559	76% 16% · 6%
1	L	559	79% 14% • 6%
2	M	2	100%
2	N	2	100%
2	О	2	100%
2	Р	2	100%
2	Q	2	50%
2	R	2	100%
2	S	2	100%
2	Т	2	100%
2	U	2	100%
2	V	2	100%



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 51848 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Glucans biosynthesis protein D.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	524	Total	С	N	О	S	0	0	0
1	A	324	4246	2704	738	780	24	U	U	
1	В	524	Total	С	N	О	S	0	0	0
1	Б	324	4246	2704	738	780	24	U	0	U
1	С	524	Total	С	N	О	S	0	0	0
1		324	4246	2704	738	780	24	U	U	U
1	D	524	Total	С	N	Ο	S	0	0	0
1	D	324	4246	2704	738	780	24	U	U	
1	E	524	Total	С	N	Ο	S	0	0	0
1	ш	324	4246	2704	738	780	24	U	U	
1	F	524	Total	С	N	О	S	0	0	0
1	I.	324	4246	2704	738	780	24	O	U	0
1	G	524	Total	\mathbf{C}	N	O	S	0	0	0
1	d	024	4246	2704	738	780	24	O	O	U
1	Н	524	Total	\mathbf{C}	N	O	S	0	0	0
1	11	024	4246	2704	738	780	24	O	U	
1	I	524	Total	С	Ν	O	S	0	0	0
	1	024	4246	2704	738	780	24	0	0	U
1	J	524	Total	С	N	О	S	0	0	0
	0	024	4246	2704	738	780	24	0	0	
1	K	524	Total	\mathbf{C}	N	O	S	0	0	0
1	17	024	4246	2704	738	780	24	0	U	U
1	L	524	Total	\mathbf{C}	N	O	S	0	0	0
	L	024	4246	2704	738	780	24			U

There are 108 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	388	ASN	ASP	engineered mutation	UNP P40120
A	552	LEU	-	expression tag	UNP P40120
A	553	GLU	-	expression tag	UNP P40120
A	554	HIS	-	expression tag	UNP P40120
A	555	HIS	-	expression tag	UNP P40120



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
A	556	HIS	-	expression tag	UNP P40120
A	557	HIS	-	expression tag	UNP P40120
A	558	HIS	_	expression tag	UNP P40120
A	559	HIS	_	expression tag	UNP P40120
В	388	ASN	ASP	engineered mutation	UNP P40120
В	552	LEU	-	expression tag	UNP P40120
В	553	GLU	-	expression tag	UNP P40120
В	554	HIS	-	expression tag	UNP P40120
В	555	HIS	-	expression tag	UNP P40120
В	556	HIS	-	expression tag	UNP P40120
В	557	HIS	-	expression tag	UNP P40120
В	558	HIS	-	expression tag	UNP P40120
В	559	HIS	-	expression tag	UNP P40120
С	388	ASN	ASP	engineered mutation	UNP P40120
С	552	LEU	-	expression tag	UNP P40120
С	553	GLU	_	expression tag	UNP P40120
С	554	HIS	-	expression tag	UNP P40120
С	555	HIS	-	expression tag	UNP P40120
С	556	HIS	-	expression tag	UNP P40120
С	557	HIS	-	expression tag	UNP P40120
С	558	HIS	-	expression tag	UNP P40120
С	559	HIS	-	expression tag	UNP P40120
D	388	ASN	ASP	engineered mutation	UNP P40120
D	552	LEU	-	expression tag	UNP P40120
D	553	GLU	-	expression tag	UNP P40120
D	554	HIS	-	expression tag	UNP P40120
D	555	HIS	-	expression tag	UNP P40120
D	556	HIS	-	expression tag	UNP P40120
D	557	HIS	-	expression tag	UNP P40120
D	558	HIS	-	expression tag	UNP P40120
D	559	HIS	-	expression tag	UNP P40120
Е	388	ASN	ASP	engineered mutation	UNP P40120
Е	552	LEU	-	expression tag	UNP P40120
Е	553	GLU	-	expression tag	UNP P40120
Е	554	HIS	-	expression tag	UNP P40120
Е	555	HIS	-	expression tag	UNP P40120
Е	556	HIS	-	expression tag	UNP P40120
Е	557	HIS	-	expression tag	UNP P40120
Е	558	HIS	-	expression tag	UNP P40120
Е	559	HIS	-	expression tag	UNP P40120
F	388	ASN	ASP	engineered mutation	UNP P40120
F	552	LEU	-	expression tag	UNP P40120



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	553	GLU	-	expression tag	UNP P40120
F	554	HIS	-	expression tag	UNP P40120
F	555	HIS	-	expression tag	UNP P40120
F	556	HIS	-	expression tag	UNP P40120
F	557	HIS	-	expression tag	UNP P40120
F	558	HIS	-	expression tag	UNP P40120
F	559	HIS	-	expression tag	UNP P40120
G	388	ASN	ASP	engineered mutation	UNP P40120
G	552	LEU	-	expression tag	UNP P40120
G	553	GLU	-	expression tag	UNP P40120
G	554	HIS	-	expression tag	UNP P40120
G	555	HIS	-	expression tag	UNP P40120
G	556	HIS	-	expression tag	UNP P40120
G	557	HIS	-	expression tag	UNP P40120
G	558	HIS	-	expression tag	UNP P40120
G	559	HIS	-	expression tag	UNP P40120
Н	388	ASN	ASP	engineered mutation	UNP P40120
Н	552	LEU	-	expression tag	UNP P40120
Н	553	GLU	-	expression tag	UNP P40120
Н	554	HIS	-	expression tag	UNP P40120
Н	555	HIS	_	expression tag	UNP P40120
Н	556	HIS	-	expression tag	UNP P40120
Н	557	HIS	-	expression tag	UNP P40120
Н	558	HIS	-	expression tag	UNP P40120
Н	559	HIS	-	expression tag	UNP P40120
I	388	ASN	ASP	engineered mutation	UNP P40120
I	552	LEU	-	expression tag	UNP P40120
I	553	GLU	-	expression tag	UNP P40120
I	554	HIS	-	expression tag	UNP P40120
I	555	HIS	-	expression tag	UNP P40120
I	556	HIS	-	expression tag	UNP P40120
I	557	HIS	-	expression tag	UNP P40120
I	558	HIS	-	expression tag	UNP P40120
I	559	HIS	-	expression tag	UNP P40120
J	388	ASN	ASP	engineered mutation	UNP P40120
J	552	LEU	-	expression tag	UNP P40120
J	553	GLU	-	expression tag	UNP P40120
J	554	HIS	-	expression tag	UNP P40120
J	555	HIS	-	expression tag	UNP P40120
J	556	HIS	-	expression tag	UNP P40120
J	557	HIS	-	expression tag	UNP P40120
J	558	HIS	-	expression tag	UNP P40120



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
J	559	HIS	-	expression tag	UNP P40120
K	388	ASN	ASP	engineered mutation	UNP P40120
K	552	LEU	-	expression tag	UNP P40120
K	553	GLU	-	expression tag	UNP P40120
K	554	HIS	-	expression tag	UNP P40120
K	555	HIS	-	expression tag	UNP P40120
K	556	HIS	-	expression tag	UNP P40120
K	557	HIS	-	expression tag	UNP P40120
K	558	HIS	-	expression tag	UNP P40120
K	559	HIS	-	expression tag	UNP P40120
L	388	ASN	ASP	engineered mutation	UNP P40120
L	552	LEU	-	expression tag	UNP P40120
L	553	GLU	-	expression tag	UNP P40120
L	554	HIS	-	expression tag	UNP P40120
L	555	HIS	-	expression tag	UNP P40120
L	556	HIS	-	expression tag	UNP P40120
L	557	HIS	-	expression tag	UNP P40120
L	558	HIS		expression tag	UNP P40120
L	559	HIS	-	expression tag	UNP P40120

• Molecule 2 is an oligosaccharide called alpha-D-glucopyranose. (1-1)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	M	2	Total C O	0	0	0
			23 12 11			_
2	N	2	Total C O	0	0	0
	11	2	23 12 11		0	0
2	0	2	Total C O	0	0	0
2	O		23 12 11	0	U	U
2	Р	2	Total C O	0	0	0
2	Г	2	23 12 11		0	U
2	0	2	Total C O	0	0	0
2	Q	2	23 12 11	0	0	0
2	D	2	Total C O	0	0	0
	R	2	23 12 11	0	0	0
9	2 S	S 2	Total C O	0	0	0
			23 12 11	0	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	Т	2	Total C O 23 12 11	0	0	0
2	U	2	Total C O 23 12 11	0	0	0
2	V	2	Total C O 23 12 11	0	0	0

• Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf						
3	A	64	Total O	0	0						
	Λ	04	64 64	0	O						
3	В	43	Total O	0	0						
	Б	10	43 43	0	Ü						
3	С	47	Total O	0	0						
		11	47 47	Ü	Ü						
3	D	71	Total O	0	0						
		1.1	71 71		Ŭ.						
3	Е	74	Total O	0	0						
		, 1	74 74	0	Ů						
3	F	52	Total O	0	0						
				52 52							
3	G	78	Total O	0	0						
			78 78								
3	Н	Н	Н	Н	Н	Н	Н	42	Total O	0	0
			42 42								
3	I	37	Total O	0	0						
			37 37								
3	J	66	Total O	0	0						
			66 66								
3	K	56	Total O	0	0						
			56 56								
3	L	36	Total O	0	0						
			36 36								




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Glucans biosynthesis protein D








• Molecule 1: Glucans biosynthesis protein D




• Molecule 1: Glucans biosynthesis protein D




• Molecule 1: Glucans biosynthesis protein D






• Molecule 1: Glucans biosynthesis protein D











• Molecule 2:	alpha-D-glucopyranose (1-1)-alpha-D-glucopyranose
Chain R:	100%
GLC2 GLC2	
• Molecule 2:	${\it alpha-D-glucopyranose} - (1\text{-}1)\text{-}alpha-D-glucopyranose}$
Chain S:	100%
GLC2	
• Molecule 2:	alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose
Chain T:	100%
GLC2 GLC2	
• Molecule 2:	${\it alpha-D-glucopyranose} - (1\text{-}1) - {\it alpha-D-glucopyranose}$
Chain U:	100%
GLC2 GLC2	
• Molecule 2:	${\it alpha-D-glucopyranose} \\ -(1-1)-{\it alpha-D-glucopyranose}$
Chain V:	100%
מרכב פרכב	



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants	226.75Å 392.76Å 324.65Å	Donositon
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.14 - 2.95	Depositor
Resolution (A)	49.09 - 2.95	EDS
% Data completeness	99.9 (49.14-2.95)	Depositor
(in resolution range)	99.9 (49.09-2.95)	EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	3.62 (at 2.96Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.194 , 0.232	Depositor
10, 10 free	0.198 , 0.233	DCC
R_{free} test set	14722 reflections $(4.89%)$	wwPDB-VP
Wilson B-factor (\mathring{A}^2)	38.1	Xtriage
Anisotropy	0.008	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	$0.35 \; , \; 7.3$	EDS
L-test for twinning ²	$< L > = 0.51, < L^2> = 0.34$	Xtriage
Estimated twinning fraction	0.469 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l	Xtriage
<u> </u>	0.477 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Ŭ .
F_o, F_c correlation	0.93	EDS
Total number of atoms	51848	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 1.76% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.67	0/4375	0.87	0/5938
1	В	0.67	0/4375	0.86	0/5938
1	С	0.68	0/4375	0.87	0/5938
1	D	0.68	0/4375	0.89	0/5938
1	Е	0.68	0/4375	0.88	0/5938
1	F	0.67	0/4375	0.87	0/5938
1	G	0.67	0/4375	0.87	0/5938
1	Н	0.67	0/4375	0.88	0/5938
1	I	0.68	0/4375	0.88	0/5938
1	J	0.68	0/4375	0.89	0/5938
1	K	0.67	0/4375	0.87	0/5938
1	L	0.67	0/4375	0.88	0/5938
All	All	0.67	0/52500	0.88	0/71256

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4246	0	4037	52	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	В	4246	0	4037	47	0
1	С	4246	0	4037	63	0
1	D	4246	0	4037	58	0
1	Ε	4246	0	4037	55	0
1	F	4246	0	4037	60	0
1	G	4246	0	4037	46	0
1	Н	4246	0	4037	47	0
1	I	4246	0	4037	56	0
1	J	4246	0	4037	48	0
1	K	4246	0	4037	59	0
1	L	4246	0	4037	57	0
2	M	23	0	21	0	0
2	N	23	0	21	0	0
2	О	23	0	21	0	0
2	Р	23	0	21	0	0
2	Q	23	0	21	1	0
2	R	23	0	21	0	0
2	S	23	0	21	0	0
2	Т	23	0	21	0	0
2	U	23	0	21	0	0
2	V	23	0	21	0	0
3	A	64	0	0	2	0
3	В	43	0	0	0	0
3	С	47	0	0	1	0
3	D	71	0	0	1	0
3	Ε	74	0	0	3	0
3	F	52	0	0	1	0
3	G	78	0	0	3	0
3	Н	42	0	0	0	0
3	I	37	0	0	3	0
3	J	66	0	0	3	0
3	K	56	0	0	0	0
3	L	36	0	0	0	0
All	All	51848	0	48654	612	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 6.

The worst 5 of 612 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:H:491:ILE:HD11	1:H:498:ARG:HB2	1.44	1.00
1:B:425:ALA:HA	1:B:461:GLY:HA3	1.48	0.94
1:C:368:ASN:HD22	1:C:369:LYS:H	0.94	0.93
1:K:491:ILE:HD11	1:K:498:ARG:HB2	1.51	0.90
1:L:491:ILE:HD11	1:L:498:ARG:HB2	1.54	0.89

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percer	ntiles
1	A	522/559 (93%)	494 (95%)	26 (5%)	2 (0%)	34	69
1	В	522/559 (93%)	497 (95%)	25 (5%)	0	100	100
1	С	522/559 (93%)	495 (95%)	26 (5%)	1 (0%)	47	79
1	D	522/559 (93%)	496 (95%)	25 (5%)	1 (0%)	47	79
1	Е	522/559 (93%)	497 (95%)	24 (5%)	1 (0%)	47	79
1	F	522/559 (93%)	489 (94%)	31 (6%)	2 (0%)	34	69
1	G	522/559 (93%)	488 (94%)	32 (6%)	2 (0%)	34	69
1	Н	522/559 (93%)	493 (94%)	29 (6%)	0	100	100
1	I	522/559 (93%)	496 (95%)	24 (5%)	2 (0%)	34	69
1	J	522/559~(93%)	498 (95%)	24 (5%)	0	100	100
1	K	522/559 (93%)	490 (94%)	29 (6%)	3 (1%)	25	60
1	L	522/559 (93%)	490 (94%)	31 (6%)	1 (0%)	47	79
All	All	6264/6708 (93%)	5923 (95%)	326 (5%)	15 (0%)	47	79

5 of 15 Ramachandran outliers are listed below:

\mathbf{Mol}	Chain	Res	Type
1	Ε	463	ASP



Continued from previous page...

Mol	Chain	Res	Type
1	A	142	ASP
1	A	287	GLU
1	G	151	GLU
1	K	287	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	446/470 (95%)	429 (96%)	17 (4%)	33 66
1	В	446/470 (95%)	418 (94%)	28 (6%)	18 48
1	C	446/470 (95%)	424 (95%)	22 (5%)	25 58
1	D	446/470 (95%)	432 (97%)	14 (3%)	40 71
1	E	446/470 (95%)	429 (96%)	17 (4%)	33 66
1	F	446/470 (95%)	424 (95%)	22 (5%)	25 58
1	G	446/470 (95%)	424 (95%)	22 (5%)	25 58
1	Н	446/470 (95%)	421 (94%)	25 (6%)	21 53
1	I	446/470 (95%)	425 (95%)	21 (5%)	26 59
1	J	446/470 (95%)	429 (96%)	17 (4%)	33 66
1	K	446/470 (95%)	426 (96%)	20 (4%)	27 61
1	L	446/470 (95%)	429 (96%)	17 (4%)	33 66
All	All	5352/5640~(95%)	5110 (96%)	242 (4%)	27 61

5 of 242 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	463	ASP
1	K	431	ARG
1	Н	110	MET
1	K	386	THR
1	L	320	ASN



Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 120 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	262	HIS
1	K	554	HIS
1	Н	320	ASN
1	K	533	GLN
1	L	554	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

20 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	ain Res Link		Bond lengths			Bond angles		
MIOI	Type	Chain	Res	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	M	1	2	11,11,12	1.08	1 (9%)	15,15,17	2.02	3 (20%)
2	GLC	M	2	2	12,12,12	0.69	0	17,17,17	1.29	3 (17%)
2	GLC	N	1	2	11,11,12	0.37	0	15,15,17	0.60	0
2	GLC	N	2	2	12,12,12	0.42	0	17,17,17	1.00	0
2	GLC	О	1	2	11,11,12	1.07	0	15,15,17	1.56	4 (26%)
2	GLC	О	2	2	12,12,12	0.52	0	17,17,17	1.25	3 (17%)
2	GLC	Р	1	2	11,11,12	0.59	0	15,15,17	1.48	3 (20%)
2	GLC	Р	2	2	12,12,12	0.92	0	17,17,17	1.68	4 (23%)
2	GLC	Q	1	2	11,11,12	0.39	0	15,15,17	1.00	1 (6%)
2	GLC	Q	2	2	12,12,12	0.39	0	17,17,17	0.53	0



Mol	Trino	Chain	Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	R	1	2	11,11,12	1.14	1 (9%)	15,15,17	2.06	5 (33%)
2	GLC	R	2	2	12,12,12	0.67	0	17,17,17	1.34	3 (17%)
2	GLC	S	1	2	11,11,12	0.58	0	15,15,17	1.73	2 (13%)
2	GLC	S	2	2	12,12,12	1.03	0	17,17,17	1.49	6 (35%)
2	GLC	Т	1	2	11,11,12	1.43	2 (18%)	15,15,17	1.44	2 (13%)
2	GLC	Т	2	2	12,12,12	1.58	1 (8%)	17,17,17	1.22	1 (5%)
2	GLC	U	1	2	11,11,12	0.54	0	15,15,17	1.68	3 (20%)
2	GLC	U	2	2	12,12,12	1.00	1 (8%)	17,17,17	1.57	4 (23%)
2	GLC	V	1	2	11,11,12	1.19	1 (9%)	15,15,17	2.20	7 (46%)
2	GLC	V	2	2	12,12,12	0.70	0	17,17,17	1.27	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	M	1	2	-	0/2/19/22	0/1/1/1
2	GLC	M	2	2	-	2/2/22/22	0/1/1/1
2	GLC	N	1	2	-	0/2/19/22	0/1/1/1
2	GLC	N	2	2	-	0/2/22/22	0/1/1/1
2	GLC	О	1	2	-	0/2/19/22	0/1/1/1
2	GLC	О	2	2	-	0/2/22/22	0/1/1/1
2	GLC	Р	1	2	-	2/2/19/22	0/1/1/1
2	GLC	Р	2	2	-	0/2/22/22	0/1/1/1
2	GLC	Q	1	2	-	0/2/19/22	0/1/1/1
2	GLC	Q	2	2	-	0/2/22/22	0/1/1/1
2	GLC	R	1	2	-	0/2/19/22	0/1/1/1
2	GLC	R	2	2	-	2/2/22/22	0/1/1/1
2	GLC	S	1	2	-	0/2/19/22	0/1/1/1
2	GLC	S	2	2	-	2/2/22/22	0/1/1/1
2	GLC	Т	1	2	-	2/2/19/22	0/1/1/1
2	GLC	Т	2	2	-	0/2/22/22	0/1/1/1
2	GLC	U	1	2	-	0/2/19/22	0/1/1/1
2	GLC	U	2	2	-	0/2/22/22	0/1/1/1
2	GLC	V	1	2	-	0/2/19/22	0/1/1/1
2	GLC	V	2	2	_	0/2/22/22	0/1/1/1

The worst 5 of 7 bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
2	Т	2	GLC	C4-C5	2.84	1.59	1.53
2	Т	1	GLC	O4-C4	2.60	1.49	1.43
2	R	1	GLC	C4-C3	2.41	1.58	1.52
2	V	1	GLC	C4-C3	2.38	1.58	1.52
2	M	1	GLC	C4-C3	2.26	1.58	1.52

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	M	1	GLC	O3-C3-C4	4.89	121.64	110.35
2	U	1	GLC	O5-C5-C6	4.39	114.09	107.20
2	V	1	GLC	O3-C3-C4	4.30	120.30	110.35
2	R	1	GLC	O3-C3-C4	4.26	120.19	110.35
2	S	1	GLC	O5-C5-C6	3.82	113.19	107.20

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	Т	1	GLC	O5-C5-C6-O6
2	M	2	GLC	C4-C5-C6-O6
2	Τ	1	GLC	C4-C5-C6-O6
2	M	2	GLC	O5-C5-C6-O6
2	Р	1	GLC	C4-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	Q	1	GLC	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.








































5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2		$ ext{OWAB}(ext{Å}^2)$	Q<0.9
1	A	524/559~(93%)	0.33	5 (0%) 82 6	8	20, 31, 57, 86	0
1	В	524/559 (93%)	0.41	9 (1%) 70 5	3	22, 36, 69, 92	0
1	С	524/559 (93%)	0.34	6 (1%) 80 6	5	20, 33, 66, 94	0
1	D	524/559 (93%)	0.33	0 100 100		20, 32, 57, 82	0
1	Е	524/559 (93%)	0.31	3 (0%) 89 7	8	21, 32, 58, 92	0
1	F	524/559 (93%)	0.34	3 (0%) 89 7	8	20, 34, 66, 98	0
1	G	524/559 (93%)	0.29	6 (1%) 80 6	5	20, 31, 58, 89	0
1	Н	524/559 (93%)	0.39	5 (0%) 82 6	8	24, 36, 68, 94	0
1	I	524/559 (93%)	0.38	11 (2%) 63	46	20, 33, 65, 104	0
1	J	524/559 (93%)	0.35	3 (0%) 89 7	8	20, 32, 59, 84	0
1	K	524/559 (93%)	0.32	4 (0%) 86 7	3	20, 31, 59, 89	0
1	L	524/559 (93%)	0.39	5 (0%) 82 6	8	24, 37, 68, 96	0
All	All	6288/6708 (93%)	0.35	60 (0%) 82	38	20, 33, 64, 104	0

The worst 5 of 60 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	35	SER	4.7
1	В	556	HIS	4.4
1	A	556	HIS	4.2
1	A	557	HIS	4.1
1	L	556	HIS	3.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.




6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
2	GLC	Τ	2	12/12	0.85	0.50	37,42,45,47	0
2	GLC	Τ	1	11/12	0.86	0.49	36,43,48,57	0
2	GLC	N	2	12/12	0.89	0.38	20,20,20,20	0
2	GLC	Q	1	11/12	0.89	0.36	20,20,20,20	0
2	GLC	Р	1	11/12	0.90	0.32	51,61,74,77	0
2	GLC	M	1	11/12	0.90	0.26	44,53,58,58	0
2	GLC	M	2	12/12	0.91	0.30	45,56,62,62	0
2	GLC	О	2	12/12	0.92	0.27	53,70,77,79	0
2	GLC	N	1	11/12	0.93	0.39	20,20,20,20	0
2	GLC	R	1	11/12	0.93	0.28	44,51,53,54	0
2	GLC	S	1	11/12	0.93	0.32	63,74,77,78	0
2	GLC	О	1	11/12	0.93	0.33	53,67,73,75	0
2	GLC	Р	2	12/12	0.93	0.29	58,67,73,74	0
2	GLC	R	2	12/12	0.94	0.27	46,54,57,62	0
2	GLC	S	2	12/12	0.94	0.24	58,68,72,73	0
2	GLC	U	1	11/12	0.94	0.31	50,64,71,73	0
2	GLC	U	2	12/12	0.94	0.29	52,64,68,70	0
2	GLC	V	1	11/12	0.94	0.22	42,51,55,56	0
2	GLC	V	2	12/12	0.94	0.25	47,57,61,64	0
2	GLC	Q	2	12/12	0.95	0.40	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







































Electron density around Chain T:


 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)
















6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.

