



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 5, 2023 – 04:09 PM JST

PDB ID : 8ISN  
Title : HLA-A24 in complex with modified 9mer WT1 peptide  
Authors : Bekker, G.J.; Numoto, N.; Kawasaki, M.; Hayashi, T.; Yabuno, S.; Kozono, Y.; Shimizu, T.; Kozono, H.; Ito, N.; Oda, M.; Kamiya, N.  
Deposited on : 2023-03-21  
Resolution : 2.48 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5857 (2.50-2.46)
Clashscore	141614	6594 (2.50-2.46)
Ramachandran outliers	138981	6469 (2.50-2.46)
Sidechain outliers	138945	6471 (2.50-2.46)
RSRZ outliers	127900	5738 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	305	 % 83% 7% 10%
1	D	305	 6% 82% 8% 10%
2	B	99	 92% 8%
2	E	99	 13% 92% 8%
3	C	9	 78% 11% 11%
3	F	9	 100%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	G	4	

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6495 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class I antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	274	2221	1382	403	426	10	0	0	0
1	D	274	2221	1382	403	426	10	0	0	0

There are 58 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	277	GLY	-	expression tag	UNP D9UAY1
A	278	ALA	-	expression tag	UNP D9UAY1
A	279	SER	-	expression tag	UNP D9UAY1
A	280	GLY	-	expression tag	UNP D9UAY1
A	281	ASP	-	expression tag	UNP D9UAY1
A	282	TYR	-	expression tag	UNP D9UAY1
A	283	LYS	-	expression tag	UNP D9UAY1
A	284	ASP	-	expression tag	UNP D9UAY1
A	285	ASP	-	expression tag	UNP D9UAY1
A	286	ASP	-	expression tag	UNP D9UAY1
A	287	ASP	-	expression tag	UNP D9UAY1
A	288	LYS	-	expression tag	UNP D9UAY1
A	289	GLY	-	expression tag	UNP D9UAY1
A	290	GLY	-	expression tag	UNP D9UAY1
A	291	GLY	-	expression tag	UNP D9UAY1
A	292	GLY	-	expression tag	UNP D9UAY1
A	293	LEU	-	expression tag	UNP D9UAY1
A	294	ASN	-	expression tag	UNP D9UAY1
A	295	ASP	-	expression tag	UNP D9UAY1
A	296	ILE	-	expression tag	UNP D9UAY1
A	297	PHE	-	expression tag	UNP D9UAY1
A	298	GLU	-	expression tag	UNP D9UAY1
A	299	ALA	-	expression tag	UNP D9UAY1
A	300	GLN	-	expression tag	UNP D9UAY1
A	301	LYS	-	expression tag	UNP D9UAY1

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	302	ILE	-	expression tag	UNP D9UAY1
A	303	GLU	-	expression tag	UNP D9UAY1
A	304	TRP	-	expression tag	UNP D9UAY1
A	305	HIS	-	expression tag	UNP D9UAY1
D	277	GLY	-	expression tag	UNP D9UAY1
D	278	ALA	-	expression tag	UNP D9UAY1
D	279	SER	-	expression tag	UNP D9UAY1
D	280	GLY	-	expression tag	UNP D9UAY1
D	281	ASP	-	expression tag	UNP D9UAY1
D	282	TYR	-	expression tag	UNP D9UAY1
D	283	LYS	-	expression tag	UNP D9UAY1
D	284	ASP	-	expression tag	UNP D9UAY1
D	285	ASP	-	expression tag	UNP D9UAY1
D	286	ASP	-	expression tag	UNP D9UAY1
D	287	ASP	-	expression tag	UNP D9UAY1
D	288	LYS	-	expression tag	UNP D9UAY1
D	289	GLY	-	expression tag	UNP D9UAY1
D	290	GLY	-	expression tag	UNP D9UAY1
D	291	GLY	-	expression tag	UNP D9UAY1
D	292	GLY	-	expression tag	UNP D9UAY1
D	293	LEU	-	expression tag	UNP D9UAY1
D	294	ASN	-	expression tag	UNP D9UAY1
D	295	ASP	-	expression tag	UNP D9UAY1
D	296	ILE	-	expression tag	UNP D9UAY1
D	297	PHE	-	expression tag	UNP D9UAY1
D	298	GLU	-	expression tag	UNP D9UAY1
D	299	ALA	-	expression tag	UNP D9UAY1
D	300	GLN	-	expression tag	UNP D9UAY1
D	301	LYS	-	expression tag	UNP D9UAY1
D	302	ILE	-	expression tag	UNP D9UAY1
D	303	GLU	-	expression tag	UNP D9UAY1
D	304	TRP	-	expression tag	UNP D9UAY1
D	305	HIS	-	expression tag	UNP D9UAY1

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	0	0
			829	528	140	158	3			
2	E	99	Total	C	N	O	S	0	0	0
			829	528	140	158	3			

- Molecule 3 is a protein called CYS-TYR-THR-TRP-ASN-GLN-MET-ASN-LEU.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	9	Total	C	N	O	S	0	0	0
			81	51	13	15	2			
3	F	9	Total	C	N	O	S	0	0	0
			81	51	13	15	2			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranos e-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	G	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	79	Total	O	0	0
			79	79		
6	B	36	Total	O	0	0
			36	36		
6	C	1	Total	O	0	0
			1	1		
6	D	32	Total	O	0	0
			32	32		
6	E	5	Total	O	0	0
			5	5		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MHC class I antigen




- Molecule 1: MHC class I antigen




- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin



- Molecule 3: CYS-TYR-THR-TRP-ASN-GLN-MET-ASN-LEU







- Molecule 3: CYS-TYR-THR-TRP-ASN-GLN-MET-ASN-LEU

Chain F:  100%

There are no outlier residues recorded for this chain.

- Molecule 4: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  50% 50%



## 4 Data and refinement statistics i

Property	Value	Source
Space group	F 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	161.40Å 166.13Å 180.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.24 – 2.48 49.24 – 2.48	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.24-2.48) 91.3 (49.24-2.48)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.25	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.74 (at 2.48Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.222 , 0.258 0.223 , 0.257	Depositor DCC
$R_{free}$ test set	2014 reflections (4.69%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.1	Xtrriage
Anisotropy	0.397	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 36.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.018 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6495	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	62.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.68% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAG, BMA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.24	0/2281	0.51	0/3092
1	D	0.24	0/2281	0.51	0/3092
2	B	0.24	0/852	0.47	0/1152
2	E	0.24	0/852	0.48	0/1152
3	C	0.24	0/83	0.38	0/111
3	F	0.26	0/83	0.36	0/111
All	All	0.24	0/6432	0.50	0/8710

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2221	0	2081	12	0
1	D	2221	0	2082	12	0
2	B	829	0	794	5	0
2	E	829	0	794	3	0
3	C	81	0	73	1	0
3	F	81	0	73	0	0
4	G	50	0	43	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	18	0	24	0	0
5	B	12	0	16	2	0
6	A	79	0	0	1	0
6	B	36	0	0	1	0
6	C	1	0	0	0	0
6	D	32	0	0	2	0
6	E	5	0	0	0	0
All	All	6495	0	5980	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 31 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:MET:HE3	1:A:171:TYR:HE2	1.53	0.73
2:B:44:GLU:HA	5:B:102:GOL:H31	1.83	0.60
1:D:83:ARG:NH1	6:D:402:HOH:O	2.37	0.57
2:B:7:ILE:HD12	2:B:91:LYS:HD3	1.89	0.55
1:A:7:TYR:OH	3:C:1:CYS:N	2.37	0.53

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/305 (89%)	264 (97%)	7 (3%)	1 (0%)	34	52
1	D	272/305 (89%)	262 (96%)	10 (4%)	0	100	100
2	B	97/99 (98%)	94 (97%)	3 (3%)	0	100	100
2	E	97/99 (98%)	93 (96%)	4 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	C	7/9 (78%)	5 (71%)	2 (29%)	0	100	100
3	F	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
All	All	752/826 (91%)	724 (96%)	27 (4%)	1 (0%)	51	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	225	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/253 (91%)	226 (98%)	4 (2%)	60	81
1	D	230/253 (91%)	228 (99%)	2 (1%)	78	91
2	B	94/94 (100%)	93 (99%)	1 (1%)	73	88
2	E	94/94 (100%)	91 (97%)	3 (3%)	39	63
3	C	9/9 (100%)	7 (78%)	2 (22%)	1	1
3	F	9/9 (100%)	9 (100%)	0	100	100
All	All	666/712 (94%)	654 (98%)	12 (2%)	59	80

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	105	SER
1	D	108	ARG
2	E	70	PHE
2	E	28	SER
1	A	256	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	114	HIS
1	D	156	GLN
3	F	5	ASN
3	C	8	ASN
1	A	180	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	NAG	G	1	1,4	14,14,15	0.25	0	17,19,21	0.47	0
4	NAG	G	2	4	14,14,15	0.23	0	17,19,21	0.64	0
4	BMA	G	3	4	11,11,12	1.40	2 (18%)	15,15,17	1.65	5 (33%)
4	MAN	G	4	4	11,11,12	0.98	1 (9%)	15,15,17	1.47	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	G	2	4	-	2/6/23/26	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BMA	G	3	4	-	2/2/19/22	0/1/1/1
4	MAN	G	4	4	-	2/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	G	3	BMA	C2-C3	2.96	1.56	1.52
4	G	4	MAN	C1-C2	2.91	1.58	1.52
4	G	3	BMA	C1-C2	2.63	1.58	1.52

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	3	BMA	C1-C2-C3	4.22	114.86	109.67
4	G	4	MAN	C1-O5-C5	3.65	117.14	112.19
4	G	3	BMA	O5-C1-C2	2.51	114.65	110.77
4	G	4	MAN	C1-C2-C3	2.23	112.40	109.67
4	G	4	MAN	O2-C2-C3	-2.22	105.69	110.14

There are no chirality outliers.


5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	G	4	MAN	O5-C5-C6-O6
4	G	4	MAN	C4-C5-C6-O6
4	G	1	NAG	O5-C5-C6-O6
4	G	1	NAG	C4-C5-C6-O6
4	G	3	BMA	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	GOL	A	401	-	5,5,5	0.94	0	5,5,5	0.96	0
5	GOL	B	101	-	5,5,5	0.92	0	5,5,5	0.95	0
5	GOL	A	403	-	5,5,5	0.98	0	5,5,5	0.91	0
5	GOL	B	102	-	5,5,5	0.95	0	5,5,5	0.95	0
5	GOL	A	402	-	5,5,5	0.90	0	5,5,5	0.99	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	401	-	-	0/4/4/4	-
5	GOL	B	101	-	-	0/4/4/4	-
5	GOL	A	403	-	-	2/4/4/4	-
5	GOL	B	102	-	-	0/4/4/4	-
5	GOL	A	402	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	403	GOL	O1-C1-C2-C3
5	A	403	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	102	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	274/305 (89%)	0.16	3 (1%) 80 82	32, 49, 73, 99	0
1	D	274/305 (89%)	0.57	17 (6%) 20 21	40, 68, 101, 120	0
2	B	99/99 (100%)	-0.17	0 100 100	32, 44, 69, 93	0
2	E	99/99 (100%)	0.91	13 (13%) 3 2	56, 88, 106, 109	0
3	C	9/9 (100%)	0.50	0 100 100	43, 46, 51, 56	0
3	F	9/9 (100%)	0.52	0 100 100	45, 50, 54, 55	0
All	All	764/826 (92%)	0.37	33 (4%) 35 37	32, 57, 98, 120	0

The worst 5 of 33 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	19	LYS	4.7
2	E	87	LEU	4.2
1	D	227	ASP	3.6
2	E	1	ILE	3.1
1	D	91	GLY	3.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	BMA	G	3	11/12	0.74	0.23	86,102,112,114	0
4	MAN	G	4	11/12	0.78	0.33	114,121,126,128	0
4	NAG	G	1	14/15	0.80	0.20	91,100,105,110	0
4	NAG	G	2	14/15	0.84	0.32	109,112,114,115	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	GOL	B	102	6/6	0.81	0.21	60,62,63,64	0
5	GOL	A	402	6/6	0.88	0.17	46,49,57,58	0
5	GOL	B	101	6/6	0.90	0.28	53,53,58,64	0
5	GOL	A	403	6/6	0.90	0.17	39,45,46,51	0
5	GOL	A	401	6/6	0.95	0.30	53,55,58,63	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.