



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 5, 2024 – 05:08 PM EDT

PDB ID : 4IX9
Title : Crystal structure of subunit F of V-ATPase from *S. cerevisiae*
Authors : Basak, S.; Balakrishna, A.M.; Manimekalai, M.S.S.; Gruber, G.
Deposited on : 2013-01-24
Resolution : 2.33 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2747 (2.36-2.32)
Clashscore	180529	2936 (2.36-2.32)
Ramachandran outliers	177936	2912 (2.36-2.32)
Sidechain outliers	177891	2912 (2.36-2.32)
RSRZ outliers	164620	2747 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	94	 70% 24% 6%
1	B	94	 64% 29% 6%
1	C	94	 70% 22% 5%
1	D	94	 57% 35% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	B	103	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3358 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called V-type proton ATPase subunit F.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	92	Total 757	C 476	N 127	O 153	Se 1	0	2	0
1	B	93	Total 777	C 487	N 129	O 160	Se 1	0	4	0
1	C	92	Total 772	C 487	N 127	O 157	Se 1	0	4	0
1	D	90	Total 743	C 467	N 123	O 152	Se 1	0	3	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	69	MSE	ILE	engineered mutation	UNP P39111
B	69	MSE	ILE	engineered mutation	UNP P39111
C	69	MSE	ILE	engineered mutation	UNP P39111
D	69	MSE	ILE	engineered mutation	UNP P39111

- Molecule 2 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	8	4	1	3	0	0
2	A	1	8	4	1	3	0	0
2	B	1	8	4	1	3	0	0
2	D	1	8	4	1	3	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	39	Total	O	0	2
			41	41		
4	B	82	Total	O	0	0
			82	82		
4	C	69	Total	O	0	0
			69	69		
4	D	55	Total	O	0	0
			55	55		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: V-type proton ATPase subunit F

Chain A: 



- Molecule 1: V-type proton ATPase subunit F

Chain B: 




- Molecule 1: V-type proton ATPase subunit F

Chain C: 



- Molecule 1: V-type proton ATPase subunit F

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	47.15Å 160.31Å 102.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.53 – 2.33 26.53 – 2.33	Depositor EDS
% Data completeness (in resolution range)	99.4 (26.53-2.33) 99.5 (26.53-2.33)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.19 (at 2.33Å)	Xtrriage
Refinement program	REFMAC 5.7.0025	Depositor
R, R_{free}	0.152 , 0.212 0.155 , 0.213	Depositor DCC
R_{free} test set	865 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	34.4	Xtrriage
Anisotropy	0.062	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3358	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.76	0/766	0.89	0/1032
1	B	0.80	0/786	0.94	3/1060 (0.3%)
1	C	0.80	0/782	0.93	2/1055 (0.2%)
1	D	0.68	0/752	0.91	1/1017 (0.1%)
All	All	0.76	0/3086	0.92	6/4164 (0.1%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	78	ARG	NE-CZ-NH1	8.62	124.61	120.30
1	C	69	MSE	N-CA-CB	-7.25	97.55	110.60
1	D	80	ARG	NE-CZ-NH1	7.03	123.82	120.30
1	B	78	ARG	NE-CZ-NH2	-6.58	117.01	120.30
1	C	69	MSE	CG-SE-CE	-6.55	84.50	98.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	757	0	744	23	0
1	B	777	0	755	31	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	772	0	752	25	0
1	D	743	0	721	36	0
2	A	16	0	24	4	0
2	B	8	0	12	4	0
2	D	8	0	12	3	0
3	B	24	0	32	7	0
3	C	6	0	8	2	0
4	A	41	0	0	0	0
4	B	82	0	0	1	0
4	C	69	0	0	3	0
4	D	55	0	0	5	0
All	All	3358	0	3060	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 108 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:76:ASN:OD1	2:A:102:TRS:H31	1.52	1.07
1:B:42:GLU:O	3:B:101:GOL:H31	1.57	1.04
1:A:80[B]:ARG:HG3	1:A:80[B]:ARG:NH1	1.65	1.02
1:A:80[B]:ARG:HH11	1:A:80[B]:ARG:CG	1.73	1.01
1:A:80[B]:ARG:HG3	1:A:80[B]:ARG:HH11	0.84	1.00

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	92/94 (98%)	89 (97%)	3 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	95/94 (101%)	92 (97%)	3 (3%)	0	100	100
1	C	94/94 (100%)	91 (97%)	3 (3%)	0	100	100
1	D	91/94 (97%)	86 (94%)	5 (6%)	0	100	100
All	All	372/376 (99%)	358 (96%)	14 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	82/79 (104%)	74 (90%)	8 (10%)	6	5
1	B	84/79 (106%)	74 (88%)	10 (12%)	4	3
1	C	84/79 (106%)	77 (92%)	7 (8%)	9	9
1	D	81/79 (102%)	75 (93%)	6 (7%)	11	11
All	All	331/316 (105%)	300 (91%)	31 (9%)	7	6

5 of 31 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	59[B]	GLU
1	D	28	ILE
1	B	92	LEU
1	D	53	LYS
1	C	92	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	71	GLN
1	C	86	ASN
1	D	72	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	36	ASN
1	B	70	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TRS	A	102	-	7,7,7	1.01	0	9,9,9	1.81	2 (22%)
3	GOL	B	105	-	5,5,5	0.31	0	5,5,5	0.47	0
2	TRS	A	101	-	7,7,7	0.59	0	9,9,9	1.18	1 (11%)
2	TRS	D	101	-	7,7,7	0.74	0	9,9,9	1.67	2 (22%)
3	GOL	C	101	-	5,5,5	0.50	0	5,5,5	0.74	0
3	GOL	B	101	-	5,5,5	0.34	0	5,5,5	0.49	0
2	TRS	B	104	-	7,7,7	0.50	0	9,9,9	1.21	1 (11%)
3	GOL	B	103	-	5,5,5	0.52	0	5,5,5	0.44	0
3	GOL	B	102	-	5,5,5	0.39	0	5,5,5	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRS	A	102	-	-	6/9/9/9	-
3	GOL	B	105	-	-	2/4/4/4	-
2	TRS	A	101	-	-	7/9/9/9	-
2	TRS	D	101	-	-	3/9/9/9	-
3	GOL	C	101	-	-	2/4/4/4	-
3	GOL	B	101	-	-	3/4/4/4	-
2	TRS	B	104	-	-	8/9/9/9	-
3	GOL	B	103	-	-	2/4/4/4	-
3	GOL	B	102	-	-	2/4/4/4	-

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	102	TRS	O3-C3-C	4.50	123.42	110.88
2	D	101	TRS	C2-C-N	3.00	115.83	108.17
2	D	101	TRS	C1-C-N	-2.74	101.19	108.17
2	A	101	TRS	C3-C-C1	-2.15	104.94	110.66
2	A	102	TRS	C3-C-N	2.06	113.43	108.17

There are no chirality outliers.

5 of 35 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	101	TRS	N-C-C2-O2
2	B	104	TRS	C2-C-C1-O1
2	B	104	TRS	C3-C-C1-O1
2	B	104	TRS	N-C-C1-O1
2	D	101	TRS	C1-C-C2-O2

There are no ring outliers.

8 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	102	TRS	2	0
2	A	101	TRS	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	101	TRS	3	0
3	C	101	GOL	2	0
3	B	101	GOL	1	0
2	B	104	TRS	4	0
3	B	103	GOL	4	0
3	B	102	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	91/94 (96%)	-0.68	0 100 100	17, 33, 52, 108	2 (2%)
1	B	92/94 (97%)	-0.71	1 (1%) 77 81	15, 33, 58, 99	4 (4%)
1	C	91/94 (96%)	-0.77	0 100 100	14, 33, 53, 93	4 (4%)
1	D	89/94 (94%)	-0.64	0 100 100	18, 38, 59, 65	3 (3%)
All	All	363/376 (96%)	-0.70	1 (0%) 90 92	14, 34, 58, 108	13 (3%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2	ALA	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	B	101	6/6	0.87	0.10	61,68,69,86	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	TRS	D	101	8/8	0.88	0.09	46,56,60,63	0
3	GOL	B	103	6/6	0.89	0.12	41,57,59,60	0
3	GOL	C	101	6/6	0.89	0.09	45,51,53,54	0
2	TRS	A	101	8/8	0.90	0.10	37,53,61,67	0
2	TRS	B	104	8/8	0.92	0.10	26,37,41,44	0
3	GOL	B	102	6/6	0.92	0.10	57,58,61,63	0
2	TRS	A	102	8/8	0.93	0.08	17,21,24,28	8
3	GOL	B	105	6/6	0.96	0.05	35,56,62,62	0

6.5 Other polymers [i](#)

There are no such residues in this entry.