

wwPDB EM Validation Summary Report (i)

Jul 2, 2023 – 12:15 AM JST

PDB ID : 8J26 EMDB ID : EMD-35945 Title : CryoEM structure of SARS CoV-2 RBD and Aptamer complex Authors Rahman, M.S.; Jang, S.K.; Lee, J.O. : Deposited on 2023-04-14 : 3.40 Å(reported) Resolution : Based on initial models 7KGJ, 6XDG :

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis		0.0.1.dev50 1.8.5 (274361), CSD as541be (2020)
0		
MolProbity	:	4.02b-467
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.9
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.33


1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.40 Å.

Sidechain outliers

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



154315

The table below summarises the geometric issues observed across the polymeric chains and their fit
to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues
that contain outliers for $>=3, 2, 1$ and 0 types of geometric quality criteria respectively. A grey
segment represents the fraction of residues that are not modelled. The numeric value for each
fraction is indicated below the corresponding segment, with a dot representing fractions $<=5\%$
The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM
map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

3826

Mol	Chain	Length	Quality of chain						
1	Е	44	7%	55%	32%	9% 5%			
2	А	224	39%	10%	51%				
3	В	248	36%	12%	51%				
4	С	253	•	9% •	23%				
5	D	48	•	67%	27%	6%			



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 5406 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a DNA chain called AM032-4.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	Е	42	Total 994	C 520	N 167	O 265	Р 42	0	0

• Molecule 2 is a protein called Fab light chain (REGN10987).

Mol	Chain	Residues	Atoms				AltConf	Trace	
2	А	109	Total	C	N	0	S	0	0
			797	495	133	166	3		

• Molecule 3 is a protein called Fab heavy chain (REGN10987).

Mol	Chain	Residues	Atoms				AltConf	Trace	
3	В	122	Total 942	$\begin{array}{c} \mathrm{C} \\ 595 \end{array}$	N 158	0 185	${S \over 4}$	0	0

• Molecule 4 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms				AltConf	Trace	
4	С	196	Total 1548	C 992	N 258	O 290	S 8	0	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	542	GLU	-	expression tag	UNP P0DTC2
С	543	ASN	-	expression tag	UNP P0DTC2
С	544	LEU	-	expression tag	UNP P0DTC2
С	545	TYR	-	expression tag	UNP P0DTC2
С	546	PHE	-	expression tag	UNP P0DTC2
С	547	GLN	-	expression tag	UNP P0DTC2
С	548	GLY	-	expression tag	UNP P0DTC2
С	549	ALA	-	expression tag	UNP P0DTC2
С	550	ALA	-	expression tag	UNP P0DTC2

Continued on next page...



	Continued from previous page									
Chain	Residue	Modelled	Actual	Comment	Reference					
C	551	ALA	-	expression tag	UNP P0DTC2					
C	552	GLY	-	expression tag	UNP P0DTC2					
С	553	GLY	-	expression tag	UNP P0DTC2					
С	554	SER	-	expression tag	UNP P0DTC2					
С	555	HIS	-	expression tag	UNP P0DTC2					
С	556	HIS	-	expression tag	UNP P0DTC2					
С	557	HIS	-	expression tag	UNP P0DTC2					
С	558	HIS	-	expression tag	UNP P0DTC2					
С	559	HIS	-	expression tag	UNP P0DTC2					
С	560	HIS	-	expression tag	UNP P0DTC2					
С	561	GLY	-	expression tag	UNP P0DTC2					
С	562	GLY	-	expression tag	UNP P0DTC2					
С	563	SER	-	expression tag	UNP P0DTC2					
С	564	ASP	-	expression tag	UNP P0DTC2					
С	565	TYR	-	expression tag	UNP P0DTC2					
С	566	LYS	-	expression tag	UNP P0DTC2					
С	567	ASP	-	expression tag	UNP P0DTC2					
С	568	ASP	-	expression tag	UNP P0DTC2					
С	569	ASP	-	expression tag	UNP P0DTC2					
С	570	ASP	-	expression tag	UNP P0DTC2					
С	571	LYS	-	expression tag	UNP P0DTC2					


Continued from previous page...

• Molecule 5 is a DNA chain called AM047-6.

Mol	Chain	Residues	Atoms				AltConf	Trace	
5	D	48	Total 1111	C 567	N 198	O 298	Р 48	0	0

• Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				AltConf
6	С	1	Total	С	Ν	0	0
0	U	1	14	8	1	5	0



Chain C:

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: AM032-4 Chain E: 55% 5% 32% 9% • Molecule 2: Fab light chain (REGN10987) Chain A: 39% 10% 51% PRO LYS ALA ALA ALA PRO PRO SER VAL • Molecule 3: Fab heavy chain (REGN10987) Chain B: 36% 12% 51% PHERE PRODUCT OF PRODUCT PROPERTING AND ADDRESS PRODUCT OF PRODUCT PROPERTING ADDRESS PRODUCT PROPERTING ADDRESS PRODUCT PROPERTING ADDRESS PRODUCT PROPERTING ADDRESS PRODUCT PRODUCT PROPERTING ADDRESS PRODUCT PRODUCT PROPERTING ADDRESS PROPERTING ADDRE • Molecule 4: Spike protein S1

68%

9%

23%

 \bullet Molecule 5: AM047-6





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	212000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	60	Depositor
Minimum defocus (nm)	700	Depositor
Maximum defocus (nm)	1600	Depositor
Magnification	105000	Depositor
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.058	Depositor
Minimum map value	-0.026	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.011	Depositor
Map size (Å)	314.496, 314.496, 314.496	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles ($^{\circ}$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.819, 0.819, 0.819	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: $85\mathrm{Y},$ NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	Е	0.51	0/740	0.75	0/1121	
2	А	0.25	0/814	0.49	0/1106	
3	В	0.28	0/963	0.53	0/1307	
4	С	0.28	0/1592	0.49	0/2168	
5	D	0.52	0/910	0.75	0/1385	
All	All	0.37	0/5019	0.60	0/7087	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Е	994	0	354	13	0
2	А	797	0	765	12	0
3	В	942	0	901	18	0
4	С	1548	0	1461	17	0
5	D	1111	0	430	9	0
6	С	14	0	13	3	0
All	All	5406	0	3924	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 64 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:35:DA:N6	4:C:494:SER:O	2.07	0.88
4:C:343:ASN:OD1	6:C:701:NAG:N2	2.17	0.77
1:E:28:DG:N2	1:E:59:85Y:O21	2.20	0.74
3:B:35:TYR:OH	3:B:104:ASP:O	2.07	0.73
1:E:36:DC:O2	4:C:505:TYR:OH	2.07	0.72

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
2	А	107/224~(48%)	94~(88%)	13 (12%)	0	100	100
3	В	120/248~(48%)	108 (90%)	12 (10%)	0	100	100
4	С	194/253~(77%)	186 (96%)	8 (4%)	0	100	100
All	All	421/725~(58%)	388 (92%)	33 (8%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	А	90/191~(47%)	87~(97%)	3~(3%)	38 66
3	В	100/212~(47%)	96~(96%)	4 (4%)	31 60
4	С	168/218~(77%)	161 (96%)	7 (4%)	30 59
All	All	358/621~(58%)	344 (96%)	14 (4%)	36 61

5 of 14 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
4	С	334	ASN
4	С	377	PHE
4	С	525	CYS
4	С	389	ASP
4	С	405	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
4	С	481	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

19 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Chain Res Linl		Bond lengths			Bond angles		
10101	туре	Chain	nes	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
1	85Y	Е	39	1	$33,\!36,\!37$	4.60	17 (51%)	47,51,54	<mark>3.60</mark>	10 (21%)
5	85Y	D	24	5	$33,\!36,\!37$	4.63	17 (51%)	47,51,54	<mark>3.54</mark>	9 (19%)
5	85Y	D	30	5	33,36,37	4.60	17 (51%)	47,51,54	3.73	11 (23%)



Mol	Type	Chain	Res	Link	В	ond leng	gths	B	ond ang	gles
	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
1	85Y	Е	51	1	$33,\!36,\!37$	4.63	17 (51%)	47,51,54	3.53	9 (19%)
5	85Y	D	17	5	$33,\!36,\!37$	4.57	17 (51%)	47,51,54	3.68	16 (34%)
5	85Y	D	34	5	33,36,37	4.61	17 (51%)	47,51,54	3.71	9 (19%)
5	85Y	D	43	5	33,36,37	4.63	17 (51%)	47,51,54	<mark>3.54</mark>	11 (23%)
5	85Y	D	20	5	33,36,37	4.63	17 (51%)	47,51,54	<mark>3.14</mark>	13 (27%)
1	85Y	Е	48	1	33,36,37	4.59	17 (51%)	47,51,54	<mark>3.54</mark>	<mark>9 (19%)</mark>
1	85Y	Е	44	1	33,36,37	4.63	17 (51%)	47,51,54	<mark>3.59</mark>	12 (25%)
1	85Y	Е	34	1	33,36,37	4.61	17 (51%)	47,51,54	<mark>3.58</mark>	13 (27%)
1	85Y	Е	55	1	33,36,37	4.59	17 (51%)	47,51,54	<mark>3.51</mark>	13 (27%)
5	85Y	D	22	5	33,36,37	4.62	17 (51%)	47,51,54	3.40	13 (27%)
1	85Y	Е	59	1	33,36,37	4.65	17 (51%)	47,51,54	<mark>3.59</mark>	9 (19%)
1	85Y	Е	38	1	33,36,37	4.57	17 (51%)	47,51,54	<mark>3.63</mark>	14 (29%)
5	85Y	D	41	5	33,36,37	4.59	17 (51%)	47,51,54	<mark>3.68</mark>	12 (25%)
5	85Y	D	35	5	33,36,37	4.56	17 (51%)	47,51,54	<mark>3.60</mark>	16 (34%)
1	85Y	Е	53	1	33,36,37	4.59	17 (51%)	47,51,54	<mark>3.60</mark>	13 (27%)
1	85Y	Е	33	1	33,36,37	4.64	16 (48%)	47,51,54	3.58	14 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	85Y	Е	39	1	-	4/16/30/31	0/4/4/4
5	85Y	D	24	5	-	9/16/30/31	0/4/4/4
5	85Y	D	30	5	-	8/16/30/31	0/4/4/4
1	85Y	Е	51	1	-	8/16/30/31	0/4/4/4
5	85Y	D	17	5	-	8/16/30/31	0/4/4/4
5	85Y	D	34	5	-	8/16/30/31	0/4/4/4
5	85Y	D	43	5	-	11/16/30/31	0/4/4/4
5	85Y	D	20	5	-	1/16/30/31	0/4/4/4
1	85Y	Е	48	1	-	10/16/30/31	0/4/4/4
1	85Y	Е	44	1	-	0/16/30/31	0/4/4/4
1	85Y	Е	34	1	-	10/16/30/31	0/4/4/4
1	85Y	Е	55	1	-	9/16/30/31	0/4/4/4
5	85Y	D	22	5	-	11/16/30/31	0/4/4/4

Continued on next page...



Mol	Type	Chain	\mathbf{Res}	Link	Chirals	Torsions	Rings
1	85Y	Е	59	1	-	8/16/30/31	0/4/4/4
1	85Y	Е	38	1	-	8/16/30/31	0/4/4/4
5	85Y	D	41	5	-	9/16/30/31	0/4/4/4
5	85Y	D	35	5	-	8/16/30/31	0/4/4/4
1	85Y	Е	53	1	-	9/16/30/31	0/4/4/4
1	85Y	Е	33	1	-	7/16/30/31	0/4/4/4

Continued from previous page...

The worst 5 of 322 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
5	D	30	85Y	C5-C4	12.59	1.68	1.45
1	Е	33	85Y	C5-C4	12.48	1.67	1.45
5	D	20	85Y	C5-C4	12.47	1.67	1.45
5	D	22	85Y	C5-C4	12.46	1.67	1.45
1	Е	59	85Y	C5-C4	12.43	1.67	1.45

The worst 5 of 226 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	D	34	85Y	C23-C24-C25	15.99	150.13	121.51
5	D	30	85Y	C23-C24-C25	15.70	149.62	121.51
1	Е	48	85Y	C23-C24-C25	15.28	148.87	121.51
5	D	43	85Y	C23-C24-C25	15.23	148.78	121.51
5	D	41	85Y	C23-C24-C25	15.22	148.75	121.51

There are no chirality outliers.

5 of 146 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	Е	33	85Y	O21-C20-C5-C4
1	Е	33	85Y	O21-C20-C5-C6
1	Е	33	85Y	N22-C20-C5-C6
1	Е	33	85Y	C5-C20-N22-C23
1	Е	33	85Y	O21-C20-N22-C23

There are no ring outliers.

7 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	34	85Y	1	0



Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	43	85Y	1	0
1	Е	55	85Y	2	0
1	Е	59	85Y	1	0
5	D	35	85Y	1	0
1	Е	53	85Y	1	0
1	Е	33	85Y	1	0

Continued from previous page...

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Dec	Link	Bo	ond leng	\mathbf{ths}	В	ond ang	les
MOI	туре	Unain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
6	NAG	C	701	4	14,14,15	0.78	0	$17,\!19,\!21$	0.95	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mo	l Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	С	701	4	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
6	С	701	NAG	C1-O5-C5	2.10	115.04	112.19



There are no chirality outliers.

Mol	Chain	Res	Type	Atoms
6	С	701	NAG	O5-C5-C6-O6
6	С	701	NAG	C4-C5-C6-O6
6	С	701	NAG	C1-C2-N2-C7
6	С	701	NAG	C3-C2-N2-C7

All (4) torsion outliers are listed below:


There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	С	701	NAG	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-35945. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map




The images above show the map projected in three orthogonal directions.




6.2 Central slices (i)

6.2.1 Primary map



X Index: 192




Y Index: 192



Z Index: 192

6.2.2 Raw map



X Index: 192

Y Index: 192




The images above show central slices of the map in three orthogonal directions.




6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 189



Y Index: 201



Z Index: 191

6.3.2 Raw map



X Index: 189

Y Index: 197




The images above show the largest variance slices of the map in three orthogonal directions.




6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



6.4.2 Raw map




The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.011. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is 46 $\rm nm^3;$ this corresponds to an approximate mass of 42 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.294 \AA^{-1}



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.294 $\mathrm{\AA^{-1}}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estim	ation	criterion (FSC cut-off)
Resolution estimate (A)	0.143	0.5	Half-bit
Reported by author	3.40	-	-
Author-provided FSC curve	3.43	4.01	3.66
Unmasked-calculated*	3.41	3.83	3.48


*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-35945 and PDB model 8J26. Per-residue inclusion information can be found in section 3 on page 6.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.011 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.011).



9.4 Atom inclusion (i)



At the recommended contour level, 89% of all backbone atoms, 83% of all non-hydrogen atoms, are inside the map.



1.0

0.0 <0.0

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.011) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8260	0.4280
А	0.8640	0.4840
В	0.8820	0.4960
С	0.8570	0.5080
D	0.7990	0.3250
Ε	0.7260	0.3090

