



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 1, 2023 – 02:53 PM JST

PDB ID : 5J4T  
Title : Structure of tetrameric jacalin complexed with GlcNAc beta-(1,3) Gal-beta-OMe  
Authors : Abhinav, K.V.; Sharma, K.; Surolia, A.; Vijayan, M.  
Deposited on : 2016-04-01  
Resolution : 1.94 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4310 (1.96-1.92)
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)
RSRZ outliers	127900	4250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	133	 92% 8%
1	C	133	 92% 8%
1	E	133	 91% 9%
1	G	133	 92% 8%
2	B	19	 74% 11% 16%
2	D	19	 5% 68% 5% 26%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	19	
2	H	19	
3	I	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	PEG	E	202	-	-	X	-

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 5043 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Agglutinin alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	133	1037	678	159	198	2	0	0	0
1	C	133	1037	677	159	199	2	0	0	0
1	E	133	1041	680	160	199	2	0	1	0
1	G	133	1034	675	159	198	2	0	0	0

- Molecule 2 is a protein called Agglutinin beta-3 chain.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	16	114	73	20	21	0	0	0
2	D	14	96	62	16	18	0	0	0
2	F	16	110	71	19	20	0	0	0
2	H	15	100	66	17	17	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-methyl beta-D-galactopyranoside.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	I	2	27	15	1	11	0	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	F	1	Total C O 4 2 2	0	0
4	G	1	Total C O 4 2 2	0	0
4	G	1	Total C O 4 2 2	0	0
4	H	1	Total C O 4 2 2	0	0

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	E	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is methyl beta-D-galactopyranoside (three-letter code: MBG) (formula:  $C_7H_{14}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	G	1	Total	C	O	0	0
			13	7	6		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	96	Total O 96 96	0	0
7	B	13	Total O 13 13	0	0
7	C	85	Total O 85 85	0	0
7	D	7	Total O 7 7	0	0
7	E	87	Total O 87 87	0	0
7	F	9	Total O 9 9	0	0
7	G	86	Total O 86 86	0	0
7	H	12	Total O 12 12	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Agglutinin alpha chain

Chain A:  92% 8%



- Molecule 1: Agglutinin alpha chain

Chain C:  92% 8%



- Molecule 1: Agglutinin alpha chain

Chain E:  91% 9%



- Molecule 1: Agglutinin alpha chain

Chain G:  92% 8%



- Molecule 2: Agglutinin beta-3 chain


Chain B:  74% 11% 16%




- Molecule 2: Agglutinin beta-3 chain

Chain D:  5% 68% 5% 26%






- Molecule 2: Agglutinin beta-3 chain



- Molecule 2: Agglutinin beta-3 chain



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-methyl beta-D-galactopyranoside



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.31Å 80.65Å 62.85Å 90.00° 107.90° 90.00°	Depositor
Resolution (Å)	48.45 – 1.94 48.04 – 1.94	Depositor EDS
% Data completeness (in resolution range)	99.5 (48.45-1.94) 99.5 (48.04-1.94)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.81 (at 1.94Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.176 , 0.225 0.175 , 0.225	Depositor DCC
$R_{free}$ test set	4055 reflections (9.97%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.8	Xtrriage
Anisotropy	0.854	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 48.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.42$ , $\langle L^2 \rangle = 0.24$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5043	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, MBG, NAG, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.94	0/1066	0.95	2/1445 (0.1%)
1	C	0.99	1/1066 (0.1%)	0.94	0/1446
1	E	0.99	0/1070	0.94	1/1451 (0.1%)
1	G	1.02	1/1063 (0.1%)	0.92	0/1441
2	B	1.18	0/116	0.89	0/157
2	D	1.05	0/98	0.91	0/134
2	F	1.16	0/112	1.03	0/152
2	H	1.03	0/102	0.85	0/139
All	All	1.00	2/4693 (0.0%)	0.94	3/6365 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	123	TRP	CD2-CE2	6.49	1.49	1.41
1	G	32	TYR	CE1-CZ	5.16	1.45	1.38

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	27	ASP	CB-CG-OD2	-6.14	112.78	118.30
1	A	27	ASP	CB-CG-OD1	6.01	123.71	118.30
1	E	5	ASP	CB-CG-OD1	5.04	122.84	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1037	0	1005	8	0
1	C	1037	0	1000	8	0
1	E	1041	0	1004	14	0
1	G	1034	0	996	7	0
2	B	114	0	116	3	0
2	D	96	0	92	1	0
2	F	110	0	110	2	0
2	H	100	0	99	2	0
3	I	27	0	26	4	0
4	A	12	0	18	1	0
4	E	4	0	6	3	0
4	F	4	0	6	0	0
4	G	8	0	12	0	0
4	H	4	0	6	3	0
5	E	7	0	10	6	0
6	G	13	0	14	1	0
7	A	96	0	0	0	0
7	B	13	0	0	0	0
7	C	85	0	0	2	0
7	D	7	0	0	0	0
7	E	87	0	0	2	0
7	F	9	0	0	0	0
7	G	86	0	0	0	0
7	H	12	0	0	1	0
All	All	5043	0	4520	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 44 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:3:ALA:HB1	5:E:202:PEG:H12	1.46	0.94
1:E:3:ALA:CB	5:E:202:PEG:H12	2.21	0.68
1:E:3:ALA:HB1	5:E:202:PEG:C3	2.23	0.68

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:3:ALA:HB1	5:E:202:PEG:H31	1.81	0.63
1:E:45[B]:LYS:NZ	4:E:201:EDO:H21	2.15	0.61

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	131/133 (98%)	127 (97%)	4 (3%)	0	100	100
1	C	131/133 (98%)	127 (97%)	4 (3%)	0	100	100
1	E	132/133 (99%)	128 (97%)	4 (3%)	0	100	100
1	G	131/133 (98%)	125 (95%)	6 (5%)	0	100	100
2	B	14/19 (74%)	14 (100%)	0	0	100	100
2	D	12/19 (63%)	12 (100%)	0	0	100	100
2	F	14/19 (74%)	14 (100%)	0	0	100	100
2	H	13/19 (68%)	13 (100%)	0	0	100	100
All	All	578/608 (95%)	560 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	112/113 (99%)	112 (100%)	0	100	100
1	C	112/113 (99%)	111 (99%)	1 (1%)	78	75
1	E	112/113 (99%)	111 (99%)	1 (1%)	78	75
1	G	111/113 (98%)	110 (99%)	1 (1%)	78	75
2	B	12/15 (80%)	12 (100%)	0	100	100
2	D	10/15 (67%)	10 (100%)	0	100	100
2	F	11/15 (73%)	11 (100%)	0	100	100
2	H	9/15 (60%)	9 (100%)	0	100	100
All	All	489/512 (96%)	486 (99%)	3 (1%)	86	85

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	100	SER
1	E	74	ASN
1	G	10	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	MBG	I	1	3	13,13,13	2.65	4 (30%)	18,18,18	2.23	7 (38%)
3	NAG	I	2	3	14,14,15	0.76	0	17,19,21	1.52	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MBG	I	1	3	-	2/4/24/24	0/1/1/1
3	NAG	I	2	3	-	2/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	I	1	MBG	O1-C1	5.87	1.50	1.40
3	I	1	MBG	C4-C5	5.68	1.65	1.53
3	I	1	MBG	O5-C1	3.48	1.50	1.41
3	I	1	MBG	O5-C5	2.02	1.49	1.44

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	1	MBG	C1-O5-C5	-4.73	104.41	113.69
3	I	1	MBG	O1-C1-C2	4.57	113.51	108.15
3	I	1	MBG	C7-O1-C1	-3.84	107.34	113.27
3	I	1	MBG	O5-C1-O1	2.91	117.72	110.97
3	I	1	MBG	O5-C5-C6	2.81	113.43	106.44

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	I	1	MBG	C2-C1-O1-C7
3	I	2	NAG	O5-C5-C6-O6
3	I	1	MBG	C4-C5-C6-O6
3	I	2	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	1	MBG	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	EDO	A	201	-	3,3,3	0.38	0	2,2,2	0.76	0
4	EDO	A	202	-	3,3,3	0.37	0	2,2,2	0.80	0
4	EDO	E	201	-	3,3,3	0.39	0	2,2,2	0.64	0
4	EDO	G	203	-	3,3,3	0.30	0	2,2,2	0.91	0
6	MBG	G	201	-	13,13,13	2.71	5 (38%)	18,18,18	3.11	8 (44%)
4	EDO	H	101	-	3,3,3	0.62	0	2,2,2	0.68	0
4	EDO	A	203	-	3,3,3	0.31	0	2,2,2	1.00	0
5	PEG	E	202	-	6,6,6	0.63	0	5,5,5	0.88	0
4	EDO	G	202	-	3,3,3	0.51	0	2,2,2	0.40	0
4	EDO	F	101	-	3,3,3	0.41	0	2,2,2	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	201	-	-	1/1/1/1	-
4	EDO	A	202	-	-	0/1/1/1	-
4	EDO	E	201	-	-	0/1/1/1	-
4	EDO	G	203	-	-	1/1/1/1	-
6	MBG	G	201	-	-	2/4/24/24	0/1/1/1
4	EDO	H	101	-	-	0/1/1/1	-
4	EDO	A	203	-	-	1/1/1/1	-
5	PEG	E	202	-	-	2/4/4/4	-
4	EDO	G	202	-	-	0/1/1/1	-
4	EDO	F	101	-	-	1/1/1/1	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	G	201	MBG	O5-C1	5.60	1.56	1.41
6	G	201	MBG	C4-C3	4.55	1.63	1.52
6	G	201	MBG	C4-C5	4.33	1.62	1.53
6	G	201	MBG	O4-C4	3.39	1.50	1.43
6	G	201	MBG	O1-C7	2.60	1.51	1.42

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	G	201	MBG	C1-O5-C5	-8.83	96.36	113.69

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	G	201	MBG	O5-C1-O1	5.85	124.53	110.97
6	G	201	MBG	C1-C2-C3	-3.80	102.09	110.00
6	G	201	MBG	O3-C3-C4	3.57	118.61	110.35
6	G	201	MBG	C7-O1-C1	-3.15	108.41	113.27

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	G	201	MBG	O5-C1-O1-C7
6	G	201	MBG	C2-C1-O1-C7
5	E	202	PEG	O2-C3-C4-O4
5	E	202	PEG	C1-C2-O2-C3
4	F	101	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	202	EDO	1	0
4	E	201	EDO	3	0
6	G	201	MBG	1	0
4	H	101	EDO	3	0
5	E	202	PEG	6	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	133/133 (100%)	-0.48	0 <a href="#">100</a>   <a href="#">100</a>	10, 16, 24, 31	0
1	C	133/133 (100%)	-0.40	1 (0%) <a href="#">86</a>   <a href="#">89</a>	12, 18, 31, 38	0
1	E	133/133 (100%)	-0.54	0 <a href="#">100</a>   <a href="#">100</a>	11, 16, 25, 38	0
1	G	133/133 (100%)	-0.47	0 <a href="#">100</a>   <a href="#">100</a>	11, 16, 29, 34	0
2	B	16/19 (84%)	-0.12	0 <a href="#">100</a>   <a href="#">100</a>	14, 20, 32, 46	0
2	D	14/19 (73%)	-0.05	1 (7%) <a href="#">16</a>   <a href="#">22</a>	15, 20, 31, 46	0
2	F	16/19 (84%)	-0.36	0 <a href="#">100</a>   <a href="#">100</a>	11, 17, 38, 41	0
2	H	15/19 (78%)	-0.23	0 <a href="#">100</a>   <a href="#">100</a>	14, 18, 30, 48	0
All	All	593/608 (97%)	-0.44	2 (0%) <a href="#">94</a>   <a href="#">96</a>	10, 17, 29, 48	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	3	GLN	3.9
1	C	98	VAL	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MBG	I	1	13/13	0.92	0.13	14,23,30,31	0
3	NAG	I	2	14/15	0.92	0.21	25,30,33,33	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	PEG	E	202	7/7	0.84	0.27	25,28,29,35	0
6	MBG	G	201	13/13	0.85	0.18	17,20,25,26	0
4	EDO	G	202	4/4	0.87	0.23	36,37,38,38	0
4	EDO	H	101	4/4	0.87	0.15	34,34,35,36	0
4	EDO	G	203	4/4	0.89	0.13	48,49,49,50	0
4	EDO	A	201	4/4	0.91	0.17	37,38,39,39	0
4	EDO	F	101	4/4	0.92	0.15	36,38,39,42	0
4	EDO	A	202	4/4	0.92	0.14	33,33,33,34	0
4	EDO	A	203	4/4	0.93	0.26	39,39,40,42	0
4	EDO	E	201	4/4	0.94	0.13	34,34,35,35	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.