



wwPDB X-ray Structure Validation Summary Report ⓘ

May 1, 2024 – 08:11 PM JST

PDB ID : 8J9B
Title : LnaB-actin binary complex
Authors : Chen, T.T.; Ouyang, S.Y.
Deposited on : 2023-05-03
Resolution : 3.42 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<https://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1486 (3.50-3.34)
Clashscore	141614	1572 (3.50-3.34)
Ramachandran outliers	138981	1534 (3.50-3.34)
Sidechain outliers	138945	1535 (3.50-3.34)
RSRZ outliers	127900	1395 (3.50-3.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	375	
1	B	375	
1	C	375	
1	D	375	
1	E	375	
1	F	375	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	441	
2	I	441	
2	J	441	
2	K	441	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 26804 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Actin gamma 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	322	2538	1619	420	479	20	0	0	0
1	B	343	2683	1706	446	511	20	0	0	0
1	C	342	2673	1700	439	514	20	0	0	0
1	D	354	2774	1763	462	529	20	0	0	0
1	E	318	2488	1585	410	474	19	0	0	0
1	F	338	2641	1676	443	502	20	0	0	0


- Molecule 2 is a protein called Type IV secretion protein Dot.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	348	2826	1803	462	556	5	0	0	0
2	I	335	2707	1723	444	535	5	0	0	0
2	J	343	2780	1772	457	546	5	0	0	0
2	K	331	2694	1726	442	521	5	0	0	0


- Molecule 1: Actin gamma 1




- Molecule 1: Actin gamma 1




- Molecule 1: Actin gamma 1







- Molecule 1: Actin gamma 1



- Molecule 2: Type IV secretion protein Dot



- Molecule 2: Type IV secretion protein Dot



4 Data and refinement statistics i

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	110.58Å 110.58Å 407.93Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	62.10 – 3.42 135.98 – 3.42	Depositor EDS
% Data completeness (in resolution range)	100.0 (62.10-3.42) 99.9 (135.98-3.42)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.82 (at 3.41Å)	Xtrriage
Refinement program	PHENIX (1.19.2_4158: ???)	Depositor
R, R_{free}	0.287 , 0.298 0.289 , 0.289	Depositor DCC
R_{free} test set	3809 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	73.3	Xtrriage
Anisotropy	0.006	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 49.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.019 for -h,-k,l 0.018 for h,-h-k,-l 0.000 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.80	EDS
Total number of atoms	26804	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.81	0/2591	0.89	2/3502 (0.1%)
1	B	0.87	0/2740	0.88	0/3706
1	C	1.03	0/2730	0.93	5/3697 (0.1%)
1	D	1.03	0/2833	0.92	6/3835 (0.2%)
1	E	0.99	0/2541	0.88	3/3438 (0.1%)
1	F	1.00	0/2696	0.89	0/3646
2	H	1.04	0/2881	0.90	3/3889 (0.1%)
2	I	1.01	0/2755	0.93	5/3713 (0.1%)
2	J	0.85	0/2831	0.85	4/3818 (0.1%)
2	K	0.72	0/2745	0.72	2/3698 (0.1%)
All	All	0.94	0/27343	0.88	30/36942 (0.1%)

There are no bond length outliers.

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	K	187	LEU	CA-CB-CG	5.99	129.08	115.30
2	I	137	GLU	N-CA-C	-5.95	94.95	111.00
2	J	228	LEU	N-CA-C	-5.95	94.94	111.00
1	D	98	PRO	N-CA-C	5.82	127.23	112.10
2	H	138	SER	N-CA-C	5.81	126.68	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2538	0	2511	62	0
1	B	2683	0	2645	100	0
1	C	2673	0	2626	48	0
1	D	2774	0	2746	60	0
1	E	2488	0	2443	49	0
1	F	2641	0	2618	52	0
2	H	2826	0	2816	34	0
2	I	2707	0	2686	56	0
2	J	2780	0	2779	47	0
2	K	2694	0	2696	36	0
All	All	26804	0	26566	540	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 540 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:157:TYR:CD2	2:H:180:MET:HE3	1.51	1.45
1:B:105:LEU:HD21	1:B:123:MET:CE	1.69	1.21
1:B:143:TYR:CD2	1:B:346:LEU:HD11	1.77	1.19
2:H:157:TYR:CD2	2:H:180:MET:CE	2.26	1.18
2:H:157:TYR:HD2	2:H:180:MET:CE	1.57	1.17

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	310/375 (83%)	292 (94%)	17 (6%)	1 (0%)	41 74
1	B	335/375 (89%)	323 (96%)	12 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	332/375 (88%)	316 (95%)	16 (5%)	0	100	100
1	D	346/375 (92%)	328 (95%)	18 (5%)	0	100	100
1	E	304/375 (81%)	283 (93%)	21 (7%)	0	100	100
1	F	330/375 (88%)	318 (96%)	12 (4%)	0	100	100
2	H	344/441 (78%)	321 (93%)	23 (7%)	0	100	100
2	I	325/441 (74%)	301 (93%)	24 (7%)	0	100	100
2	J	337/441 (76%)	315 (94%)	21 (6%)	1 (0%)	41	74
2	K	321/441 (73%)	305 (95%)	16 (5%)	0	100	100
All	All	3284/4014 (82%)	3102 (94%)	180 (6%)	2 (0%)	51	83

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	70	PRO
2	J	214	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	275/318 (86%)	261 (95%)	14 (5%)	24	57
1	B	289/318 (91%)	277 (96%)	12 (4%)	30	62
1	C	292/318 (92%)	282 (97%)	10 (3%)	37	68
1	D	302/318 (95%)	291 (96%)	11 (4%)	35	66
1	E	270/318 (85%)	259 (96%)	11 (4%)	30	62
1	F	287/318 (90%)	275 (96%)	12 (4%)	30	62
2	H	322/408 (79%)	303 (94%)	19 (6%)	19	52
2	I	308/408 (76%)	296 (96%)	12 (4%)	32	64
2	J	317/408 (78%)	308 (97%)	9 (3%)	43	73
2	K	305/408 (75%)	299 (98%)	6 (2%)	55	79

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	2967/3540 (84%)	2851 (96%)	116 (4%)	32 64

5 of 116 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	334	GLU
2	K	21	TYR
2	H	59	GLU
2	J	299	TYR
2	I	331	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
2	I	74	GLN
2	I	331	ASN
2	I	320	GLN
2	I	346	GLN
1	B	360	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	322/375 (85%)	1.29	77 (23%) 0 0	30, 96, 106, 116	0
1	B	343/375 (91%)	1.00	52 (15%) 2 3	40, 84, 100, 161	0
1	C	342/375 (91%)	0.55	20 (5%) 23 24	30, 43, 57, 75	0
1	D	354/375 (94%)	0.53	18 (5%) 28 28	30, 46, 58, 78	0
1	E	318/375 (84%)	0.68	30 (9%) 8 11	32, 49, 63, 77	0
1	F	338/375 (90%)	0.68	22 (6%) 18 21	38, 52, 62, 92	0
2	H	348/441 (78%)	0.49	18 (5%) 27 28	30, 47, 65, 88	0
2	I	335/441 (75%)	0.56	26 (7%) 13 16	33, 46, 61, 77	0
2	J	343/441 (77%)	0.46	13 (3%) 40 40	27, 43, 59, 75	0
2	K	331/441 (75%)	0.69	31 (9%) 8 11	36, 52, 72, 515	0
All	All	3374/4014 (84%)	0.69	307 (9%) 9 12	27, 50, 98, 515	0

The worst 5 of 307 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	J	220	ASN	7.1
1	A	271	SER	6.5
1	A	272	CYS	5.9
1	A	159	VAL	5.8
1	C	250	ILE	5.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.