



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 11, 2024 – 06:10 PM JST

PDB ID : 8JZ2  
Title : Crystal structure of AetF in complex with FAD  
Authors : Li, H.; Dai, L.; Chen, C.-C.; Guo, R.-T.  
Deposited on : 2023-07-04  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	679	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5661 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called AetF.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	629	5207	3348	880	964	15	0	0	0

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	GLY	-	expression tag	UNP A0A861B9Z9
A	-9	ALA	-	expression tag	UNP A0A861B9Z9
A	-8	GLY	-	expression tag	UNP A0A861B9Z9
A	-7	ALA	-	expression tag	UNP A0A861B9Z9
A	-6	GLY	-	expression tag	UNP A0A861B9Z9
A	-5	ALA	-	expression tag	UNP A0A861B9Z9
A	-4	GLY	-	expression tag	UNP A0A861B9Z9
A	-3	ALA	-	expression tag	UNP A0A861B9Z9
A	-2	GLY	-	expression tag	UNP A0A861B9Z9
A	-1	ALA	-	expression tag	UNP A0A861B9Z9
A	0	GLY	-	expression tag	UNP A0A861B9Z9

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	401	401	401	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.48Å 74.96Å 143.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.47 – 1.85 24.45 – 1.85	Depositor EDS
% Data completeness (in resolution range)	95.8 (24.47-1.85) 95.9 (24.45-1.85)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.90 (at 1.85Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, $R_{free}$	0.178 , 0.229 0.185 , 0.236	Depositor DCC
$R_{free}$ test set	2724 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.7	Xtrriage
Anisotropy	0.645	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 41.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5661	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.28% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.71	0/5342	0.82	0/7236

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5207	0	5070	60	0
2	A	53	0	31	0	0
3	A	401	0	0	9	0
All	All	5661	0	5101	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:GLY:HA2	1:A:257:ILE:HD11	1.67	0.77

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:633:GLN:HE21	1:A:633:GLN:HA	1.59	0.67
1:A:46:ARG:NH2	1:A:141:PHE:O	2.29	0.66
1:A:305:ASN:HD21	1:A:307:ASP:HB3	1.63	0.62
1:A:553:ASP:HB3	1:A:556:VAL:HG13	1.80	0.62
1:A:134:MET:HG2	1:A:327:ASP:HB2	1.83	0.61
1:A:114:ARG:HD3	3:A:836:HOH:O	2.00	0.60
1:A:560:GLU:O	1:A:564:MET:HG3	2.02	0.59
1:A:253:GLY:HA2	1:A:424:HIS:O	2.03	0.59
1:A:633:GLN:HA	1:A:633:GLN:NE2	2.19	0.58
1:A:1:MET:N	3:A:813:HOH:O	2.38	0.56
1:A:220:VAL:HG11	1:A:254:LEU:HD11	1.87	0.56
1:A:304:ILE:H	1:A:304:ILE:HD13	1.69	0.56
1:A:169:ASN:OD1	1:A:550:ILE:HD13	2.05	0.56
1:A:146:SER:O	1:A:148:LYS:HD3	2.07	0.55
1:A:176:ARG:NH2	1:A:304:ILE:HD12	2.22	0.55
1:A:134:MET:CG	1:A:327:ASP:HB2	2.38	0.54
1:A:134:MET:HG2	1:A:327:ASP:CB	2.37	0.53
1:A:192:LYS:HB3	3:A:898:HOH:O	2.08	0.53
1:A:323:TYR:CE2	1:A:325:LYS:HG3	2.44	0.53
1:A:453:VAL:HA	1:A:606:ILE:HD11	1.90	0.53
1:A:486:LYS:NZ	3:A:818:HOH:O	2.42	0.53
1:A:134:MET:CG	1:A:327:ASP:CB	2.88	0.52
1:A:305:ASN:C	1:A:305:ASN:HD22	2.11	0.51
1:A:188:GLN:HE21	1:A:188:GLN:HA	1.76	0.50
1:A:345:GLU:OE2	1:A:347:LEU:HD23	2.12	0.50
1:A:323:TYR:HE2	1:A:325:LYS:HG3	1.75	0.49
1:A:156:GLY:HA2	1:A:257:ILE:CD1	2.39	0.49
1:A:365:PHE:HB3	1:A:368:TYR:CD2	2.48	0.48
1:A:332:ARG:NE	3:A:824:HOH:O	2.46	0.48
1:A:327:ASP:O	1:A:327:ASP:CG	2.53	0.47
1:A:153:ASP:OD1	1:A:326:GLY:N	2.48	0.47
1:A:444:HIS:H	1:A:444:HIS:CD2	2.32	0.47
1:A:189:VAL:CG1	1:A:496:GLN:HG3	2.46	0.46
1:A:188:GLN:HE21	1:A:188:GLN:CA	2.28	0.45
1:A:531:GLU:OE1	1:A:534:ARG:HD2	2.17	0.45
1:A:134:MET:HG3	1:A:327:ASP:HB3	1.99	0.45
1:A:420:LYS:HD2	3:A:1193:HOH:O	2.16	0.45
1:A:397:HIS:HD2	3:A:1147:HOH:O	2.00	0.44
1:A:304:ILE:HG22	1:A:311:ILE:HG13	2.00	0.44
1:A:173:ILE:HB	1:A:295:VAL:HG22	1.99	0.44
1:A:490:LYS:N	1:A:490:LYS:HE3	2.33	0.44

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:414:HIS:HE1	3:A:937:HOH:O	2.00	0.44
1:A:134:MET:HG3	1:A:327:ASP:CB	2.49	0.43
1:A:593:LYS:HA	1:A:596:ARG:HG2	1.99	0.43
1:A:169:ASN:OD1	1:A:550:ILE:CD1	2.67	0.43
1:A:4:VAL:O	1:A:27:PHE:HA	2.19	0.43
1:A:332:ARG:CD	3:A:824:HOH:O	2.67	0.42
1:A:192:LYS:HA	1:A:193:PRO:HD3	1.94	0.42
1:A:134:MET:CG	1:A:327:ASP:HB3	2.48	0.42
1:A:420:LYS:HG3	1:A:421:PRO:HD2	2.01	0.42
1:A:588:ASP:O	1:A:592:ASN:HB2	2.20	0.41
1:A:370:ARG:O	1:A:427:THR:HA	2.20	0.41
1:A:156:GLY:CA	1:A:257:ILE:HD11	2.42	0.41
1:A:289:TRP:HB3	1:A:295:VAL:HG23	2.02	0.41
1:A:636:GLU:C	1:A:640:ARG:HE	2.24	0.41
1:A:338:ILE:HD13	1:A:359:ARG:NH2	2.36	0.40
1:A:339:ASN:ND2	1:A:341:GLU:OE1	2.55	0.40
1:A:453:VAL:CA	1:A:606:ILE:HD11	2.52	0.40
1:A:186:GLU:CD	1:A:268:HIS:HE2	2.25	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	623/679 (92%)	606 (97%)	17 (3%)	0	<a href="#">100</a> <a href="#">100</a>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	571/603 (95%)	540 (95%)	31 (5%)	22 8

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	93	LYS
1	A	138	LEU
1	A	143	ASN
1	A	184	ASP
1	A	188	GLN
1	A	190	LEU
1	A	202	PRO
1	A	225	VAL
1	A	231	TYR
1	A	236	LEU
1	A	254	LEU
1	A	304	ILE
1	A	305	ASN
1	A	316	ILE
1	A	327	ASP
1	A	347	LEU
1	A	420	LYS
1	A	426	HIS
1	A	457	LEU
1	A	490	LYS
1	A	496	GLN
1	A	550	ILE
1	A	556	VAL
1	A	557	VAL
1	A	566	LYS
1	A	588	ASP
1	A	593	LYS
1	A	596	ARG
1	A	598	ILE
1	A	636	GLU
1	A	640	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	GLN
1	A	140	ASN
1	A	170	ASN
1	A	188	GLN
1	A	264	GLN
1	A	305	ASN
1	A	401	HIS
1	A	414	HIS
1	A	444	HIS
1	A	446	GLN
1	A	529	HIS
1	A	562	ASN
1	A	633	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FAD	A	700	-	53,58,58	0.77	0	68,89,89	0.93	3 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	700	-	-	2/30/50/50	0/6/6/6

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	700	FAD	C4-N3-C2	-2.17	121.63	125.64
2	A	700	FAD	O4B-C1B-C2B	-2.17	103.76	106.93
2	A	700	FAD	O2P-P-O1P	2.05	122.37	112.24

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	700	FAD	N10-C1'-C2'-C3'
2	A	700	FAD	O4B-C4B-C5B-O5B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	629/679 (92%)	0.22	33 (5%) <span style="border: 1px solid red; padding: 2px;">27</span> <span style="border: 1px solid red; padding: 2px;">26</span>	17, 33, 63, 96	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	144	HIS	6.3
1	A	577	PRO	5.9
1	A	314	ALA	4.4
1	A	630	PHE	4.2
1	A	575	VAL	4.2
1	A	578	ASN	4.0
1	A	579	ILE	3.4
1	A	549	ASN	3.2
1	A	589	GLU	2.9
1	A	303	PRO	2.9
1	A	574	LYS	2.7
1	A	612	SER	2.7
1	A	191	GLY	2.6
1	A	341	GLU	2.6
1	A	635	ILE	2.5
1	A	520	ARG	2.5
1	A	552	VAL	2.5
1	A	653	ARG	2.4
1	A	596	ARG	2.4
1	A	645	LYS	2.4
1	A	315	GLY	2.4
1	A	631	GLU	2.4
1	A	240	ASP	2.3
1	A	573	ASP	2.2
1	A	304	ILE	2.2
1	A	506	ASN	2.2
1	A	317	GLU	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	633	GLN	2.1
1	A	125	VAL	2.1
1	A	551	SER	2.1
1	A	339	ASN	2.0
1	A	563	PHE	2.0
1	A	570	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	FAD	A	700	53/53	0.98	0.08	17,21,27,33	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.