



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 20, 2023 – 06:00 PM EDT

PDB ID : 5K1A  
Title : Crystal structure of the UAF1-USP12 complex in C2 space group  
Authors : Li, H.; D'Andrea, A.D.; Zheng, N.  
Deposited on : 2016-05-18  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 63%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 28%; height: 10px; background-color: grey;"></div> </div>
1	C	331	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 59%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 29%; height: 10px; background-color: grey;"></div> </div>
1	E	331	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 58%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 33%; height: 10px; background-color: grey;"></div> </div>
1	G	331	<div style="display: flex; align-items: center;"> <div style="width: 6%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 58%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 32%; height: 10px; background-color: grey;"></div> </div>
2	B	677	<div style="display: flex; align-items: center;"> <div style="width: 0%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 76%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 12%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 12%; height: 10px; background-color: grey;"></div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	D	677	 2% 76% 11% 12%
2	F	677	 75% 13% 11%
2	H	677	 2% 76% 11% 12%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 27654 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin carboxyl-terminal hydrolase 12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	239	Total 1886	C 1217	N 310	O 345	S 14	0	0	0
1	C	236	Total 1882	C 1215	N 308	O 345	S 14	0	0	0
1	E	222	Total 1747	C 1131	N 285	O 317	S 14	0	0	0
1	G	224	Total 1768	C 1136	N 290	O 328	S 14	0	1	0

- Molecule 2 is a protein called WD repeat-containing protein 48.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	595	Total 4690	C 2973	N 818	O 872	S 27	0	0	0
2	D	597	Total 4701	C 2976	N 822	O 876	S 27	0	0	0
2	F	600	Total 4731	C 2994	N 825	O 885	S 27	0	0	0
2	H	594	Total 4664	C 2948	N 819	O 870	S 27	0	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Zn 1	0	0
3	C	1	Total 1	Zn 1	0	0
3	E	1	Total 1	Zn 1	0	0
3	G	1	Total 1	Zn 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	81	Total O 81 81	0	0
4	B	286	Total O 286 286	0	0
4	C	82	Total O 82 82	0	0
4	D	334	Total O 334 334	0	0
4	E	68	Total O 68 68	0	0
4	F	364	Total O 364 364	0	0
4	G	57	Total O 57 57	0	0
4	H	309	Total O 309 309	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 12




- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 12




- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 12








- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 12




- Molecule 2: WD repeat-containing protein 48




- Molecule 2: WD repeat-containing protein 48




• Molecule 2: WD repeat-containing protein 48



• Molecule 2: WD repeat-containing protein 48







## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	262.43Å 103.30Å 178.47Å 90.00° 117.47° 90.00°	Depositor
Resolution (Å)	49.10 – 2.30 49.10 – 2.29	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.10-2.30) 92.4 (49.10-2.29)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.63 (at 2.29Å)	Xtrriage
Refinement program	PHENIX dev_1760	Depositor
R, $R_{free}$	0.188 , 0.239 0.191 , 0.241	Depositor DCC
$R_{free}$ test set	9518 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.7	Xtrriage
Anisotropy	0.153	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 43.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	27654	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.41	0/1923	0.56	0/2601
1	C	0.42	0/1918	0.66	4/2591 (0.2%)
1	E	0.40	0/1782	0.59	0/2408
1	G	0.44	1/1803 (0.1%)	0.56	0/2435
2	B	0.50	2/4783 (0.0%)	0.63	4/6485 (0.1%)
2	D	0.49	1/4795 (0.0%)	0.64	4/6502 (0.1%)
2	F	0.51	1/4825 (0.0%)	0.64	3/6543 (0.0%)
2	H	0.50	1/4757 (0.0%)	0.63	4/6452 (0.1%)
All	All	0.48	6/26586 (0.0%)	0.63	19/36017 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	1
All	All	0	2

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	189	CYS	CB-SG	-8.79	1.67	1.82
2	D	84	CYS	CB-SG	-6.10	1.71	1.82
2	F	84	CYS	CB-SG	-6.05	1.72	1.82
2	H	84	CYS	CB-SG	-5.82	1.72	1.81
2	B	84	CYS	CB-SG	-5.16	1.73	1.81

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	340	ASP	O-C-N	-13.95	100.37	122.70
2	F	73	HIS	N-CA-CB	10.34	129.21	110.60
2	B	73	HIS	N-CA-C	7.95	132.47	111.00
2	B	72	GLU	N-CA-C	7.00	129.90	111.00
1	C	340	ASP	CA-C-N	6.96	132.51	117.20

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	342	GLN	Peptide
2	B	599	HIS	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1886	0	1796	29	0
1	C	1882	0	1825	34	0
1	E	1747	0	1656	27	1
1	G	1768	0	1659	17	0
2	B	4690	0	4693	56	0
2	D	4701	0	4698	57	1
2	F	4731	0	4732	53	0
2	H	4664	0	4649	54	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0
3	E	1	0	0	0	0
3	G	1	0	0	0	0
4	A	81	0	0	0	0
4	B	286	0	0	9	0
4	C	82	0	0	4	0
4	D	334	0	0	11	0
4	E	68	0	0	7	0
4	F	364	0	0	7	0
4	G	57	0	0	1	0
4	H	309	0	0	8	0
All	All	27654	0	25708	319	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 319 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:284:LEU:HD13	1:C:366:TYR:CE1	1.81	1.16
1:A:314:ASN:HB3	1:A:319:ILE:CD1	1.84	1.07
1:A:314:ASN:HB3	1:A:319:ILE:HD11	1.45	0.95
1:C:284:LEU:CD1	1:C:366:TYR:CE1	2.49	0.95
2:F:30:ARG:NH1	2:F:361:ALA:O	2.05	0.89

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:249:ARG:NH2	1:E:338:LYS:O[2_555]	2.11	0.09

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	221/331 (67%)	211 (96%)	10 (4%)	0	100	100
1	C	218/331 (66%)	212 (97%)	6 (3%)	0	100	100
1	E	204/331 (62%)	197 (97%)	7 (3%)	0	100	100
1	G	205/331 (62%)	198 (97%)	6 (3%)	1 (0%)	29	35
2	B	583/677 (86%)	558 (96%)	23 (4%)	2 (0%)	41	50
2	D	587/677 (87%)	570 (97%)	17 (3%)	0	100	100
2	F	590/677 (87%)	570 (97%)	19 (3%)	1 (0%)	47	58
2	H	584/677 (86%)	563 (96%)	20 (3%)	1 (0%)	47	58
All	All	3192/4032 (79%)	3079 (96%)	108 (3%)	5 (0%)	47	58

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	334	ASP
2	B	73	HIS
2	F	73	HIS
2	B	307	ASP
2	H	307	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/305 (66%)	202 (100%)	0	100	100
1	C	207/305 (68%)	207 (100%)	0	100	100
1	E	186/305 (61%)	185 (100%)	1 (0%)	88	95
1	G	188/305 (62%)	185 (98%)	3 (2%)	62	78
2	B	518/597 (87%)	517 (100%)	1 (0%)	93	97
2	D	519/597 (87%)	514 (99%)	5 (1%)	76	87
2	F	525/597 (88%)	522 (99%)	3 (1%)	86	94
2	H	514/597 (86%)	508 (99%)	6 (1%)	71	84
All	All	2859/3608 (79%)	2840 (99%)	19 (1%)	84	92

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	84	CYS
2	H	307	ASP
2	H	527	ARG
2	H	282	ARG
2	F	169	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	146	GLN
2	D	283	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	239/331 (72%)	0.06	12 (5%) 28 35	25, 49, 80, 92	0
1	C	236/331 (71%)	0.17	10 (4%) 36 43	27, 53, 84, 95	0
1	E	222/331 (67%)	0.25	8 (3%) 42 49	25, 54, 84, 89	0
1	G	224/331 (67%)	0.49	21 (9%) 8 11	24, 55, 87, 97	0
2	B	595/677 (87%)	-0.15	7 (1%) 79 83	23, 35, 57, 90	0
2	D	597/677 (88%)	-0.15	11 (1%) 68 74	21, 33, 60, 81	0
2	F	600/677 (88%)	-0.26	3 (0%) 91 94	22, 31, 58, 73	0
2	H	594/677 (87%)	-0.11	16 (2%) 54 62	20, 32, 70, 87	0
All	All	3307/4032 (82%)	-0.05	88 (2%) 54 62	20, 37, 73, 97	0

The worst 5 of 88 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	44	PHE	5.4
2	H	500	VAL	5.1
1	G	358	SER	4.3
1	G	359	GLU	4.3
1	C	206	VAL	4.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	ZN	G	401	1/1	0.97	0.22	45,45,45,45	0
3	ZN	C	401	1/1	0.99	0.11	30,30,30,30	0
3	ZN	E	401	1/1	1.00	0.15	32,32,32,32	0
3	ZN	A	401	1/1	1.00	0.17	38,38,38,38	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.