



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 6, 2023 – 02:26 AM EDT

PDB ID : 1K4P
Title : Crystal Structure of 3,4-dihydroxy-2-butanone 4-phosphate synthase in complex with zinc ions
Authors : Liao, D.-I.; Zheng, Y.-J.; Viitanen, P.V.; Jordan, D.B.
Deposited on : 2001-10-08
Resolution : 1.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1117 (1.06-0.94)
Ramachandran outliers	138981	1043 (1.06-0.94)
Sidechain outliers	138945	1045 (1.06-0.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	233	 73% 17% 7%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1858 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3,4-Dihydroxy-2-Butanone 4-Phosphate Synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	216	1631	1009	303	308	11	0	0	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
3	A	5	5	5	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	212	Total 212	O 212	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: 3,4-Dihydroxy-2-Butanone 4-Phosphate Synthase



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	53.73Å 88.53Å 43.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 1.00	Depositor
% Data completeness (in resolution range)	(Not available) (15.00-1.00)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	TNT	Depositor
R, R_{free}	0.178 , 0.201	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1858	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.95	13/1661 (0.8%)	1.34	28/2255 (1.2%)

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	198	GLU	CD-OE2	5.97	1.32	1.25
1	A	101	GLU	CD-OE2	5.83	1.32	1.25
1	A	26	GLU	CD-OE2	5.82	1.32	1.25
1	A	155	GLU	CD-OE2	5.70	1.31	1.25
1	A	178	ASP	CG-OD2	5.62	1.38	1.25
1	A	174	GLU	CD-OE2	5.51	1.31	1.25
1	A	159	GLU	CD-OE2	5.37	1.31	1.25
1	A	72	GLU	CD-OE2	5.37	1.31	1.25
1	A	183	GLU	CD-OE2	5.37	1.31	1.25
1	A	181	GLU	CD-OE2	5.34	1.31	1.25
1	A	224	GLU	CD-OE2	5.29	1.31	1.25
1	A	37	GLU	CD-OE2	5.28	1.31	1.25
1	A	47	GLU	CD-OE1	-5.16	1.20	1.25

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	59	ARG	NE-CZ-NH1	8.96	124.78	120.30
1	A	148	ARG	NE-CZ-NH1	8.34	124.47	120.30
1	A	205	ARG	NE-CZ-NH1	7.94	124.27	120.30
1	A	197	ASP	CB-CG-OD2	-7.79	111.29	118.30
1	A	125	ASP	CB-CG-OD1	7.70	125.23	118.30
1	A	73	ARG	NE-CZ-NH1	7.02	123.81	120.30
1	A	177	ASP	CB-CG-OD1	6.83	124.45	118.30
1	A	59	ARG	NE-CZ-NH2	-6.79	116.91	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	178	ASP	N-CA-CB	6.68	122.62	110.60
1	A	33	ASP	CB-CG-OD1	6.67	124.30	118.30
1	A	148	ARG	NE-CZ-NH2	-6.60	117.00	120.30
1	A	78	ASP	CB-CG-OD1	6.56	124.21	118.30
1	A	215	ASP	CB-CG-OD1	6.45	124.11	118.30
1	A	197	ASP	CB-CG-OD1	6.34	124.01	118.30
1	A	33	ASP	CB-CG-OD2	-6.25	112.67	118.30
1	A	177	ASP	CB-CG-OD2	-6.21	112.71	118.30
1	A	113	ASP	CB-CG-OD1	6.16	123.85	118.30
1	A	151	ARG	NE-CZ-NH2	-6.15	117.22	120.30
1	A	36	ARG	NE-CZ-NH1	6.07	123.33	120.30
1	A	215	ASP	CB-CG-OD2	-5.91	112.98	118.30
1	A	78	ASP	CB-CG-OD2	-5.83	113.05	118.30
1	A	73	ARG	NE-CZ-NH2	-5.78	117.41	120.30
1	A	90	ARG	NE-CZ-NH1	5.75	123.17	120.30
1	A	88	ASP	CB-CG-OD2	-5.61	113.26	118.30
1	A	204	ARG	NE-CZ-NH2	-5.59	117.50	120.30
1	A	99	ASP	CB-CG-OD1	5.55	123.30	118.30
1	A	41	ASP	CB-CG-OD2	-5.30	113.53	118.30
1	A	125	ASP	CB-CG-OD2	-5.17	113.64	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1631	0	1618	18	1
2	A	10	0	0	0	0
3	A	5	0	0	0	0
4	A	212	0	0	8	1
All	All	1858	0	1618	18	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12:PHE:N	4:A:1147:HOH:O	2.16	0.78
1:A:20:GLN:OE1	4:A:1143:HOH:O	2.12	0.67
1:A:133:ARG:HH11	1:A:133:ARG:HG2	1.63	0.62
1:A:223:THR:HG22	1:A:224:GLU:OE2	2.01	0.59
1:A:83:VAL:HG21	1:A:86:ASN:HA	1.84	0.59
1:A:209:LYS:NZ	4:A:1056:HOH:O	2.37	0.56
1:A:151:ARG:HE	1:A:217:ILE:HD13	1.73	0.53
1:A:133:ARG:HG2	1:A:133:ARG:NH1	2.25	0.52
1:A:26:GLU:OE1	4:A:1206:HOH:O	2.20	0.48
1:A:133:ARG:HG2	4:A:1210:HOH:O	2.13	0.48
1:A:133:ARG:NH1	4:A:1210:HOH:O	2.29	0.46
1:A:133:ARG:HH11	1:A:133:ARG:CG	2.25	0.45
1:A:86:ASN:CB	4:A:1061:HOH:O	2.66	0.44
1:A:133:ARG:NH1	1:A:133:ARG:CG	2.82	0.42
1:A:223:THR:HG22	1:A:224:GLU:CD	2.40	0.42
1:A:227:LEU:HD12	1:A:227:LEU:HA	1.79	0.41
1:A:207:GLY:O	4:A:1206:HOH:O	2.22	0.41
1:A:127:GLN:HA	1:A:127:GLN:OE1	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:HIS:NE2	4:A:1143:HOH:O[4_455]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	214/233 (92%)	209 (98%)	4 (2%)	1 (0%)	29 8

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	196	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	171/186 (92%)	163 (95%)	8 (5%)	26 4

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	LEU
1	A	69	LEU
1	A	86	ASN
1	A	133	ARG
1	A	178	ASP
1	A	180	GLN
1	A	197	ASP
1	A	224	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	GLN
1	A	24	ASN
1	A	38	ASN
1	A	81	GLN
1	A	85	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 5 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	1002	-	4,4,4	0.61	0	6,6,6	1.28	1 (16%)
2	SO4	A	1001	3	4,4,4	0.72	0	6,6,6	0.84	0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1002	SO4	O4-S-O3	2.88	121.35	109.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.