



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2024 – 04:57 PM EST

PDB ID : 4K4I
Title : Ternary crystal structures of a human DNA POLYMERASE LAMBDA IN COMPLEX WITH DNA AND (-)FTC-TP.
Authors : Vyas, R.; Suo, Z.
Deposited on : 2013-04-12
Resolution : 2.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 78%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 17%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">5% 78% 17% . . .</p>
1	E	340	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 75%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 19%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">3% 75% 19% . .</p>
1	I	340	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 82%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 13%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">4% 82% 13% . .</p>
1	M	340	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 76%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 17%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">4% 76% 17% . .</p>
2	B	11	<div style="display: flex; align-items: center;"> <div style="width: 73%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 27%; height: 10px; background-color: yellow; margin-right: 5px;"></div> </div> <p style="text-align: center;">73% 27%</p>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	11	 73% 9% 18%
2	J	11	 91% 9%
2	N	11	 73% 27%
3	C	6	 100%
3	G	6	 100%
3	K	6	 83% 17%
3	O	6	 83% 17%
4	D	4	 25% 25% 50%
4	H	4	 75% 25%
4	L	4	 75% 25%
4	P	4	 75% 25%

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 12760 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase lambda.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	333	Total 2649	C 1663	N 489	O 485	S 12	0	3	0
1	E	329	Total 2589	C 1626	N 474	O 477	S 12	0	0	0
1	I	325	Total 2574	C 1617	N 473	O 473	S 11	0	0	0
1	M	329	Total 2592	C 1628	N 474	O 478	S 12	0	1	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	244	MET	-	expression tag	UNP Q9UGP5
A	576	LEU	-	expression tag	UNP Q9UGP5
A	577	GLU	-	expression tag	UNP Q9UGP5
A	578	HIS	-	expression tag	UNP Q9UGP5
A	579	HIS	-	expression tag	UNP Q9UGP5
A	580	HIS	-	expression tag	UNP Q9UGP5
A	581	HIS	-	expression tag	UNP Q9UGP5
A	582	HIS	-	expression tag	UNP Q9UGP5
A	583	HIS	-	expression tag	UNP Q9UGP5
E	244	MET	-	expression tag	UNP Q9UGP5
E	576	LEU	-	expression tag	UNP Q9UGP5
E	577	GLU	-	expression tag	UNP Q9UGP5
E	578	HIS	-	expression tag	UNP Q9UGP5
E	579	HIS	-	expression tag	UNP Q9UGP5
E	580	HIS	-	expression tag	UNP Q9UGP5
E	581	HIS	-	expression tag	UNP Q9UGP5
E	582	HIS	-	expression tag	UNP Q9UGP5
E	583	HIS	-	expression tag	UNP Q9UGP5
I	244	MET	-	expression tag	UNP Q9UGP5
I	576	LEU	-	expression tag	UNP Q9UGP5
I	577	GLU	-	expression tag	UNP Q9UGP5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
I	578	HIS	-	expression tag	UNP Q9UGP5
I	579	HIS	-	expression tag	UNP Q9UGP5
I	580	HIS	-	expression tag	UNP Q9UGP5
I	581	HIS	-	expression tag	UNP Q9UGP5
I	582	HIS	-	expression tag	UNP Q9UGP5
I	583	HIS	-	expression tag	UNP Q9UGP5
M	244	MET	-	expression tag	UNP Q9UGP5
M	576	LEU	-	expression tag	UNP Q9UGP5
M	577	GLU	-	expression tag	UNP Q9UGP5
M	578	HIS	-	expression tag	UNP Q9UGP5
M	579	HIS	-	expression tag	UNP Q9UGP5
M	580	HIS	-	expression tag	UNP Q9UGP5
M	581	HIS	-	expression tag	UNP Q9UGP5
M	582	HIS	-	expression tag	UNP Q9UGP5
M	583	HIS	-	expression tag	UNP Q9UGP5

- Molecule 2 is a DNA chain called DNA (5'-D(*CP*GP*GP*CP*GP*GP*TP*AP*CP*TP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	B	11	Total 225	C 107	N 43	O 65	P 10	0	0	0
2	F	11	Total 225	C 107	N 43	O 65	P 10	0	0	0
2	J	11	Total 225	C 107	N 43	O 65	P 10	0	0	0
2	N	11	Total 225	C 107	N 43	O 65	P 10	0	0	0


- Molecule 3 is a DNA chain called DNA (5'-D(*CP*AP*GP*TP*AP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	C	6	Total 119	C 58	N 23	O 33	P 5	0	0	0
3	G	6	Total 119	C 58	N 23	O 33	P 5	0	0	0
3	K	6	Total 119	C 58	N 23	O 33	P 5	0	0	0
3	O	6	Total 119	C 58	N 23	O 33	P 5	0	0	0

- Molecule 4 is a DNA chain called DNA (5'-D(P*GP*CP*CP*G)-3').

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
4	D	4	Total	C	N	O	P	0	0	0
			83	38	16	25	4			
4	H	4	Total	C	N	O	P	0	0	0
			83	38	16	25	4			
4	L	4	Total	C	N	O	P	0	0	0
			83	38	16	25	4			
4	P	4	Total	C	N	O	P	0	0	0
			83	38	16	25	4			

- Molecule 5 is [[(2R,5S)-5-(4-azanyl-5-fluoranyl-2-oxidanylidene-pyrimidin-1-yl)-1,3-oxathiolan-2-yl]methoxy-oxidanyl-phosphoryl] phosphono hydrogen phosphate (three-letter code: 1RY) (formula: C₈H₁₃FN₃O₁₂P₃S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	
5	A	1	Total	C	F	N	O	P	S	0	0
			28	8	1	3	12	3	1		
5	E	1	Total	C	F	N	O	P	S	0	0
			28	8	1	3	12	3	1		
5	I	1	Total	C	F	N	O	P	S	0	0
			28	8	1	3	12	3	1		
5	M	1	Total	C	F	N	O	P	S	0	0
			28	8	1	3	12	3	1		

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	6	Total	Ca	0	0
			6	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total Ca 1 1	0	0
6	E	5	Total Ca 5 5	0	0
6	F	1	Total Ca 1 1	0	0
6	I	3	Total Ca 3 3	0	0
6	J	1	Total Ca 1 1	0	0
6	K	1	Total Ca 1 1	0	0
6	M	4	Total Ca 4 4	0	0
6	N	1	Total Ca 1 1	0	0

- Molecule 7 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 4 2 2	0	0
7	B	1	Total C O 4 2 2	0	0
7	E	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	F	1	Total	C	O	0	0
			4	2	2		
7	I	1	Total	C	O	0	0
			4	2	2		
7	M	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is CACODYLATE ION (three-letter code: CAC) (formula: C₂H₆AsO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	E	1	Total	As	C	O	0	0
			5	1	2	2		
8	M	1	Total	As	C	O	0	0
			5	1	2	2		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	129	Total	O	0	0
			129	129		
9	B	20	Total	O	0	0
			20	20		
9	C	13	Total	O	0	0
			13	13		
9	E	78	Total	O	0	0
			78	78		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	F	9	Total O 9 9	0	0
9	G	2	Total O 2 2	0	0
9	H	2	Total O 2 2	0	0
9	I	76	Total O 76 76	0	0
9	J	4	Total O 4 4	0	0
9	K	9	Total O 9 9	0	0
9	M	124	Total O 124 124	0	0
9	N	8	Total O 8 8	0	0
9	O	3	Total O 3 3	0	0
9	P	2	Total O 2 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase lambda



- Molecule 1: DNA polymerase lambda






- Molecule 1: DNA polymerase lambda






- Molecule 1: DNA polymerase lambda




- Molecule 2: DNA (5'-D(*CP*GP*GP*CP*GP*GP*TP*AP*CP*TP*G)-3')



- Molecule 2: DNA (5'-D(*CP*GP*GP*CP*GP*GP*TP*AP*CP*TP*G)-3')



- Molecule 2: DNA (5'-D(*CP*GP*GP*CP*GP*GP*TP*AP*CP*TP*G)-3')



- Molecule 2: DNA (5'-D(*CP*GP*GP*CP*GP*GP*TP*AP*CP*TP*G)-3')



- Molecule 3: DNA (5'-D(*CP*AP*GP*TP*AP*C)-3')




There are no outlier residues recorded for this chain.

- Molecule 3: DNA (5'-D(*CP*AP*GP*TP*AP*C)-3')

Chain G:  100%


There are no outlier residues recorded for this chain.

- Molecule 3: DNA (5'-D(*CP*AP*GP*TP*AP*C)-3')

Chain K:  83% 17%




- Molecule 3: DNA (5'-D(*CP*AP*GP*TP*AP*C)-3')

Chain O:  83% 17%




- Molecule 4: DNA (5'-D(P*GP*CP*CP*G)-3')

Chain D:  25% 25% 50%



- Molecule 4: DNA (5'-D(P*GP*CP*CP*G)-3')

Chain H:  75% 25%



- Molecule 4: DNA (5'-D(P*GP*CP*CP*G)-3')

Chain L:  75% 25%



- Molecule 4: DNA (5'-D(P*GP*CP*CP*G)-3')

Chain P:  75% 25%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	194.44Å 97.58Å 105.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.03 – 2.25 45.03 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.9 (45.03-2.25) 100.0 (45.03-2.25)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.97 (at 2.24Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.212 , 0.274 0.216 , 0.274	Depositor DCC
R_{free} test set	4788 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	32.5	Xtrriage
Anisotropy	0.202	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 40.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	12760	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ACT, 1RY, CAC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.85	1/2716 (0.0%)	0.94	4/3666 (0.1%)
1	E	0.76	0/2642	0.88	5/3566 (0.1%)
1	I	0.74	1/2628 (0.0%)	0.89	6/3546 (0.2%)
1	M	0.85	2/2648 (0.1%)	1.04	11/3574 (0.3%)
2	B	0.68	0/252	1.09	1/388 (0.3%)
2	F	0.51	0/252	1.01	3/388 (0.8%)
2	J	0.64	0/252	0.89	0/388
2	N	0.65	0/252	0.96	0/388
3	C	0.86	0/133	1.04	0/203
3	G	0.60	0/133	0.93	0/203
3	K	0.60	0/133	1.08	2/203 (1.0%)
3	O	0.73	0/133	0.98	1/203 (0.5%)
4	D	1.17	1/92 (1.1%)	1.11	2/138 (1.4%)
4	H	1.33	1/92 (1.1%)	0.88	0/138
4	L	1.08	1/92 (1.1%)	0.81	0/138
4	P	1.23	1/92 (1.1%)	0.84	0/138
All	All	0.80	8/12542 (0.1%)	0.95	35/17268 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	M	0	1

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	1	DG	OP3-P	-10.91	1.48	1.61
4	P	1	DG	OP3-P	-10.16	1.49	1.61

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	1	DG	OP3-P	-9.65	1.49	1.61
4	L	1	DG	OP3-P	-8.95	1.50	1.61
1	M	429	ASP	CB-CG	6.77	1.66	1.51

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	M	488	ARG	NE-CZ-NH2	-12.84	113.88	120.30
1	M	427	ASP	CB-CG-OD1	10.82	128.04	118.30
1	M	488	ARG	NE-CZ-NH1	10.42	125.51	120.30
1	A	427	ASP	CB-CG-OD1	9.96	127.27	118.30
1	M	441	ARG	NE-CZ-NH2	-9.63	115.48	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	M	290	HIS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2649	0	2639	54	0
1	E	2589	0	2590	58	0
1	I	2574	0	2566	23	0
1	M	2592	0	2595	52	0
2	B	225	0	125	1	0
2	F	225	0	125	4	0
2	J	225	0	125	1	0
2	N	225	0	125	2	0
3	C	119	0	68	0	0
3	G	119	0	69	0	0
3	K	119	0	69	0	0
3	O	119	0	69	0	0
4	D	83	0	45	2	0
4	H	83	0	45	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	L	83	0	45	0	0
4	P	83	0	45	0	0
5	A	28	0	11	0	0
5	E	28	0	11	0	0
5	I	28	0	11	0	0
5	M	28	0	12	1	0
6	A	6	0	0	0	0
6	B	1	0	0	0	0
6	E	5	0	0	0	0
6	F	1	0	0	0	0
6	I	3	0	0	0	0
6	J	1	0	0	0	0
6	K	1	0	0	0	0
6	M	4	0	0	0	0
6	N	1	0	0	0	0
7	A	4	0	3	0	0
7	B	4	0	3	0	0
7	E	4	0	3	1	0
7	F	4	0	3	0	0
7	I	4	0	3	0	0
7	M	4	0	3	0	0
8	E	5	0	0	0	0
8	M	5	0	0	3	0
9	A	129	0	0	12	0
9	B	20	0	0	0	0
9	C	13	0	0	0	0
9	E	78	0	0	7	0
9	F	9	0	0	0	0
9	G	2	0	0	0	0
9	H	2	0	0	0	0
9	I	76	0	0	1	0
9	J	4	0	0	1	0
9	K	9	0	0	0	0
9	M	124	0	0	3	0
9	N	8	0	0	1	0
9	O	3	0	0	0	0
9	P	2	0	0	0	0
All	All	12760	0	11408	192	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 192 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:397:GLN:HG3	9:E:770:HOH:O	1.58	1.01
1:M:535:ALA:HB1	1:M:536:VAL:HG23	1.38	1.01
1:M:536:VAL:H	1:M:537:VAL:HA	1.23	1.01
1:E:521:LYS:NZ	2:F:4:DC:OP1	2.03	0.90
1:E:501:CYS:SG	1:E:531:ALA:HA	2.15	0.86

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	334/340 (98%)	313 (94%)	17 (5%)	4 (1%)	13	9
1	E	327/340 (96%)	296 (90%)	25 (8%)	6 (2%)	8	4
1	I	321/340 (94%)	308 (96%)	12 (4%)	1 (0%)	41	46
1	M	328/340 (96%)	306 (93%)	15 (5%)	7 (2%)	7	3
All	All	1310/1360 (96%)	1223 (93%)	69 (5%)	18 (1%)	11	7

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	538	ARG
1	M	251	ASN
1	M	289	PHE
1	M	290	HIS
1	M	536	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/287 (99%)	270 (95%)	14 (5%)	25	27
1	E	276/287 (96%)	260 (94%)	16 (6%)	20	20
1	I	275/287 (96%)	264 (96%)	11 (4%)	31	37
1	M	277/287 (96%)	256 (92%)	21 (8%)	13	12
All	All	1112/1148 (97%)	1050 (94%)	62 (6%)	21	21

5 of 62 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	561	ARG
1	M	493	VAL
1	I	445	SER
1	M	489	LEU
1	M	544	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	354	GLN
1	I	470	GLN
1	I	397	GLN
1	M	255	HIS
1	A	541	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 35 ligands modelled in this entry, 23 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	1RY	E	601	6	25,29,29	2.19	6 (24%)	35,45,45	1.42	5 (14%)
7	ACT	A	608	-	3,3,3	0.73	0	3,3,3	1.98	1 (33%)
5	1RY	I	601	6	25,29,29	1.99	4 (16%)	35,45,45	2.24	8 (22%)
5	1RY	A	601	6	25,29,29	1.98	7 (28%)	35,45,45	1.67	8 (22%)
8	CAC	M	606	-	0,4,4	-	-	0,6,6	-	-
7	ACT	E	608	-	3,3,3	0.64	0	3,3,3	1.03	0
8	CAC	E	607	-	0,4,4	-	-	0,6,6	-	-
7	ACT	F	102	-	3,3,3	0.94	0	3,3,3	0.71	0
5	1RY	M	601	6	25,29,29	1.70	6 (24%)	35,45,45	2.38	11 (31%)
7	ACT	I	605	-	3,3,3	0.76	0	3,3,3	1.30	0
7	ACT	M	607	-	3,3,3	0.93	0	3,3,3	0.65	0
7	ACT	B	102	-	3,3,3	0.70	0	3,3,3	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	1RY	M	601	6	-	7/22/31/31	0/2/2/2
5	1RY	I	601	6	-	5/22/31/31	0/2/2/2
5	1RY	A	601	6	-	3/22/31/31	0/2/2/2
5	1RY	E	601	6	-	4/22/31/31	0/2/2/2

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	I	601	1RY	OAB-CAV	7.67	1.37	1.23
5	E	601	1RY	OAB-CAV	6.49	1.35	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	E	601	1RY	CAT-CAU	-6.31	1.37	1.42
5	A	601	1RY	CAT-CAU	-5.15	1.38	1.42
5	M	601	1RY	OAB-CAV	4.67	1.32	1.23

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	I	601	1RY	FAJ-CAU-CAT	9.72	124.37	118.02
5	M	601	1RY	OAP-CAX-NAY	6.90	120.19	107.86
5	M	601	1RY	PBB-OAQ-PAZ	-6.83	109.38	132.83
5	A	601	1RY	OAB-CAV-NAN	-5.05	114.12	122.33
5	M	601	1RY	OAB-CAV-NAN	-4.67	114.74	122.33

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	601	1RY	PBB-OAQ-PAZ-OAF
5	E	601	1RY	PBB-OAQ-PAZ-OAF
5	I	601	1RY	PBB-OAQ-PAZ-OAC
5	I	601	1RY	PBB-OAQ-PAZ-OAF
5	M	601	1RY	OAO-CAL-CAW-OAP


There are no ring outliers.


3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	M	606	CAC	3	0
7	E	608	ACT	1	0
5	M	601	1RY	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	333/340 (97%)	-0.07	17 (5%) 28 30	17, 32, 69, 132	0
1	E	329/340 (96%)	-0.08	11 (3%) 46 48	20, 37, 75, 127	0
1	I	325/340 (95%)	-0.17	12 (3%) 41 44	19, 37, 63, 95	0
1	M	329/340 (96%)	-0.13	13 (3%) 38 40	15, 33, 56, 112	0
2	B	11/11 (100%)	-0.54	0 100 100	21, 25, 42, 46	0
2	F	11/11 (100%)	-0.45	0 100 100	36, 43, 55, 60	0
2	J	11/11 (100%)	-0.49	0 100 100	33, 35, 42, 43	0
2	N	11/11 (100%)	-0.56	0 100 100	22, 29, 40, 44	0
3	C	6/6 (100%)	-0.42	0 100 100	20, 24, 37, 39	0
3	G	6/6 (100%)	-0.18	0 100 100	28, 32, 57, 68	0
3	K	6/6 (100%)	-0.33	0 100 100	23, 25, 64, 69	0
3	O	6/6 (100%)	-0.67	0 100 100	24, 28, 47, 50	0
4	D	4/4 (100%)	-0.87	0 100 100	33, 35, 37, 39	0
4	H	4/4 (100%)	-0.69	0 100 100	28, 32, 33, 37	0
4	L	4/4 (100%)	-0.56	0 100 100	36, 38, 44, 46	0
4	P	4/4 (100%)	-0.88	0 100 100	32, 36, 41, 44	0
All	All	1400/1444 (96%)	-0.14	53 (3%) 40 43	15, 35, 66, 132	0

The worst 5 of 53 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	538	ARG	8.0
1	A	540	THR	7.5
1	M	249	ALA	6.1
1	M	543	CYS	5.8
1	M	542	GLY	5.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	CAC	E	607	5/5	0.53	0.32	132,134,146,155	0
7	ACT	M	607	4/4	0.60	0.22	55,55,57,61	0
6	CA	M	604	1/1	0.60	0.10	49,49,49,49	0
6	CA	J	101	1/1	0.72	0.07	58,58,58,58	0
6	CA	A	607	1/1	0.83	0.16	59,59,59,59	0
6	CA	M	605	1/1	0.84	0.11	61,61,61,61	0
6	CA	I	604	1/1	0.85	0.11	64,64,64,64	0
6	CA	K	101	1/1	0.85	0.14	64,64,64,64	0
7	ACT	E	608	4/4	0.86	0.18	54,55,57,58	0
6	CA	M	602	1/1	0.87	0.15	65,65,65,65	0
6	CA	E	603	1/1	0.89	0.06	50,50,50,50	0
6	CA	E	604	1/1	0.89	0.08	64,64,64,64	0
7	ACT	B	102	4/4	0.89	0.17	48,49,49,57	0
6	CA	E	606	1/1	0.91	0.11	62,62,62,62	0
6	CA	A	602	1/1	0.92	0.17	33,33,33,33	0
7	ACT	F	102	4/4	0.92	0.17	43,48,48,50	0
7	ACT	I	605	4/4	0.92	0.24	39,43,45,49	0
6	CA	N	101	1/1	0.92	0.07	58,58,58,58	0
6	CA	E	605	1/1	0.92	0.18	39,39,39,39	0
8	CAC	M	606	5/5	0.92	0.12	74,77,82,88	0
6	CA	F	101	1/1	0.93	0.05	47,47,47,47	0
6	CA	A	606	1/1	0.93	0.06	29,29,29,29	0
7	ACT	A	608	4/4	0.93	0.19	41,41,45,51	0
6	CA	M	603	1/1	0.93	0.10	42,42,42,42	0
6	CA	A	603	1/1	0.93	0.15	53,53,53,53	0
5	1RY	I	601	28/28	0.94	0.15	41,52,65,66	0
6	CA	I	603	1/1	0.94	0.08	38,38,38,38	0


Continued on next page...


Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	1RY	M	601	28/28	0.95	0.11	23,32,43,47	0
6	CA	B	101	1/1	0.95	0.16	66,66,66,66	0
6	CA	A	604	1/1	0.97	0.03	37,37,37,37	0
5	1RY	A	601	28/28	0.98	0.11	18,20,23,24	0
6	CA	A	605	1/1	0.98	0.08	21,21,21,21	0
5	1RY	E	601	28/28	0.98	0.10	23,32,36,40	0
6	CA	I	602	1/1	0.99	0.14	35,35,35,35	0
6	CA	E	602	1/1	0.99	0.10	30,30,30,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers [i](#)

There are no such residues in this entry.