wwPDB X-ray Structure Validation Summary Report (i)

Sep 10, 2023 – 06:07 PM EDT

PDB ID	:	4K5Y
Title	:	Crystal structure of human corticotropin-releasing factor receptor 1 (CRF1R) in complex with the antagonist CP-376395
Authors	:	Hollenstein, K.; Kean, J.; Bortolato, A.; Cheng, R.K.Y.; Dore, A.S.; Jazayeri, A.; Cooke, R.M.; Weir, M.; Marshall, F.H.
Deposited on	:	2013-04-15
Resolution	:	2.98 Å(reported)

The following versions of software and data (see references (i)) were used in the production of this report:

|		
MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2754 (3.00-2.96)
Clashscore	141614	3103 (3.00-2.96)
Ramachandran outliers	138981	2993 (3.00-2.96)
Sidechain outliers	138945	2996 (3.00-2.96)
RSRZ outliers	127900	2644 (3.00-2.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	441	7%; 79% 13% 8%
1	B	441	3%; 78% 12% 10%
1	C	441	6%; 50% 6% 44%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	SO4	A	504	-	-	X	-

2 Entry composition ⓘ

There are 8 unique types of molecules in this entry. The entry contains 8823 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Corticotropin-releasing factor receptor 1, T4-Lysozyme chimeric construct.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	407	Total 3278	C 2145	N 562	O 554	S 17	0	0	0
1	B	396	Total 3194	C 2091	N 547	O 539	S 17	0	0	0
1	C	248	Total 2005	C 1341	N 330	O 321	S 13	0	0	0

There are 90 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	103	MET	-	initiating methionine	UNP P34998
A	120	ALA	VAL	engineered mutation	UNP P34998
A	144	ALA	LEU	engineered mutation	UNP P34998
A	156	ALA	TRP	engineered mutation	UNP P34998
A	160	ALA	SER	engineered mutation	UNP P34998
A	1040	SER	ASN	engineered mutation	UNP P00720
A	1041	VAL	ALA	engineered mutation	UNP P00720
A	1054	SER	CYS	engineered mutation	UNP P00720
A	1097	SER	CYS	engineered mutation	UNP P00720
A	1151	ALA	THR	engineered mutation	UNP P00720
A	228	ALA	LYS	engineered mutation	UNP P34998
A	260	ALA	PHE	engineered mutation	UNP P34998
A	277	ALA	ILE	engineered mutation	UNP P34998
A	309	ALA	TYR	engineered mutation	UNP P34998
A	330	ALA	PHE	engineered mutation	UNP P34998
A	349	ALA	SER	engineered mutation	UNP P34998
A	363	ALA	TYR	engineered mutation	UNP P34998
A	374	ALA	-	expression tag	UNP P34998
A	375	ALA	-	expression tag	UNP P34998
A	376	ALA	-	expression tag	UNP P34998
A	377	HIS	-	expression tag	UNP P34998
A	378	HIS	-	expression tag	UNP P34998

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	379	HIS	-	expression tag	UNP P34998
A	380	HIS	-	expression tag	UNP P34998
A	381	HIS	-	expression tag	UNP P34998
A	382	HIS	-	expression tag	UNP P34998
A	383	HIS	-	expression tag	UNP P34998
A	384	HIS	-	expression tag	UNP P34998
A	385	HIS	-	expression tag	UNP P34998
A	386	HIS	-	expression tag	UNP P34998
B	103	MET	-	initiating methionine	UNP P34998
B	120	ALA	VAL	engineered mutation	UNP P34998
B	144	ALA	LEU	engineered mutation	UNP P34998
B	156	ALA	TRP	engineered mutation	UNP P34998
B	160	ALA	SER	engineered mutation	UNP P34998
B	1040	SER	ASN	engineered mutation	UNP P00720
B	1041	VAL	ALA	engineered mutation	UNP P00720
B	1054	SER	CYS	engineered mutation	UNP P00720
B	1097	SER	CYS	engineered mutation	UNP P00720
B	1151	ALA	THR	engineered mutation	UNP P00720
B	228	ALA	LYS	engineered mutation	UNP P34998
B	260	ALA	PHE	engineered mutation	UNP P34998
B	277	ALA	ILE	engineered mutation	UNP P34998
B	309	ALA	TYR	engineered mutation	UNP P34998
B	330	ALA	PHE	engineered mutation	UNP P34998
B	349	ALA	SER	engineered mutation	UNP P34998
B	363	ALA	TYR	engineered mutation	UNP P34998
B	374	ALA	-	expression tag	UNP P34998
B	375	ALA	-	expression tag	UNP P34998
B	376	ALA	-	expression tag	UNP P34998
B	377	HIS	-	expression tag	UNP P34998
B	388	HIS	-	expression tag	UNP P34998
B	389	HIS	-	expression tag	UNP P34998
B	390	HIS	-	expression tag	UNP P34998
B	391	HIS	-	expression tag	UNP P34998
B	392	HIS	-	expression tag	UNP P34998
B	393	HIS	-	expression tag	UNP P34998
B	394	HIS	-	expression tag	UNP P34998
B	395	HIS	-	expression tag	UNP P34998
B	396	HIS	-	expression tag	UNP P34998
C	103	MET	-	initiating methionine	UNP P34998
C	120	ALA	VAL	engineered mutation	UNP P34998
C	144	ALA	LEU	engineered mutation	UNP P34998
C	156	ALA	TRP	engineered mutation	UNP P34998

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	160	ALA	SER	engineered mutation	UNP P34998
C	1040	SER	ASN	engineered mutation	UNP P00720
C	1041	VAL	ALA	engineered mutation	UNP P00720
C	1054	SER	CYS	engineered mutation	UNP P00720
C	1097	SER	CYS	engineered mutation	UNP P00720
C	1151	ALA	THR	engineered mutation	UNP P00720
C	228	ALA	LYS	engineered mutation	UNP P34998
C	260	ALA	PHE	engineered mutation	UNP P34998
C	277	ALA	ILE	engineered mutation	UNP P34998
C	309	ALA	TYR	engineered mutation	UNP P34998
C	330	ALA	PHE	engineered mutation	UNP P34998
C	349	ALA	SER	engineered mutation	UNP P34998
C	363	ALA	TYR	engineered mutation	UNP P34998
C	374	ALA	-	expression tag	UNP P34998
C	375	ALA	-	expression tag	UNP P34998
C	376	ALA	-	expression tag	UNP P34998
C	377	HIS	-	expression tag	UNP P34998
C	378	HIS	-	expression tag	UNP P34998
C	379	HIS	-	expression tag	UNP P34998
C	380	HIS	-	expression tag	UNP P34998
C	381	HIS	-	expression tag	UNP P34998
C	382	HIS	-	expression tag	UNP P34998
C	383	HIS	-	expression tag	UNP P34998
C	384	HIS	-	expression tag	UNP P34998
C	385	HIS	-	expression tag	UNP P34998
C	386	HIS	-	expression tag	UNP P34998

- Molecule 2 is 3,6-dimethyl-N-(pentan-3-yl)-2-(2,4,6-trimethylphenoxy)pyridin-4-amine (three-letter code: 1Q5) (formula: $C_{21}H_{30}N_2O$).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 24	C 21	N 2	O 1	0	0
2	B	1	Total 24	C 21	N 2	O 1	0	0
2	C	1	Total 24	C 21	N 2	O 1	0	0

- Molecule 3 is OLEIC ACID (three-letter code: OLA) (formula: $C_{18}H_{34}O_2$).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 13	C 13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total 20	C 18	O 2	0	0

- Molecule 4 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula: $C_{21}H_{40}O_4$).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total 25	C 21	O 4	0	0
4	A	1	Total 25	C 21	O 4	0	0
4	B	1	Total 25	C 21	O 4	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O_4S).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total 5	O 4	S 1	0	0
5	B	1	Total 5	O 4	S 1	0	0
5	B	1	Total 5	O 4	S 1	0	0
5	B	1	Total 5	O 4	S 1	0	0

- Molecule 6 is (1R)-2-{[(S)-{[(2S)-2,3-dihydroxypropyl]oxy}(hydroxy)phosphoryl]oxy}-1-[(hexadecanoyloxy)methyl]ethyl (9Z)-octadec-9-enoate (three-letter code: PGW) (formula: $C_{40}H_{77}O_{10}P$).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total 39	C 30	O 8	P 1	0	0
6	B	1	Total 46	C 37	O 8	P 1	0	0
6	B	1	Total 37	C 32	O 5		0	0

- Molecule 7 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: $C_{10}H_{22}O_6$).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total 16	C 10	O 6	0	0

- Molecule 8 is water.

| Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	4	Total 4	O 4	0	0
8	B	4	Total 4	O 4	0	0

3 Residue-property plots ⓘ

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Corticotropin-releasing factor receptor 1, T4-Lysozyme chimeric construct

• Molecule 1: Corticotropin-releasing factor receptor 1, T4-Lysozyme chimeric construct

• Molecule 1: Corticotropin-releasing factor receptor 1, T4-Lysozyme chimeric construct

GLY LEU ARG LEU LYS ILE TYR LYS ASP THR GLU GLY TYR TYR THR ILE GLY ILE GLY HIS LEU LEU THR LYS SER PRO SER LEU ASN SER LEU VAL ALA LYS SER GLU LEU ASP LYS ALA ILE GLY ARG ASN SER ASN GLY VAL ILE THR LYS ASP GLU ALA GLU LYS LEU PHE ASN GLN ASP VAL

ASP ALA ALA VAL ARG GLY ILE LEU ARG ASN ALA LYS LEU LYS PRO VAL TYR ASP SER LEU ASP ALA VAL ARG ARG SER ALA LEU ILE ASN ASN MET VAL PHE GLN MET GLU GLU THR GLY VAL ALA GLY PHE THR ASN SER LEU ARG MET LEU GLN GLN LYS ARG TRP ASP GLU ALA ALA VAL

ASN LEU ALA LYS SER ARG TRP TYR ASN GLN THR PRO ASN ARG ALA LYS ARG VAL ILE ALA THR PHE ARG THR GLY THR TRP ASP ALA TYR **D224** **R225** W236 I243 **Y253** **R263** **P264** **G265** **V266** **Y267** **Y270** V279 I293 **L294** **M295** **T296** R299 **A300** **S301** **T302** L323

L329 **N333** **P334** **G335** **E336** **D337** E338 **V339** **S340** F344 F347 L351 G356 **N367** SER GLU VAL ARG SER ALA ALA ALA HIS HIS HIS HIS HIS HIS HIS HIS HIS HIS

4 Data and refinement statistics ⓘ

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α, β, γ	86.56Å 123.97Å 166.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.14 – 2.98 34.15 – 2.98	Depositor EDS
% Data completeness (in resolution range)	85.7 (34.14-2.98) 85.7 (34.15-2.98)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$<I/\sigma(I)>$ [1]	1.73 (at 3.00Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.242 , 0.265 0.245 , 0.269	Depositor DCC
R_{free} test set	1586 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å^2)	55.4	Xtriage
Anisotropy	0.324	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.31 , 50.2	EDS
L-test for twinning[2]	$<\lvert L\rvert> = 0.45$, $<L^2> = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.89	EDS
Total number of atoms	8823	wwPDB-VP
Average B, all atoms (Å^2)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

---

[1]Intensities estimated from amplitudes.
[2]Theoretical values of $<\lvert L\rvert>$, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 1PE, PGW, OLA, 1Q5, OLC, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#|Z| >5	RMSZ	#|Z| >5
1	A	0.62	0/3358	0.68	3/4556 (0.1%)
1	B	0.65	0/3271	0.65	1/4435 (0.0%)
1	C	0.60	0/2063	0.69	0/2810
All	All	0.63	0/8692	0.67	4/11801 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	262	LYS	N-CA-C	-7.98	89.46	111.00
1	B	296	THR	N-CA-C	6.72	129.15	111.00
1	A	1048	LYS	CD-CE-NZ	-6.53	96.68	111.70
1	A	1007	LEU	CB-CG-CD2	-5.55	101.56	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	256	GLU	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3278	0	3333	40	1
1	B	3194	0	3247	40	1
1	C	2005	0	2038	19	1
2	A	24	0	30	2	0
2	B	24	0	30	5	0
2	C	24	0	30	3	0
3	A	13	0	23	1	0
3	C	20	0	33	1	0
4	A	50	0	80	4	0
4	B	25	0	40	2	0
5	A	5	0	0	1	1
5	B	15	0	0	0	1
6	B	122	0	174	10	0
7	B	16	0	22	0	0
8	A	4	0	0	0	0
8	B	4	0	0	1	0
All	All	8823	0	9080	102	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 102 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:338:GLU:HB2	1:C:340:SER:H	1.44	0.82
1:B:329:LEU:HB3	1:B:344:PHE:HE1	1.48	0.77
1:A:1006:MET:HG2	1:A:1007:LEU:HD12	1.69	0.75
1:A:263:ARG:O	1:A:265:GLY:HA3	1.87	0.74
1:C:263:ARG:O	1:C:265:GLY:HA3	1.92	0.70

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1137:ARG:NH1	5:B:506:SO4:O1[1_655]	2.16	0.04
1:B:1137:ARG:NH1	5:A:504:SO4:O2[1_455]	2.17	0.03
1:C:270:TYR:OH	1:C:338:GLU:OE1[2_556]	2.19	0.01

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	403/441 (91%)	394 (98%)	9 (2%)	0	100	100
1	B	390/441 (88%)	383 (98%)	7 (2%)	0	100	100
1	C	244/441 (55%)	240 (98%)	4 (2%)	0	100	100
All	All	1037/1323 (78%)	1017 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	345/374 (92%)	344 (100%)	1 (0%)	92	97
1	B	335/374 (90%)	333 (99%)	2 (1%)	86	94
1	C	211/374 (56%)	210 (100%)	1 (0%)	88	95
All	All	891/1122 (79%)	887 (100%)	4 (0%)	91	97

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	253	TYR
1	B	283	ASN
1	B	354	PHE
1	C	333	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	152	ASN
1	A	367	ASN
1	C	333	ASN

5.3.3   RNA ⓘ

There are no RNA molecules in this entry.

5.4   Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5   Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6   Ligand geometry ⓘ

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	#$|Z| > 2$	Counts	RMSZ	#$|Z| > 2$
4	OLC	A	502	-	24,24,24	0.80	2 (8%)	25,25,25	0.99	1 (4%)
6	PGW	B	503	-	45,45,50	0.89	3 (6%)	49,50,56	1.18	3 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	#|Z| > 2	Counts	RMSZ	#|Z| > 2
2	1Q5	A	401	-	24,25,25	1.59	3 (12%)	31,35,35	1.96	9 (29%)
6	PGW	B	504	-	36,36,50	0.92	3 (8%)	38,38,56	1.23	2 (5%)
2	1Q5	B	401	-	24,25,25	1.57	3 (12%)	31,35,35	1.82	7 (22%)
5	SO4	B	507	-	4,4,4	0.15	0	6,6,6	0.14	0
4	OLC	A	503	-	24,24,24	0.83	2 (8%)	25,25,25	0.98	2 (8%)
7	1PE	B	505	-	15,15,15	0.52	0	14,14,14	0.86	0
5	SO4	A	504	-	4,4,4	0.14	0	6,6,6	0.10	0
4	OLC	B	501	-	24,24,24	0.80	2 (8%)	25,25,25	0.98	1 (4%)
3	OLA	A	501	-	12,12,19	0.31	0	11,11,19	0.67	0
5	SO4	B	508	-	4,4,4	0.15	0	6,6,6	0.07	0
2	1Q5	C	401	-	24,25,25	1.56	3 (12%)	31,35,35	1.79	7 (22%)
3	OLA	C	501	-	19,19,19	0.50	0	19,19,19	1.03	0
5	SO4	B	506	-	4,4,4	0.13	0	6,6,6	0.06	0
6	PGW	B	502	-	38,38,50	0.97	3 (7%)	42,43,56	1.25	3 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OLC	A	502	-	-	9/24/24/24	-
6	PGW	B	503	-	-	17/47/47/55	-
2	1Q5	A	401	-	-	2/12/12/12	0/2/2/2
6	PGW	B	504	-	-	18/38/38/55	-
2	1Q5	B	401	-	-	3/12/12/12	0/2/2/2
4	OLC	A	503	-	-	17/24/24/24	-
7	1PE	B	505	-	-	6/13/13/13	-
4	OLC	B	501	-	-	9/24/24/24	-
3	OLA	A	501	-	-	4/10/10/17	-
2	1Q5	C	401	-	-	3/12/12/12	0/2/2/2
3	OLA	C	501	-	-	9/17/17/17	-
6	PGW	B	502	-	-	14/40/40/55	-

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	1Q5	C1-C6	4.40	1.47	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	1Q5	C1-C6	4.39	1.47	1.40
2	C	401	1Q5	C1-C6	4.30	1.47	1.40
2	A	401	1Q5	C1-C2	4.11	1.47	1.40
2	C	401	1Q5	C1-C2	3.98	1.47	1.40

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	401	1Q5	C1-O10-C11	5.37	122.38	116.88
2	B	401	1Q5	C1-O10-C11	4.27	121.26	116.88
6	B	503	PGW	O01-C1-C2	4.16	120.47	111.50
6	B	504	PGW	O01-C1-C2	4.14	120.43	111.50
2	C	401	1Q5	C1-O10-C11	3.95	120.92	116.88

There are no chirality outliers.

5 of 111 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	1Q5	C23-C20-C21-C22
2	B	401	1Q5	N19-C20-C21-C22
2	C	401	1Q5	N19-C20-C23-C24
3	A	501	OLA	C6-C7-C8-C9
4	A	503	OLC	C21-C22-C24-O25

There are no ring outliers.

13 monomers are involved in 31 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	502	OLC	3	0
6	B	503	PGW	1	0
2	A	401	1Q5	2	0
6	B	504	PGW	5	0
2	B	401	1Q5	5	0
4	A	503	OLC	1	0
5	A	504	SO4	1	1
4	B	501	OLC	2	0
3	A	501	OLA	1	0
2	C	401	1Q5	3	0
3	C	501	OLA	1	0
5	B	506	SO4	0	1
6	B	502	PGW	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand OLC A 502

Bond lengths   Bond angles

Torsions   Rings

Ligand PGW B 503

Bond lengths

Bond angles

Torsions

Rings

Ligand 1Q5 A 401

Bond lengths

Bond angles

Torsions

Rings

Ligand PGW B 504

Bond lengths

Bond angles

Torsions

Rings

Ligand 1Q5 B 401

Bond lengths

Bond angles

Torsions

Rings

Ligand OLC A 503

Bond lengths

Bond angles

Torsions

Rings

Ligand OLC B 501

Bond lengths

Bond angles

Torsions

Rings

Ligand OLA A 501

Bond lengths

Bond angles

Torsions

Rings

Ligand 1Q5 C 401

Bond lengths

Bond angles

Torsions

Rings

Ligand OLA C 501

Bond lengths

Bond angles

Torsions

Rings

Ligand PGW B 502

Bond lengths

Bond angles

Torsions

Rings

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	407/441 (92%)	0.24	33 (8%)	12	6	32, 56, 112, 131	0
1	B	396/441 (89%)	0.02	13 (3%)	46	28	28, 50, 92, 116	0
1	C	248/441 (56%)	0.37	27 (10%)	5	3	37, 66, 115, 139	0
All	All	1051/1323 (79%)	0.19	73 (6%)	16	9	28, 55, 109, 139	0

The worst 5 of 73 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	264	PRO	6.5
1	A	116	TYR	6.3
1	A	1054	SER	5.4
1	C	300	ALA	5.1
1	C	117	HIS	5.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

WORLDWIDE
wwPDB
PROTEIN DATA BANK

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
6	PGW	B	504	37/51	0.83	0.26	41,54,75,81	0
6	PGW	B	502	39/51	0.84	0.31	31,53,90,114	0
4	OLC	A	503	25/25	0.84	0.22	28,50,67,70	0
6	PGW	B	503	46/51	0.85	0.26	29,49,93,111	0
4	OLC	B	501	25/25	0.87	0.27	25,58,79,83	0
4	OLC	A	502	25/25	0.89	0.19	41,53,77,82	0
3	OLA	C	501	20/20	0.89	0.22	35,47,79,79	0
7	1PE	B	505	16/16	0.91	0.30	45,56,64,66	0
3	OLA	A	501	13/20	0.92	0.20	28,39,47,47	0
5	SO4	B	508	5/5	0.93	0.17	69,80,92,103	0
2	1Q5	C	401	24/24	0.94	0.21	47,52,58,69	0
2	1Q5	B	401	24/24	0.95	0.19	35,47,52,54	0
2	1Q5	A	401	24/24	0.96	0.20	43,52,58,60	0
5	SO4	B	506	5/5	0.97	0.12	62,64,72,76	0
5	SO4	A	504	5/5	0.97	0.10	61,62,73,74	0
5	SO4	B	507	5/5	0.99	0.18	40,48,54,55	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around PGW B 504:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around PGW B 502:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around OLC A 503:

$2mF_o\text{-}DF_c$ (at 0.7 rmsd) in gray
$mF_o\text{-}DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around PGW B 503:

$2mF_o\text{-}DF_c$ (at 0.7 rmsd) in gray
$mF_o\text{-}DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around OLC B 501:**

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around OLC A 502:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around OLA C 501:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around OLA A 501:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around 1Q5 C 401:

$2mF_o\text{-}DF_c$ (at 0.7 rmsd) in gray
$mF_o\text{-}DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around 1Q5 B 401:

$2mF_o\text{-}DF_c$ (at 0.7 rmsd) in gray
$mF_o\text{-}DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around 1Q5 A 401:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)

6.5 Other polymers ⓘ

There are no such residues in this entry.