



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 19, 2023 – 08:50 PM EDT

PDB ID : 5K5R
Title : AspA-32mer DNA,crystal form 2
Authors : Schumacher, M.
Deposited on : 2016-05-23
Resolution : 3.09 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	98	 2% (poor fit), 57% (0 outliers), 32% (1 outlier), 9% (2+ outliers)
1	B	98	 58% (0 outliers), 32% (1 outlier), 6% (2+ outliers)
1	C	98	 2% (poor fit), 51% (0 outliers), 35% (1 outlier), 7% (2 outliers), 7% (2+ outliers)
1	D	98	 16% (poor fit), 42% (0 outliers), 41% (1 outlier), 9% (2 outliers), 8% (2+ outliers)
1	E	98	 % (poor fit), 38% (0 outliers), 48% (1 outlier), 7% (2 outliers), 7% (2+ outliers)

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	98	
2	P	32	
3	N	32	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PO4	E	103	-	-	-	X
4	PO4	F	102	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5778 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AspA.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	C	91	738	481	122	135	0	0	0
1	D	90	731	476	121	134	0	0	0
1	A	89	726	472	120	134	0	0	0
1	B	92	747	486	124	137	0	0	0
1	E	91	738	481	122	135	0	0	0
1	F	92	747	486	124	137	0	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	94	HIS	-	expression tag	UNP O93706
C	95	HIS	-	expression tag	UNP O93706
C	96	HIS	-	expression tag	UNP O93706
C	97	HIS	-	expression tag	UNP O93706
C	98	HIS	-	expression tag	UNP O93706
C	99	HIS	-	expression tag	UNP O93706
D	94	HIS	-	expression tag	UNP O93706
D	95	HIS	-	expression tag	UNP O93706
D	96	HIS	-	expression tag	UNP O93706
D	97	HIS	-	expression tag	UNP O93706
D	98	HIS	-	expression tag	UNP O93706
D	99	HIS	-	expression tag	UNP O93706
A	94	HIS	-	expression tag	UNP O93706
A	95	HIS	-	expression tag	UNP O93706
A	96	HIS	-	expression tag	UNP O93706
A	97	HIS	-	expression tag	UNP O93706
A	98	HIS	-	expression tag	UNP O93706

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	99	HIS	-	expression tag	UNP O93706
B	94	HIS	-	expression tag	UNP O93706
B	95	HIS	-	expression tag	UNP O93706
B	96	HIS	-	expression tag	UNP O93706
B	97	HIS	-	expression tag	UNP O93706
B	98	HIS	-	expression tag	UNP O93706
B	99	HIS	-	expression tag	UNP O93706
E	94	HIS	-	expression tag	UNP O93706
E	95	HIS	-	expression tag	UNP O93706
E	96	HIS	-	expression tag	UNP O93706
E	97	HIS	-	expression tag	UNP O93706
E	98	HIS	-	expression tag	UNP O93706
E	99	HIS	-	expression tag	UNP O93706
F	94	HIS	-	expression tag	UNP O93706
F	95	HIS	-	expression tag	UNP O93706
F	96	HIS	-	expression tag	UNP O93706
F	97	HIS	-	expression tag	UNP O93706
F	98	HIS	-	expression tag	UNP O93706
F	99	HIS	-	expression tag	UNP O93706

- Molecule 2 is a DNA chain called DNA (32-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	P	32	653	315	114	193	31	0	0	0


- Molecule 3 is a DNA chain called DNA (32-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	N	32	653	315	120	187	31	0	0	0



- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		
4	E	1	Total	O	P	0	0
			5	4	1		
4	E	1	Total	O	P	0	0
			5	4	1		
4	E	1	Total	O	P	0	0
			5	4	1		
4	F	1	Total	O	P	0	0
			5	4	1		
4	F	1	Total	O	P	0	0
			5	4	1		
4	F	1	Total	O	P	0	0
			5	4	1		



- Molecule 1: AspA



- Molecule 2: DNA (32-MER)



- Molecule 3: DNA (32-MER)



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	150.24Å 55.88Å 101.62Å 90.00° 110.50° 90.00°	Depositor
Resolution (Å)	48.97 – 3.09 51.94 – 3.09	Depositor EDS
% Data completeness (in resolution range)	96.7 (48.97-3.09) 96.6 (51.94-3.09)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.99 (at 3.07Å)	Xtrriage
Refinement program	PHENIX 1.6.4_486	Depositor
R, R_{free}	0.202 , 0.275 0.198 , 0.270	Depositor DCC
R_{free} test set	1430 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å ²)	82.2	Xtrriage
Anisotropy	0.467	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 65.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5778	wwPDB-VP
Average B, all atoms (Å ²)	88.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.21	0/737	0.39	0/994
1	B	0.22	0/758	0.41	0/1021
1	C	0.21	0/749	0.40	0/1009
1	D	0.21	0/742	0.39	0/999
1	E	0.22	0/749	0.41	0/1009
1	F	0.22	0/758	0.41	0/1021
2	P	0.41	0/731	1.25	5/1127 (0.4%)
3	N	0.42	0/733	1.18	2/1129 (0.2%)
All	All	0.28	0/5957	0.72	7/8309 (0.1%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	P	25	DC	O4'-C1'-N1	8.27	113.79	108.00
2	P	25	DC	C3'-C2'-C1'	-6.51	94.69	102.50
2	P	8	DG	O4'-C4'-C3'	-6.07	102.07	104.50
3	N	19	DG	C3'-C2'-C1'	-6.06	95.23	102.50
3	N	19	DG	O4'-C1'-N9	5.76	112.03	108.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	726	0	766	31	0
1	B	747	0	793	41	0
1	C	738	0	785	42	0
1	D	731	0	776	56	0
1	E	738	0	785	44	0
1	F	747	0	793	43	0
2	P	653	0	366	33	0
3	N	653	0	364	29	0
4	A	10	0	0	0	0
4	D	5	0	0	0	0
4	E	15	0	0	0	0
4	F	15	0	0	0	0
All	All	5778	0	5428	264	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

The worst 5 of 264 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:92:VAL:O	1:A:93:GLN:HB3	1.69	0.90
3:N:43:DC:H2'	3:N:44:DA:C8	2.11	0.86
1:C:37:THR:HG23	1:C:39:ILE:H	1.45	0.82
2:P:26:DG:H1	3:N:25:DT:H3	1.28	0.81
1:C:27:LYS:HB2	1:C:69:LYS:HB3	1.63	0.80

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	87/98 (89%)	81 (93%)	6 (7%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	90/98 (92%)	83 (92%)	7 (8%)	0	100	100
1	C	89/98 (91%)	76 (85%)	13 (15%)	0	100	100
1	D	88/98 (90%)	76 (86%)	12 (14%)	0	100	100
1	E	89/98 (91%)	81 (91%)	8 (9%)	0	100	100
1	F	90/98 (92%)	76 (84%)	13 (14%)	1 (1%)	14	46
All	All	533/588 (91%)	473 (89%)	59 (11%)	1 (0%)	47	79

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	41	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	79/87 (91%)	75 (95%)	4 (5%)	24	56
1	B	81/87 (93%)	74 (91%)	7 (9%)	10	37
1	C	80/87 (92%)	67 (84%)	13 (16%)	2	10
1	D	79/87 (91%)	68 (86%)	11 (14%)	3	15
1	E	80/87 (92%)	64 (80%)	16 (20%)	1	5
1	F	81/87 (93%)	73 (90%)	8 (10%)	8	29
All	All	480/522 (92%)	421 (88%)	59 (12%)	4	19

5 of 59 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	6	THR
1	F	66	ILE
1	E	4	ILE
1	F	64	GLU
1	E	86	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	93	GLN
1	E	47	ASN
1	F	76	GLN
1	F	38	GLN
1	F	46	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PO4	E	101	-	4,4,4	0.89	0	6,6,6	0.45	0
4	PO4	A	101	-	4,4,4	0.91	0	6,6,6	0.38	0
4	PO4	A	102	-	4,4,4	0.88	0	6,6,6	0.42	0
4	PO4	D	101	-	4,4,4	0.91	0	6,6,6	0.42	0
4	PO4	E	102	-	4,4,4	0.88	0	6,6,6	0.43	0
4	PO4	F	102	-	4,4,4	0.91	0	6,6,6	0.41	0
4	PO4	E	103	-	4,4,4	0.90	0	6,6,6	0.43	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PO4	F	101	-	4,4,4	0.90	0	6,6,6	0.45	0
4	PO4	F	103	-	4,4,4	0.90	0	6,6,6	0.43	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	89/98 (90%)	-0.07	2 (2%) 62 41	59, 80, 112, 121	0
1	B	92/98 (93%)	-0.19	0 100 100	45, 59, 94, 117	0
1	C	91/98 (92%)	-0.03	2 (2%) 62 41	58, 85, 121, 126	0
1	D	90/98 (91%)	0.98	16 (17%) 1 0	57, 98, 141, 155	0
1	E	91/98 (92%)	0.13	1 (1%) 80 64	62, 86, 136, 145	0
1	F	92/98 (93%)	0.37	6 (6%) 18 8	71, 106, 136, 149	0
2	P	32/32 (100%)	-0.36	0 100 100	58, 85, 119, 141	0
3	N	32/32 (100%)	-0.30	0 100 100	62, 84, 122, 132	0
All	All	609/652 (93%)	0.14	27 (4%) 34 17	45, 86, 127, 155	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	62	LYS	7.2
1	D	61	GLN	5.9
1	D	60	GLU	5.3
1	C	7	ASP	5.3
1	D	64	GLU	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PO4	F	102	5/5	0.70	0.48	138,168,174,176	0
4	PO4	E	103	5/5	0.79	0.41	143,145,154,163	0
4	PO4	A	101	5/5	0.84	0.57	107,107,125,134	0
4	PO4	A	102	5/5	0.84	0.40	88,105,129,136	0
4	PO4	E	102	5/5	0.86	0.33	87,93,118,130	0
4	PO4	F	103	5/5	0.86	0.43	113,120,129,145	0
4	PO4	D	101	5/5	0.90	0.26	109,110,127,129	0
4	PO4	F	101	5/5	0.91	0.63	106,110,125,134	0
4	PO4	E	101	5/5	0.93	0.56	110,110,119,140	0

6.5 Other polymers [i](#)

There are no such residues in this entry.