

wwPDB X-ray Structure Validation Summary Report (i)

Jul 21, 2021 – 12:06 PM EDT

PDB ID	:	7K5M
Title	:	CRYSTAL STRUCTURE OF HBV CAPSID Y132A MUTANT IN COM-
		PLEX WITH N-(3-chloro-4-fluorophenyl)-3-phenyl-1,4,6,7-tetrahydro-5
		H-pyrazolo[4,3-c]pyridine-5-carboxamide AT 2.65A RESOLUTION
Authors	:	Kuduk, S.D.; Stoops, B.; Alexander, R.; Lam, A.M.; Espiritu, C.; Vogel, R.;
		Lau, V.; Klumpp, K.; Flores, O.A.; Hartman, G.D.; Lukacs, C.M.; Abendroth,
		J.
Deposited on	:	2020-09-17
Resolution	:	2.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


Xtriage (Phenix) EDS buster-report Percentile statistics Refmac	:::::::::::::::::::::::::::::::::::::::	20191225.v01 (using entries in the PDB archive December 25th 2019) 5.8.0158 7.0.044 (Gargrove) Engh & Huber (2001)
Ideal geometry (DNA, RNA) Validation Pipeline (wwPDB-VP)		Parkinson et al. (1996) 2.22

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	1332 (2.68-2.64)
Clashscore	141614	1374(2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	А	158	70%	16%	14%		
1	В	158	% 7 0%	13%	16%		
1	С	158	% 69%	15%	16%		
1	D	158	71%	12%	17%		
1	Е	158	% 66%	11% •	22%		



Mol	Chain	Length	Quality of chain				
			2%				
1	\mathbf{F}	158	66%	16%	18%		



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 6216 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	136	Total	С	Ν	0	S	0	0	0
1	Π	150	1037	674	175	183	5	0	0	0
1	В	132	Total	С	Ν	Ο	\mathbf{S}	0	1	0
1	D	152	1031	669	172	185	5	0	1	0
1	С	133	Total	С	Ν	Ο	\mathbf{S}	0	0	0
1	U	100	1015	665	169	176	5	0	0	0
1	D	131	Total	С	Ν	Ο	\mathbf{S}	0	0	0
1	D	101	1001	653	168	175	5	0	0	U
1	Е	124	Total	С	Ν	Ο	\mathbf{S}	0	0	0
	Ľ	124	963	628	161	169	5	0	0	0
1	F	129	Total	С	Ν	Ο	S	0	0	0
	Ľ	129	990	645	164	176	5		0	U

• Molecule 1 is a protein called Capsid protein.

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	0	SER	-	expression tag	UNP P03147
А	132	ALA	TYR	engineered mutation	UNP P03147
А	150	LYS	-	expression tag	UNP P03147
А	151	LEU	-	expression tag	UNP P03147
А	152	GLU	-	expression tag	UNP P03147
А	153	ASN	-	expression tag	UNP P03147
А	154	LEU	-	expression tag	UNP P03147
А	155	TYR	-	expression tag	UNP P03147
А	156	PHE	-	expression tag	UNP P03147
А	157	GLN	-	expression tag	UNP P03147
В	0	SER	-	expression tag	UNP P03147
В	132	ALA	TYR	engineered mutation	UNP P03147
В	150	LYS	-	expression tag	UNP P03147
В	151	LEU	-	expression tag	UNP P03147
В	152	GLU	-	expression tag	UNP P03147
В	153	ASN	-	expression tag	UNP P03147
В	154	LEU	-	expression tag	UNP P03147



Chain	Residue	Modelled	Actual	Comment	Reference
В	155	TYR	-	expression tag	UNP P03147
В	156	PHE	-	expression tag	UNP P03147
В	157	GLN	-	expression tag	UNP P03147
С	0	SER	-	expression tag	UNP P03147
С	132	ALA	TYR	engineered mutation	UNP P03147
С	150	LYS	-	expression tag	UNP P03147
С	151	LEU	_	expression tag	UNP P03147
С	152	GLU	-	expression tag	UNP P03147
С	153	ASN	-	expression tag	UNP P03147
С	154	LEU	-	expression tag	UNP P03147
С	155	TYR	-	expression tag	UNP P03147
С	156	PHE	-	expression tag	UNP P03147
С	157	GLN	-	expression tag	UNP P03147
D	0	SER	-	expression tag	UNP P03147
D	132	ALA	TYR	engineered mutation	UNP P03147
D	150	LYS	-	expression tag	UNP P03147
D	151	LEU	-	expression tag	UNP P03147
D	152	GLU	-	expression tag	UNP P03147
D	153	ASN	_	expression tag	UNP P03147
D	154	LEU	-	expression tag	UNP P03147
D	155	TYR	-	expression tag	UNP P03147
D	156	PHE	-	expression tag	UNP P03147
D	157	GLN	-	expression tag	UNP P03147
Е	0	SER	-	expression tag	UNP P03147
Е	132	ALA	TYR	engineered mutation	UNP P03147
Е	150	LYS	-	expression tag	UNP P03147
Е	151	LEU	-	expression tag	UNP P03147
Е	152	GLU	-	expression tag	UNP P03147
Е	153	ASN	-	expression tag	UNP P03147
Е	154	LEU	-	expression tag	UNP P03147
Е	155	TYR	-	expression tag	UNP P03147
Е	156	PHE	-	expression tag	UNP P03147
Е	157	GLN	-	expression tag	UNP P03147
F	0	SER	-	expression tag	UNP P03147
F	132	ALA	TYR	engineered mutation	UNP P03147
F	150	LYS	-	expression tag	UNP P03147
F	151	LEU	-	expression tag	UNP P03147
F	152	GLU	-	expression tag	UNP P03147
F	153	ASN	-	expression tag	UNP P03147
F	154	LEU	-	expression tag	UNP P03147
F	155	TYR	-	expression tag	UNP P03147
F	156	PHE	-	expression tag	UNP P03147


Continued from previous page...

expression tag UNP P03147 Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
F	157	GLN	-	expression tag	UNP P03147


• Molecule 2 is N-(3-chloro-4-fluorophenyl)-3-phenyl-2,4,6,7-tetrahydro-5H-pyrazolo[4,3-c]pyri dine-5-carboxamide (three-letter code: VXJ) (formula: C₁₉H₁₆ClFN₄O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total C Cl F N O 26 19 1 1 4 1	0	0
2	В	1	Total C Cl F N O 26 19 1 1 4 1	0	0
2	С	1	Total C Cl F N O 26 19 1 1 4 1	0	0
2	D	1	Total C Cl F N O 26 19 1 1 4 1	0	0
2	Е	1	Total C Cl F N O 26 19 1 1 4 1	0	0
2	F	1	Total C Cl F N O 26 19 1 1 4 1	0	0

• Molecule 3 is ISOPROPYL ALCOHOL (three-letter code: IPA) (formula: C₃H₈O).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 3 & 1 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 3 & 1 \end{array}$	0	0
3	D	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 3 & 1 \end{array}$	0	0
3	Е	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 3 & 1 \end{array}$	0	0
3	Е	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 3 & 1 \end{array}$	0	0


• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	С	2	Total O 2 2	0	0
4	Е	1	Total O 1 1	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Capsid protein







4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	151.58Å 87.96Å 106.71Å	Depositor
a, b, c, α , β , γ	90.00° 103.53° 90.00°	Depositor
Resolution (Å)	48.03 - 2.65	Depositor
Resolution (A)	48.04 - 2.65	EDS
% Data completeness	92.9 (48.03-2.65)	Depositor
(in resolution range)	93.2(48.04-2.65)	EDS
R _{merge}	0.07	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.79 (at 2.65 \text{\AA})$	Xtriage
Refinement program	PHENIX dev-1932	Depositor
D D.	0.242 , 0.276	Depositor
R, R_{free}	0.241 , 0.275	DCC
R_{free} test set	1983 reflections (5.35%)	wwPDB-VP
Wilson B-factor $(Å^2)$	42.1	Xtriage
Anisotropy	1.274	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.29, 58.0	EDS
L-test for twinning ²	$ \langle L \rangle = 0.46, \langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6216	wwPDB-VP
Average B, all atoms $(Å^2)$	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.61% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: IPA, VXJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
WIOI	Ullalli	RMSZ # Z > 5		RMSZ	# Z > 5	
1	А	0.24	0/1069	0.41	0/1466	
1	В	0.22	0/1063	0.38	0/1455	
1	С	0.22	0/1047	0.38	0/1437	
1	D	0.22	0/1032	0.40	0/1415	
1	Ε	0.22	0/992	0.40	0/1357	
1	F	0.24	0/1021	0.39	0/1400	
All	All	0.23	0/6224	0.39	0/8530	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1037	0	966	15	0
1	В	1031	0	973	17	0
1	С	1015	0	961	18	0
1	D	1001	0	948	14	0
1	Е	963	0	930	15	0
1	F	990	0	934	18	0
2	А	26	0	0	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	В	26	0	0	0	0
2	С	26	0	0	0	0
2	D	26	0	0	0	0
2	Ε	26	0	0	0	0
2	F	26	0	0	0	0
3	А	4	0	8	1	0
3	В	4	0	8	0	0
3	D	4	0	8	1	0
3	Е	8	0	16	2	0
4	С	2	0	0	0	0
4	Е	1	0	0	0	0
All	All	6216	0	5752	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 92 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:16:LEU:HD23	1:E:99:GLN:HB3	1.66	0.77
1:A:1:MET:HA	1:B:39:ARG:HH12	1.58	0.68
1:E:95:LEU:O	1:E:99:GLN:HG3	1.98	0.64
1:D:39:ARG:NH1	1:E:14:GLU:OE1	2.33	0.62
1:C:21:SER:O	1:C:98:ARG:NH2	2.32	0.62

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	132/158~(84%)	131 (99%)	1 (1%)	0	100 100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	\mathbf{ntiles}
1	В	129/158~(82%)	128~(99%)	1 (1%)	0	100	100
1	С	129/158~(82%)	128 (99%)	1 (1%)	0	100	100
1	D	127/158~(80%)	126 (99%)	1 (1%)	0	100	100
1	Ε	120/158~(76%)	119 (99%)	1 (1%)	0	100	100
1	F	125/158~(79%)	124 (99%)	1 (1%)	0	100	100
All	All	762/948~(80%)	756 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	103/140~(74%)	103 (100%)	0	100	100
1	В	106/140~(76%)	106 (100%)	0	100	100
1	С	102/140~(73%)	102 (100%)	0	100	100
1	D	101/140~(72%)	101 (100%)	0	100	100
1	Е	100/140~(71%)	99~(99%)	1 (1%)	76	86
1	F	101/140~(72%)	101 (100%)	0	100	100
All	All	613/840~(73%)	612~(100%)	1 (0%)	93	97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Ε	16	LEU

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such side chains are listed below:

Mol	Chain	Res	Type
1	А	47	HIS
1	С	99	GLN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	les
	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
3	IPA	Е	203	-	$3,\!3,\!3$	0.53	0	3,3,3	0.33	0
3	IPA	А	202	-	3,3,3	0.54	0	3,3,3	0.29	0
2	VXJ	D	201	-	27,29,29	1.94	9 (33%)	32,41,41	0.96	2 (6%)
3	IPA	D	202	-	3,3,3	0.53	0	3,3,3	0.33	0
3	IPA	Е	202	-	3,3,3	0.52	0	3,3,3	0.31	0
2	VXJ	С	500	-	27,29,29	1.94	8 (29%)	32,41,41	1.04	3 (9%)
3	IPA	В	202	-	3,3,3	0.54	0	3,3,3	0.28	0
2	VXJ	А	201	-	27,29,29	1.93	8 (29%)	32,41,41	1.01	3 (9%)
2	VXJ	Е	201	-	27,29,29	1.93	8 (29%)	32,41,41	1.14	4 (12%)
2	VXJ	F	500	-	$27,\!29,\!29$	2.00	8 (29%)	32,41,41	0.99	2 (6%)
2	VXJ	В	201	-	$27,\!29,\!29$	1.96	8 (29%)	32,41,41	1.06	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	VXJ	D	201	-	-	0/12/21/21	0/4/4/4
2	VXJ	С	500	-	-	0/12/21/21	0/4/4/4
2	VXJ	А	201	-	-	0/12/21/21	0/4/4/4
2	VXJ	Е	201	-	-	2/12/21/21	0/4/4/4
2	VXJ	F	500	-	-	2/12/21/21	0/4/4/4
2	VXJ	В	201	-	-	4/12/21/21	0/4/4/4

The worst 5 of 49 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
2	F	500	VXJ	C02-N12	5.14	1.46	1.36
2	В	201	VXJ	C02-N12	5.11	1.45	1.36
2	А	201	VXJ	C02-N12	5.07	1.45	1.36
2	С	500	VXJ	C02-N12	5.02	1.45	1.36
2	D	201	VXJ	C02-N12	5.01	1.45	1.36

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
2	Е	201	VXJ	C14-C13-N12	2.97	113.54	110.04
2	Е	201	VXJ	N03-C02-N12	2.74	119.02	115.89
2	В	201	VXJ	C14-C13-N12	2.62	113.12	110.04
2	С	500	VXJ	N03-C02-N12	2.51	118.76	115.89
2	D	201	VXJ	C14-C13-N12	2.48	112.97	110.04

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	500	VXJ	C16-C18-C19-C20
2	F	500	VXJ	C16-C18-C19-C24
2	Е	201	VXJ	O01-C02-N03-C04
2	В	201	VXJ	N12-C02-N03-C04
2	Е	201	VXJ	N12-C02-N03-C04

There are no ring outliers.

3 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	А	202	IPA	1	0
3	D	202	IPA	1	0




Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	Ε	202	IPA	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$\mathbf{OWAB}(\mathbf{\AA}^2)$	Q < 0.9
1	А	136/158~(86%)	-0.28	0 100 100	48, 63, 106, 129	0
1	В	132/158~(83%)	-0.20	2 (1%) 73 71	55, 74, 118, 128	0
1	С	133/158~(84%)	-0.11	1 (0%) 86 85	62, 77, 119, 141	0
1	D	131/158~(82%)	-0.27	0 100 100	60, 75, 114, 153	0
1	Ε	124/158~(78%)	-0.16	2 (1%) 72 69	50, 76, 110, 148	0
1	F	129/158~(81%)	-0.10	3 (2%) 60 56	49, 67, 113, 135	0
All	All	785/948~(82%)	-0.19	8 (1%) 82 81	48, 73, 115, 153	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Е	68	LEU	3.6
1	F	71	TRP	3.3
1	F	68	LEU	2.8
1	F	70	THR	2.6
1	В	89	VAL	2.5

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} extsf{-}\mathbf{B} extsf{-}\mathbf{factors}(\mathbf{A}^2)$	Q<0.9
3	IPA	Е	203	4/4	0.75	0.14	78,82,83,86	0
3	IPA	А	202	4/4	0.78	0.13	63,67,73,80	0
3	IPA	Е	202	4/4	0.85	0.16	74,79,80,83	0
3	IPA	D	202	4/4	0.85	0.18	75,84,87,91	0
3	IPA	В	202	4/4	0.86	0.15	76,86,86,90	0
2	VXJ	С	500	26/26	0.89	0.24	$55,\!69,\!78,\!85$	1
2	VXJ	D	201	26/26	0.90	0.23	$55,\!63,\!80,\!87$	1
2	VXJ	В	201	26/26	0.91	0.19	55,66,74,78	1
2	VXJ	А	201	26/26	0.91	0.21	$55,\!68,\!79,\!85$	1
2	VXJ	Е	201	26/26	0.92	0.21	55,71,78,83	1
2	VXJ	F	500	26/26	0.93	0.22	55,64,71,76	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
























6.5 Other polymers (i)

There are no such residues in this entry.

