



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 15, 2024 – 12:11 PM JST

PDB ID : 8K55  
Title : Crystal structure of sulfur transferase from Frondihabitans sp. PAMC28461  
crystallized in the P1 space group  
Authors : Nguyen, D.L.; Do, H.  
Deposited on : 2023-07-21  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	 4% 90% 7% ..
1	B	306	 4% 86% 9% 5%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4924 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called sulfur transferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	299	2331	1471	406	453	1	0	0	0
1	B	292	2279	1437	399	442	1	0	0	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	158	158	158	0	0
2	B	156	156	156	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: sulfur transferase



- Molecule 1: sulfur transferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	35.25Å 61.60Å 74.08Å 89.06° 76.30° 79.64°	Depositor
Resolution (Å)	28.19 – 2.00 28.19 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.5 (28.19-2.00) 96.5 (28.19-2.00)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	15.87 (at 1.99Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.215 , 0.269 0.221 , 0.269	Depositor DCC
$R_{free}$ test set	1935 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.6	Xtrriage
Anisotropy	0.035	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 35.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.000 for -h,-k,-h+1	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4924	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.72	0/2396	0.86	2/3275 (0.1%)
1	B	0.73	0/2340	0.84	0/3194
All	All	0.72	0/4736	0.85	2/6469 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	136	TYR	CB-CG-CD2	-5.63	117.62	121.00
1	A	136	TYR	CB-CG-CD1	5.37	124.22	121.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2331	0	2188	12	0
1	B	2279	0	2141	18	0
2	A	158	0	0	1	0
2	B	156	0	0	3	0
All	All	4924	0	4329	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:174:ILE:HD11	1:A:237:LEU:HD11	1.68	0.74
1:B:10:ARG:NH1	1:B:220:THR:OG1	2.30	0.65
1:B:174:ILE:HD11	1:B:237:LEU:HD11	1.78	0.63
1:A:198:LEU:HD12	1:A:199:ARG:HG3	1.81	0.62
1:B:300:ILE:HG22	1:B:300:ILE:O	2.04	0.58
1:B:135:ALA:N	2:B:401:HOH:O	2.24	0.56
1:B:185:ALA:O	1:B:186:ARG:HG2	2.07	0.55
1:A:286:PRO:HA	1:B:69:GLN:HE22	1.72	0.54
1:A:231:TYR:HB3	1:A:237:LEU:HD12	1.90	0.53
1:B:173:LEU:HD23	1:B:244:VAL:HB	1.91	0.53
1:B:141:PRO:O	1:B:143:VAL:HG13	2.11	0.51
1:A:186:ARG:HD2	2:A:416:HOH:O	2.11	0.50
1:B:174:ILE:HD12	1:B:260:LEU:HD11	1.91	0.50
1:B:217:GLU:H	1:B:217:GLU:CD	2.14	0.50
1:A:90:ASP:OD1	1:A:90:ASP:N	2.47	0.46
1:B:51:HIS:NE2	1:B:139:ASP:O	2.48	0.46
1:A:38:VAL:O	1:A:95:ILE:HA	2.16	0.46
1:A:141:PRO:O	1:A:142:THR:O	2.34	0.46
1:B:42:GLU:OE2	1:B:99:LYS:NZ	2.50	0.44
1:B:143:VAL:HG23	2:B:451:HOH:O	2.17	0.44
1:A:188:THR:OG1	1:A:189:ALA:N	2.50	0.43
1:B:174:ILE:CD1	1:B:260:LEU:HD11	2.48	0.43
1:A:106:TYR:HA	1:A:254:SER:CB	2.49	0.42
1:B:204:PRO:HG3	1:B:288:VAL:HG13	2.01	0.42
1:B:47:TYR:HB2	2:B:446:HOH:O	2.20	0.41
1:A:10:ARG:NH1	1:A:220:THR:OG1	2.53	0.41
1:B:172:PRO:HD2	1:B:242:ASP:O	2.21	0.41
1:A:73:ILE:HB	1:A:77:ALA:HB3	2.02	0.41
1:B:106:TYR:HA	1:B:254:SER:CB	2.51	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	297/306 (97%)	290 (98%)	6 (2%)	1 (0%)	41	37
1	B	288/306 (94%)	274 (95%)	12 (4%)	2 (1%)	22	16
All	All	585/612 (96%)	564 (96%)	18 (3%)	3 (0%)	29	23

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	142	THR
1	B	142	THR
1	B	205	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/241 (98%)	232 (98%)	4 (2%)	60	65
1	B	231/241 (96%)	228 (99%)	3 (1%)	69	74
All	All	467/482 (97%)	460 (98%)	7 (2%)	65	69

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	90	ASP
1	A	103	TRP
1	A	198	LEU
1	A	253	SER
1	B	103	TRP
1	B	139	ASP
1	B	198	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:



Mol	Chain	Res	Type
1	A	61	HIS
1	A	168	HIS
1	B	69	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	299/306 (97%)	0.22	12 (4%) 38 37	14, 24, 44, 56	0
1	B	292/306 (95%)	0.15	11 (3%) 40 39	13, 24, 48, 78	0
All	All	591/612 (96%)	0.19	23 (3%) 39 38	13, 24, 46, 78	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	192	TYR	3.6
1	A	188	THR	3.4
1	A	191	ALA	3.3
1	B	187	THR	3.3
1	B	198	LEU	3.3
1	A	190	PRO	3.2
1	A	189	ALA	3.1
1	B	68	VAL	3.0
1	A	194	GLU	2.9
1	B	186	ARG	2.8
1	A	180	GLU	2.6
1	A	142	THR	2.6
1	B	95	ILE	2.5
1	A	193	PRO	2.4
1	B	121	LEU	2.3
1	A	141	PRO	2.3
1	B	107	ALA	2.3
1	B	185	ALA	2.3
1	B	139	ASP	2.2
1	A	185	ALA	2.2
1	B	300	ILE	2.0
1	A	140	ALA	2.0
1	B	141	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.