



# wwPDB X-ray Structure Validation Summary Report

Jan 30, 2024 – 10:45 PM EST

PDB ID : 1K74  
Title : The 2.3 Angstrom resolution crystal structure of the heterodimer of the human PPARgamma and RXRalpha ligand binding domains respectively bound with GW409544 and 9-cis retinoic acid and co-activator peptides.  
Authors : Xu, H.E.; Lambert, M.H.; Montana, V.G.; Moore, L.B.; Collins, J.L.; Oplinger, J.A.; Kliewer, S.A.; Gampe Jr., R.T.; McKee, D.D.; Moore, J.T.; Willson, T.M.  
Deposited on : 2001-10-18  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	238	63% 29% 5% . . .
2	D	283	58% 32% 5% . . .
3	B	25	24% 16% 60%
3	E	25	28% 36% 36%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	544	D	478	X	-	-	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4405 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Retinoic acid receptor RXR-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	232	1754	1117	306	321	10	0	0	0

- Molecule 2 is a protein called Peroxisome proliferator activated receptor gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	272	2168	1399	354	405	10	0	0	0


There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	195	MET	-	expression tag	UNP P37231
D	196	LYS	-	expression tag	UNP P37231
D	197	LYS	-	expression tag	UNP P37231
D	198	GLY	-	expression tag	UNP P37231
D	199	HIS	-	expression tag	UNP P37231
D	200	HIS	-	expression tag	UNP P37231
D	201	HIS	-	expression tag	UNP P37231
D	202	HIS	-	expression tag	UNP P37231
D	203	HIS	-	expression tag	UNP P37231
D	204	HIS	-	expression tag	UNP P37231
D	205	GLY	-	expression tag	UNP P37231

- Molecule 3 is a protein called steroid receptor coactivator.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	B	10	86	55	18	13	0	0	0
3	E	16	129	79	27	23	0	0	0

- Molecule 4 is (9cis)-retinoic acid (three-letter code: 9CR) (formula: C<sub>20</sub>H<sub>28</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			22	20	2		

- Molecule 5 is 2-(1-METHYL-3-OXO-3-PHENYL-PROPYLAMINO)-3-{4-[2-(5-METHYL-2-PHENYL-OXAZOL-4-YL)-ETHOXY]-PHENYL}-PROPIONIC ACID (three-letter code: 544) (formula: C<sub>31</sub>H<sub>30</sub>N<sub>2</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	D	1	Total	C	N	O	0	0
			38	31	2	5		

- Molecule 6 is water.


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
6	A	67	Total O 67 67	0	0
6	D	133	Total O 133 133	0	0
6	B	2	Total O 2 2	0	0
6	E	6	Total O 6 6	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: Retinoic acid receptor RXR-alpha



- Molecule 2: Peroxisome proliferator activated receptor gamma



- Molecule 3: steroid receptor coactivator



- Molecule 3: steroid receptor coactivator



CYS	E685
PRO	R686
SER	H687
SER	K688
HIS	I689
SER	L690
SER	H691
SER	R692
LEU	L693
THR	I694
	Q695
	E696
	G697
	S698
	P699
	S700



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.62Å 55.10Å 214.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.30	Depositor
% Data completeness (in resolution range)	(Not available) (20.00-2.30)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
Refinement program	CNX	Depositor
R, $R_{free}$	0.238 , 0.279	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	4405	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 9CR, 544

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.36	0/1789	0.62	2/2414 (0.1%)
2	D	0.41	0/2206	0.66	1/2974 (0.0%)
3	B	0.31	0/87	0.62	0/116
3	E	0.29	0/131	0.51	0/175
All	All	0.39	0/4213	0.64	3/5679 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	243	GLU	C-N-CD	-10.50	97.50	120.60
1	A	243	GLU	C-N-CA	6.57	149.59	122.00
2	D	206	PRO	N-CA-CB	5.53	109.94	103.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1754	0	1733	73	0
2	D	2168	0	2213	126	1
3	B	86	0	86	5	0
3	E	129	0	125	14	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	22	0	27	7	0
5	D	38	0	29	1	0
6	A	67	0	0	0	0
6	B	2	0	0	1	0
6	D	133	0	0	2	0
6	E	6	0	0	2	0
All	All	4405	0	4213	203	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

The worst 5 of 203 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:270:LEU:H	2:D:270:LEU:HD12	1.19	1.05
2:D:270:LEU:HD22	2:D:283:GLN:HB3	1.37	1.05
1:A:230:MET:HE3	1:A:235:ILE:HD11	1.38	1.05
2:D:268:THR:H	2:D:269:PRO:HD2	1.26	1.00
2:D:270:LEU:H	2:D:270:LEU:CD1	1.83	0.89

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:250:TYR:OH	2:D:269:PRO:O[4_455]	2.03	0.17

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	230/238 (97%)	194 (84%)	21 (9%)	15 (6%)	<b>1</b> <b>0</b>

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	D	270/283 (95%)	242 (90%)	16 (6%)	12 (4%)	2	1
3	B	8/25 (32%)	6 (75%)	1 (12%)	1 (12%)	0	0
3	E	14/25 (56%)	10 (71%)	3 (21%)	1 (7%)	1	0
All	All	522/571 (91%)	452 (87%)	41 (8%)	29 (6%)	2	1

5 of 29 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	243	GLU
1	A	244	PRO
1	A	256	LEU
1	A	257	ASN
1	A	259	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/205 (89%)	169 (93%)	13 (7%)	14	19
2	D	240/254 (94%)	221 (92%)	19 (8%)	12	15
3	B	9/24 (38%)	9 (100%)	0	100	100
3	E	14/24 (58%)	13 (93%)	1 (7%)	14	19
All	All	445/507 (88%)	412 (93%)	33 (7%)	13	17

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	440	THR
2	D	442	LEU
3	E	695	GLN
1	A	450	PHE
1	A	426	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	454	GLN
3	E	691	HIS
3	E	695	GLN
3	B	638	GLN
2	D	335	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	9CR	A	463	-	22,22,22	3.64	13 (59%)	30,30,30	1.83	8 (26%)
5	544	D	478	-	36,41,41	2.55	15 (41%)	41,55,55	1.47	4 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	9CR	A	463	-	-	6/15/32/32	0/1/1/1
5	544	D	478	-	1/1/4/6	0/25/30/30	0/4/4/4

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	463	9CR	C1-C6	10.11	1.67	1.53
4	A	463	9CR	C5-C6	9.58	1.51	1.34
5	D	478	544	C1A-N	8.25	1.44	1.33
5	D	478	544	C3E-C3D	4.60	1.54	1.48
5	D	478	544	C1F-C1G	4.57	1.58	1.43

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	478	544	C1F-C1A-N	-5.17	117.16	121.29
5	D	478	544	C1B-C1A-N	5.07	124.50	118.82
4	A	463	9CR	C19-C9-C10	-4.56	116.53	122.92
4	A	463	9CR	C19-C9-C8	3.46	123.52	118.08
4	A	463	9CR	C20-C13-C12	3.36	123.37	118.08

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	D	478	544	C1A

5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	463	9CR	C1-C6-C7-C8
4	A	463	9CR	C5-C6-C7-C8
4	A	463	9CR	C13-C14-C15-O2
4	A	463	9CR	C13-C14-C15-O1
4	A	463	9CR	C11-C10-C9-C19


There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	463	9CR	7	0
5	D	478	544	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.