

Full wwPDB X-ray Structure Validation Report (i)

Sep 10, 2023 – 09:40 AM EDT

PDB ID : 4KE3

Title : Crystal structure of a glutathione transferase family member from Burkholde-

ria graminis, target efi-507264, no gsh, disordered domains, space group P21,

form(2)

Authors: Vetting, M.W.; Toro, R.; Bhosle, R.; Al Obaidi, N.F.; Morisco, L.L.; Wasser-

man, S.R.; Sojitra, S.; Stead, M.; Washington, E.; Scott Glenn, A.; Chowdhury, S.; Evans, B.; Hammonds, J.; Hillerich, B.; Love, J.; Seidel, R.D.; Imker, H.J.; Gerlt, J.A.; Armstrong, R.N.; Almo, S.C.; Enzyme Function Initiative

(EFI)

Deposited on : 2013-04-25

Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467 Xtriage (Phenix) : 1.13

EDS : 2.35.1

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158


CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	229	82%	7% • 10%
1	В	229	83%	6% • 10%
1	С	229	76%	19%

Continued on next page...

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1



 $Continued\ from\ previous\ page...$

Mol	Chain	Length	Quality of chain		
1	D	229	75%	5% •	19%



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 12973 atoms, of which 6134 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Glutathione S-transferase domain.

Mol	Chain	Residues		Atoms					ZeroOcc	AltConf	Trace
1	A	207	Total	С	Н	N	О	S	0	0	0
1	Λ	201	3273	1053	1613	300	300	7	0	U	U
1	В	207	Total	С	Н	N	О	S	0	0	0
1	Ъ	201	3273	1053	1613	300	300	7	0		U
1	С	185	Total	С	Н	N	О	S	0	0	0
1		100	2944	943	1454	271	270	6	0	U	U
1	D	185	Total	С	Н	N	О	S	0	0	0
1	ש	100	2944	943	1454	271	270	6		U	U

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	expression tag	UNP B1FZ96
A	-20	HIS	-	expression tag	UNP B1FZ96
A	-19	HIS	-	expression tag	UNP B1FZ96
A	-18	HIS	-	expression tag	UNP B1FZ96
A	-17	HIS	-	expression tag	UNP B1FZ96
A	-16	HIS	-	expression tag	UNP B1FZ96
A	-15	HIS	-	expression tag	UNP B1FZ96
A	-14	SER	-	expression tag	UNP B1FZ96
A	-13	SER	-	expression tag	UNP B1FZ96
A	-12	GLY	-	expression tag	UNP B1FZ96
A	-11	VAL	-	expression tag	UNP B1FZ96
A	-10	ASP	-	expression tag	UNP B1FZ96
A	-9	LEU	-	expression tag	UNP B1FZ96
A	-8	GLY	-	expression tag	UNP B1FZ96
A	-7	THR	-	expression tag	UNP B1FZ96
A	-6	GLU	-	expression tag	UNP B1FZ96
A	-5	ASN	-	expression tag	UNP B1FZ96
A	-4	LEU	-	expression tag	UNP B1FZ96
A	-3	TYR	-	expression tag	UNP B1FZ96
A	-2	PHE	-	expression tag	UNP B1FZ96
A	-1	GLN	-	expression tag	UNP B1FZ96



 $Continued\ from\ previous\ page...$

B -1 B -1 B -1	21 20 .9	SER MET HIS HIS HIS	- - -	expression tag expression tag expression tag	UNP B1FZ96 UNP B1FZ96 UNP B1FZ96
B -2 B -1 B -1 B -1 B -1	20 .9 .8 .7	HIS HIS HIS		1 0	
B -1 B -1 B -1	9 .8 .7	HIS HIS	-	expression tag	UNP R1F706
B -1 B -1 B -1	.8	HIS	-		OMI DIL 190
B -1	.7			expression tag	UNP B1FZ96
В -1			-	expression tag	UNP B1FZ96
	.6	HIS	_	expression tag	UNP B1FZ96
В -1		HIS	-	expression tag	UNP B1FZ96
1	.5	HIS	-	expression tag	UNP B1FZ96
В -1	.4	SER	-	expression tag	UNP B1FZ96
В -1	.3	SER	-	expression tag	UNP B1FZ96
В -1	.2	GLY	-	expression tag	UNP B1FZ96
В -1	.1	VAL	-	expression tag	UNP B1FZ96
В -1	.0	ASP	-	expression tag	UNP B1FZ96
В -	9	LEU	-	expression tag	UNP B1FZ96
В -	8	GLY	-	expression tag	UNP B1FZ96
В -	7	THR	-	expression tag	UNP B1FZ96
В -	6	GLU	-	expression tag	UNP B1FZ96
В -	5	ASN	-	expression tag	UNP B1FZ96
В -	4	LEU	-	expression tag	UNP B1FZ96
В -	3	TYR	-	expression tag	UNP B1FZ96
В -	2	PHE	-	expression tag	UNP B1FZ96
В -	1	GLN	-	expression tag	UNP B1FZ96
В ()	SER	-	expression tag	UNP B1FZ96
C -2	21	MET	-	expression tag	UNP B1FZ96
C -2	20	HIS	-	expression tag	UNP B1FZ96
C -1	.9	HIS	-	expression tag	UNP B1FZ96
C -1	.8	HIS	-	expression tag	UNP B1FZ96
C -1	.7	HIS	-	expression tag	UNP B1FZ96
C -1	.6	HIS	-	expression tag	UNP B1FZ96
C -1	.5	HIS	-	expression tag	UNP B1FZ96
C -1	.4	SER	-	expression tag	UNP B1FZ96
C -1	.3	SER	-	expression tag	UNP B1FZ96
C -1	.2	GLY	-	expression tag	UNP B1FZ96
C -1	.1	VAL	-	expression tag	UNP B1FZ96
C -1	.0	ASP	-	expression tag	UNP B1FZ96
С -	9	LEU	-	expression tag	UNP B1FZ96
С -	8	GLY	-	expression tag	UNP B1FZ96
С -	7	THR	-	expression tag	UNP B1FZ96
С -	6	GLU	-	expression tag	UNP B1FZ96
С -	5	ASN	-	expression tag	UNP B1FZ96
С -	4	LEU	-	expression tag	UNP B1FZ96
С -	3	TYR	-	expression tag	UNP B1FZ96



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
С	-2	PHE	-	expression tag	UNP B1FZ96
С	-1	GLN	-	expression tag	UNP B1FZ96
С	0	SER	-	expression tag	UNP B1FZ96
D	-21	MET	-	expression tag	UNP B1FZ96
D	-20	HIS	-	expression tag	UNP B1FZ96
D	-19	HIS	-	expression tag	UNP B1FZ96
D	-18	HIS	-	expression tag	UNP B1FZ96
D	-17	HIS	-	expression tag	UNP B1FZ96
D	-16	HIS	-	expression tag	UNP B1FZ96
D	-15	HIS	-	expression tag	UNP B1FZ96
D	-14	SER	-	expression tag	UNP B1FZ96
D	-13	SER	-	expression tag	UNP B1FZ96
D	-12	GLY	-	expression tag	UNP B1FZ96
D	-11	VAL	-	expression tag	UNP B1FZ96
D	-10	ASP	-	expression tag	UNP B1FZ96
D	-9	LEU	-	expression tag	UNP B1FZ96
D	-8	GLY	-	expression tag	UNP B1FZ96
D	-7	THR	-	expression tag	UNP B1FZ96
D	-6	GLU	-	expression tag	UNP B1FZ96
D	-5	ASN	-	expression tag	UNP B1FZ96
D	-4	LEU	-	expression tag	UNP B1FZ96
D	-3	TYR	-	expression tag	UNP B1FZ96
D	-2	PHE	-	expression tag	UNP B1FZ96
D	-1	GLN	-	expression tag	UNP B1FZ96
D	0	SER	-	expression tag	UNP B1FZ96

• Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	145	Total O 145 145	0	0
2	В	133	Total O 133 133	0	0
2	С	122	Total O 122 122	0	0
2	D	139	Total O 139 139	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glutathione S-transferase domain





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	78.21Å 56.14Å 98.30Å	Donogiton
a, b, c, α , β , γ	90.00° 91.35° 90.00°	Depositor
Resolution (Å)	32.76 - 1.90	Depositor
Resolution (A)	32.76 - 1.89	EDS
% Data completeness	85.3 (32.76-1.90)	Depositor
(in resolution range)	85.3 (32.76-1.89)	EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$< I/\sigma(I) > 1$	3.70 (at 1.89Å)	Xtriage
Refinement program	PHENIX 1.8.1_1168	Depositor
D D.	0.153 , 0.199	Depositor
R, R_{free}	0.155 , 0.199	DCC
R_{free} test set	2979 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	21.5	Xtriage
Anisotropy	0.380	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.41 , 25.8	EDS
L-test for twinning ²	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	0.329 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12973	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.06% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Moi Chain		RMSZ $ \# Z > 5$		# Z > 5	
1	A	0.58	0/1704	0.71	2/2319 (0.1%)	
1	В	0.58	0/1704	0.69	0/2319	
1	С	0.59	0/1526	0.70	$2/2072 \ (0.1\%)$	
1	D	0.59	0/1526	0.68	0/2072	
All	All	0.59	0/6460	0.70	4/8782 (0.0%)	

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	С	71	ARG	NE-CZ-NH2	-5.98	117.31	120.30
1	A	191	ARG	NE-CZ-NH1	5.76	123.18	120.30
1	С	199	ARG	NE-CZ-NH1	5.71	123.16	120.30
1	A	33	GLY	N-CA-C	5.51	126.87	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1660	1613	1614	20	0
1	В	1660	1613	1614	10	0
1	С	1490	1454	1455	5	0
1	D	1490	1454	1455	10	0
2	A	145	0	0	5	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
2	В	133	0	0	1	1
2	С	122	0	0	1	1
2	D	139	0	0	3	0
All	All	6839	6134	6138	44	1

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 4.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A 1 - 1	A. 0	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:A:33:GLY:N	1:A:34:ALA:HB2	1.74	1.01
1:A:34:ALA:HB1	1:A:37:ARG:HB2	1.65	0.76
1:D:53:VAL:N	2:D:340:HOH:O	2.20	0.73
1:A:181:ASP:OD2	2:A:443:HOH:O	2.07	0.71
1:B:29:ARG:NH1	1:B:31:ASP:OD2	2.26	0.67
1:A:34:ALA:CB	1:A:37:ARG:O	2.43	0.66
1:A:1:MET:N	2:A:445:HOH:O	2.28	0.65
1:A:33:GLY:N	1:A:34:ALA:CB	2.58	0.65
1:B:50:ASN:OD1	1:C:141:ILE:HD13	2.03	0.59
1:A:117:ARG:NH2	2:A:404:HOH:O	2.24	0.57
1:A:34:ALA:HB1	1:A:37:ARG:CB	2.34	0.56
1:B:123:ARG:NH1	2:B:417:HOH:O	2.38	0.55
1:C:6:GLY:HA2	1:C:53:VAL:HG21	1.91	0.52
1:A:34:ALA:HB3	1:A:37:ARG:O	2.10	0.52
1:A:76:ARG:O	1:A:76:ARG:HG3	2.10	0.51
1:D:132:ARG:NH1	2:D:355:HOH:O	2.39	0.50
1:D:120:PRO:HA	1:D:123:ARG:HD2	1.94	0.50
1:A:32:TRP:C	1:A:34:ALA:HB2	2.31	0.50
1:A:33:GLY:H	1:A:34:ALA:HB2	1.68	0.50
1:D:1:MET:HE2	1:D:28:GLU:HG3	1.95	0.49
1:A:37:ARG:NE	2:A:363:HOH:O	2.39	0.49
1:D:25:LEU:HD21	1:D:77:TYR:HB3	1.96	0.48
1:D:53:VAL:HG22	1:D:53:VAL:O	2.14	0.47
1:D:1:MET:CE	1:D:28:GLU:HG3	2.45	0.46
1:A:33:GLY:CA	1:A:34:ALA:CB	2.95	0.45
1:B:127:GLN:OE1	1:B:127:GLN:HA	2.17	0.45
1:A:33:GLY:CA	1:A:34:ALA:HB2	2.46	0.45
1:B:34:ALA:O	1:B:36:PHE:N	2.50	0.44
1:A:110:TYR:CE1	1:A:122:HIS:HB3	2.54	0.43
1:B:60:ASP:C	1:B:60:ASP:OD1	2.57	0.43



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${ m distance}({ m \AA})$	overlap (Å)
1:D:108:TRP:C	1:D:108:TRP:CD1	2.92	0.42
1:C:148:ARG:NH1	2:C:382:HOH:O	2.46	0.42
1:A:37:ARG:CD	2:A:363:HOH:O	2.67	0.42
1:C:6:GLY:HA2	1:C:53:VAL:CG2	2.50	0.42
1:B:8:PRO:HD3	1:B:29:ARG:NH1	2.34	0.42
1:C:25:LEU:HD21	1:C:77:TYR:HB3	2.02	0.41
1:D:175:THR:HG23	1:D:176:PRO:HD2	2.02	0.41
1:A:124:ASP:O	1:A:128:ILE:HG13	2.20	0.41
1:B:60:ASP:OD1	1:B:60:ASP:O	2.39	0.41
1:A:30:GLU:OE1	1:A:37:ARG:NH1	2.51	0.41
1:B:170:ASN:OD1	1:B:203:ARG:NH1	2.53	0.40
1:A:4:ILE:HD12	1:A:27:PHE:CD1	2.55	0.40
1:B:108:TRP:CD1	1:B:108:TRP:C	2.95	0.40
1:D:95:GLN:CD	2:D:354:HOH:O	2.60	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
2:B:396:HOH:O	2:C:392:HOH:O[1_545]	2.18	0.02

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	$205/229 \ (90\%)$	199 (97%)	4 (2%)	2 (1%)	15 6
1	В	$205/229 \ (90\%)$	197 (96%)	7 (3%)	1 (0%)	29 18
1	С	181/229 (79%)	175 (97%)	6 (3%)	0	100 100
1	D	181/229 (79%)	174 (96%)	7 (4%)	0	100 100
All	All	772/916 (84%)	745 (96%)	24 (3%)	3 (0%)	34 24



All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	34	ALA
1	В	35	GLY
1	A	65	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	167/187 (89%)	165 (99%)	2 (1%)	71 70
1	В	167/187 (89%)	162 (97%)	5 (3%)	41 33
1	\mathbf{C}	150/187 (80%)	147 (98%)	3 (2%)	55 51
1	D	150/187 (80%)	147 (98%)	3 (2%)	55 51
All	All	$634/748 \; (85\%)$	621 (98%)	13 (2%)	53 48

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	108	TRP
1	A	119	SER
1	В	1	MET
1	В	10	SER
1	В	60	ASP
1	В	108	TRP
1	В	181	ASP
1	С	30	GLU
1	С	53	VAL
1	С	181	ASP
1	D	5	LEU
1	D	53	VAL
1	D	108	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$OWAB(Å^2)$	Q < 0.9
1	A	207/229 (90%)	-0.29	1 (0%) 91 92	6, 20, 47, 70	1 (0%)
1	В	207/229 (90%)	-0.21	1 (0%) 91 92	8, 21, 52, 71	2 (0%)
1	С	185/229 (80%)	-0.21	1 (0%) 91 92	9, 21, 52, 69	0
1	D	185/229 (80%)	-0.27	0 100 100	7, 19, 46, 63	1 (0%)
All	All	784/916 (85%)	-0.24	3 (0%) 92 93	6, 20, 51, 71	4 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	36	PHE	6.8
1	A	34	ALA	2.2
1	С	123	ARG	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.

