

### wwPDB X-ray Structure Validation Summary Report (i)

Nov 27, 2023 – 06:20 PM JST

PDB ID : 8KHQ

Title: Bifunctional sulfoxide synthase OvoA Th2 in complex with histidine and cys-

teine

Authors: Wang, J.; Ye, K.; Wang, X.Y.; Yan, W.P.

Deposited on : 2023-08-22

Resolution : 2.69 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

### 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	728	88%	8%	<del>-</del>
1	В	728	87%	9%	•
1	С	728	87%	9%	•
1	D	728	6% 85%	11%	•



### 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 23253 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called 5-histidylcysteine sulfoxide synthase/putative 4-mercaptohistidine N1-methyltransferase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Λ	700	Total	С	N	О	S	0	0	0
1	A	700	5784	3742	967	1056	19	0	0	
1	В	600	Total	С	N	О	S	0	0	0
1	Б	698	5763	3728	967	1050	18		U	
1	C	699	Total	С	N	О	S	0	0	0
1		099	5776	3735	970	1053	18	0	0	0
1	D	697	Total	С	N	О	S	0	0	0
1	ש	097	5693 3678 958 1038 19		U	U				

There are 116 discrepancies between the modelled and reference sequences:

Chain	Residue	Residue   Modelled   Actual   Comment		Comment	Reference
A	1	MET	-	initiating methionine	UNP A0A1I5R890
A	2	GLY	-	expression tag	UNP A0A1I5R890
A	3	ASP	-	expression tag	UNP A0A1I5R890
A	4	ARG	-	expression tag	UNP A0A1I5R890
A	5	GLY	-	expression tag	UNP A0A1I5R890
A	6	PRO	-	expression tag	UNP A0A1I5R890
A	7	GLU	-	expression tag	UNP A0A1I5R890
A	8	PHE	-	expression tag	UNP A0A1I5R890
A	708	LEU	-	expression tag	UNP A0A1I5R890
A	709	GLU	-	expression tag	UNP A0A1I5R890
A	710	VAL	_	expression tag	UNP A0A1I5R890
A	711	ASP	-	expression tag	UNP A0A1I5R890
A	712	LEU	_	expression tag	UNP A0A1I5R890
A	713	GLN	-	expression tag	UNP A0A1I5R890
A	714	GLY	-	expression tag	UNP A0A1I5R890
A	715	ASP	-	expression tag	UNP A0A1I5R890
A	716	HIS	-	expression tag	UNP A0A1I5R890
A	717	GLY	-	expression tag	UNP A0A1I5R890
A	718	LEU	-	expression tag	UNP A0A1I5R890
A	719	SER	-	expression tag	UNP A0A1I5R890



Continued from previous page...

Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
A	720	ALA	-	expression tag	UNP A0A1I5R890
A	721	TRP	_	expression tag	UNP A0A1I5R890
A	722	SER	-	expression tag	UNP A0A1I5R890
A	723	HIS	_	expression tag	UNP A0A1I5R890
A	724	PRO	-	expression tag	UNP A0A1I5R890
A	725	GLN	-	expression tag	UNP A0A1I5R890
A	726	PHE	-	expression tag	UNP A0A1I5R890
A	727	GLU	-	expression tag	UNP A0A1I5R890
A	728	LYS	-	expression tag	UNP A0A1I5R890
В	1	MET	-	initiating methionine	UNP A0A1I5R890
В	2	GLY	-	expression tag	UNP A0A1I5R890
В	3	ASP	-	expression tag	UNP A0A1I5R890
В	4	ARG	-	expression tag	UNP A0A1I5R890
В	5	GLY	-	expression tag	UNP A0A1I5R890
В	6	PRO	-	expression tag	UNP A0A1I5R890
В	7	GLU	-	expression tag	UNP A0A1I5R890
В	8	PHE	-	expression tag	UNP A0A1I5R890
В	708	LEU	-	expression tag	UNP A0A1I5R890
В	709	GLU	-	expression tag	UNP A0A1I5R890
В	710	VAL	-	expression tag	UNP A0A1I5R890
В	711	ASP	-	expression tag	UNP A0A1I5R890
В	712	LEU	-	expression tag	UNP A0A1I5R890
В	713	GLN	-	expression tag	UNP A0A1I5R890
В	714	GLY	-	expression tag	UNP A0A1I5R890
В	715	ASP	-	expression tag	UNP A0A1I5R890
В	716	HIS	-	expression tag	UNP A0A1I5R890
В	717	GLY	-	expression tag	UNP A0A1I5R890
В	718	LEU	-	expression tag	UNP A0A1I5R890
В	719	SER	-	expression tag	UNP A0A1I5R890
В	720	ALA	-	expression tag	UNP A0A1I5R890
В	721	TRP	-	expression tag	UNP A0A1I5R890
В	722	SER	-	expression tag	UNP A0A1I5R890
В	723	HIS	-	expression tag	UNP A0A1I5R890
В	724	PRO	-	expression tag	UNP A0A1I5R890
В	725	GLN	-	expression tag	UNP A0A1I5R890
В	726	PHE	-	expression tag	UNP A0A1I5R890
В	727	GLU		expression tag	UNP A0A1I5R890
В	728	LYS	-	expression tag	UNP A0A1I5R890
С	1	MET	-	initiating methionine	UNP A0A1I5R890
С	2	GLY	-	expression tag	UNP A0A1I5R890
С	3	ASP	-	expression tag	UNP A0A1I5R890
С	4	ARG	-	expression tag	UNP A0A1I5R890



Continued from previous page...

Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
С	5	GLY	-	expression tag	UNP A0A1I5R890
С	6	PRO	-	expression tag	UNP A0A1I5R890
С	7	GLU	-	expression tag	UNP A0A1I5R890
С	8	PHE	-	expression tag	UNP A0A1I5R890
С	708	LEU	-	expression tag	UNP A0A1I5R890
С	709	GLU	-	expression tag	UNP A0A1I5R890
С	710	VAL	-	expression tag	UNP A0A1I5R890
С	711	ASP	-	expression tag	UNP A0A1I5R890
С	712	LEU	-	expression tag	UNP A0A1I5R890
С	713	GLN	-	expression tag	UNP A0A1I5R890
С	714	GLY	-	expression tag	UNP A0A1I5R890
С	715	ASP	-	expression tag	UNP A0A1I5R890
С	716	HIS	-	expression tag	UNP A0A1I5R890
С	717	GLY	-	expression tag	UNP A0A1I5R890
С	718	LEU	-	expression tag	UNP A0A1I5R890
С	719	SER	-	expression tag	UNP A0A1I5R890
С	720	ALA	-	expression tag	UNP A0A1I5R890
С	721	TRP	-	expression tag	UNP A0A1I5R890
С	722	SER	-	expression tag	UNP A0A1I5R890
С	723	HIS	-	expression tag	UNP A0A1I5R890
С	724	PRO	-	expression tag	UNP A0A1I5R890
С	725	GLN	-	expression tag	UNP A0A1I5R890
С	726	PHE	_	expression tag	UNP A0A1I5R890
С	727	GLU	-	expression tag	UNP A0A1I5R890
С	728	LYS	-	expression tag	UNP A0A1I5R890
D	1	MET	_	initiating methionine	UNP A0A1I5R890
D	2	GLY	-	expression tag	UNP A0A1I5R890
D	3	ASP	-	expression tag	UNP A0A1I5R890
D	4	ARG	_	expression tag	UNP A0A1I5R890
D	5	GLY	-	expression tag	UNP A0A1I5R890
D	6	PRO	_	expression tag	UNP A0A1I5R890
D	7	GLU	_	expression tag	UNP A0A1I5R890
D	8	PHE	-	expression tag	UNP A0A1I5R890
D	708	LEU	-	expression tag	UNP A0A1I5R890
D	709	GLU	-	expression tag	UNP A0A1I5R890
D	710	VAL	-	expression tag	UNP A0A1I5R890
D	711	ASP	-	expression tag	UNP A0A1I5R890
D	712	LEU	-	expression tag	UNP A0A1I5R890
D	713	GLN	-	expression tag	UNP A0A1I5R890
D	714	GLY	-	expression tag	UNP A0A1I5R890
D	715	ASP	-	expression tag	UNP A0A1I5R890
D	716	HIS	-	expression tag	UNP A0A1I5R890



Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	717	GLY	-	expression tag	UNP A0A1I5R890
D	718	LEU	-	expression tag	UNP A0A1I5R890
D	719	SER	-	expression tag	UNP A0A1I5R890
D	720	ALA	-	expression tag	UNP A0A1I5R890
D	721	TRP	-	expression tag	UNP A0A1I5R890
D	722	SER	-	expression tag	UNP A0A1I5R890
D	723	HIS	-	expression tag	UNP A0A1I5R890
D	724	PRO	-	expression tag	UNP A0A1I5R890
D	725	GLN	-	expression tag	UNP A0A1I5R890
D	726	PHE	-	expression tag	UNP A0A1I5R890
D	727	GLU	-	expression tag	UNP A0A1I5R890
D	728	LYS	-	expression tag	UNP A0A1I5R890

• Molecule 2 is COBALT (II) ION (three-letter code: CO) (formula: Co) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Co 1 1	0	0
2	В	1	Total Co 1 1	0	0
2	С	1	Total Co 1 1	0	0
2	D	1	Total Co 1 1	0	0


• Molecule 3 is HISTIDINE (three-letter code: HIS) (formula:  $C_6H_{10}N_3O_2$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
3	Δ	1	Total C N O	0	0	
	Λ	1	11 6 3 2	U		
3	В	1	Total C N O	0	0	
3	D	1	11 6 3 2			
3	C	1	Total C N O	0	0	
3	C	1	11 6 3 2	0		
2	D	1	Total C N O	0	0	
3	D	$D \mid I \mid$	11 6 3 2	0	. 0	


 $\bullet$  Molecule 4 is CYSTEINE (three-letter code: CYS) (formula:  $C_3H_7NO_2S)$  (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
4	Λ	1	Total	С	N	О	S	0	0
4	A	1	7	3	1	2	1		0
1	R	1	Total	С	N	О	S	0	0
4	Б	1	7	3	1	2	1		U
1	С	1	Total	С	N	О	S	0	0
4		1	7	3	1	2	1	U	U
1	D	1	Total	С	N	О	S	0	0
4	D	1	7	3	1	2	1	0	0

• Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total C O 7 4 3	0	0
5	С	1	Total C O 7 4 3	0	0

### • Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	49	Total O 49 49	0	0
6	В	40	Total O 40 40	0	0
6	С	44	Total O 44 44	0	0
6	D	14	Total O 14 14	0	0




### 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 5-histidylcysteine sulfoxide synthase/putative 4-mercaptohistidine N1-methyltranferase




• Molecule 1: 5-histidylcysteine sulfoxide synthase/putative 4-mercaptohistidine N1-methyltranferase




• Molecule 1: 5-histidylcysteine sulfoxide synthase/putative 4-mercaptohistidine N1-methyltranferase





 $\bullet \ Molecule \ 1: \ 5-histidyl cysteine \ sulfoxide \ synthase/putative \ 4-mercapto histidine \ N1-methyl transferase$ 





### 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	72.26Å 119.11Å 412.43Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	38.70 - 2.69	Depositor
resolution (A)	39.58 - 2.69	EDS
% Data completeness	99.6 (38.70-2.69)	Depositor
(in resolution range)	99.8 (39.58-2.69)	EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.07  (at  2.69Å)	Xtriage
Refinement program	PHENIX (1.18.2_3874: ???)	Depositor
P.P.	0.201 , $0.243$	Depositor
$R, R_{free}$	0.201 , $0.242$	DCC
$R_{free}$ test set	4983 reflections $(5.00\%)$	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	58.1	Xtriage
Anisotropy	0.313	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.32 , 29.8	EDS
L-test for twinning <sup>2</sup>	$ < L > = 0.49, < L^2> = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	23253	wwPDB-VP
Average B, all atoms $(Å^2)$	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 12.39% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

### 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, CO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.26	0/5953	0.42	0/8066	
1	В	0.25	0/5932	0.42	0/8039	
1	С	0.25	0/5945	0.42	0/8054	
1	D	0.24	0/5858	0.41	0/7943	
All	All	0.25	0/23688	0.41	0/32102	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	5784	0	5616	32	0
1	В	5763	0	5591	35	0
1	С	5776	0	5609	37	0
1	D	5693	0	5488	43	0
2	A	1	0	0	0	0
2	В	1	0	0	0	0
2	С	1	0	0	0	0
2	D	1	0	0	0	0
3	A	11	0	6	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	11	0	6	0	0
3	С	11	0	6	0	0
3	D	11	0	6	0	0
4	A	7	0	3	0	0
4	В	7	0	3	0	0
4	С	7	0	3	0	0
4	D	7	0	3	0	0
5	В	7	0	10	0	0
5	С	7	0	10	0	0
6	A	49	0	0	0	0
6	В	40	0	0	0	0
6	С	44	0	0	0	0
6	D	14	0	0	0	0
All	All	23253	0	22360	146	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 146 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} & & & & & & & & & & & & & & & & & & &$	Clash overlap (Å)	
1:B:549:ARG:HG2	1:B:580:VAL:HG13	1.69	0.73	
1:A:108:ASP:HB3	1:B:624:LYS:HD3	1.77	0.67	
1:C:105:MET:O	1:C:444:HIS:NE2	2.23	0.67	
1:A:240:SER:O	1:A:452:ARG:NH1	2.29	0.66	
1:C:549:ARG:HG2	1:C:580:VAL:HG13	1.79	0.64	

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	$696/728 \; (96\%)$	676 (97%)	20 (3%)	0	100 100
1	В	694/728 (95%)	670 (96%)	24 (4%)	0	100 100
1	С	695/728~(96%)	672 (97%)	23 (3%)	0	100 100
1	D	693/728 (95%)	661 (95%)	32 (5%)	0	100 100
All	All	2778/2912 (95%)	2679 (96%)	99 (4%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Rotameric		Percent	iles
1	A	623/651~(96%)	619 (99%)	4 (1%)	86	95
1	В	620/651~(95%)	616 (99%)	4 (1%)	86	95
1	С	622/651~(96%)	617 (99%)	5 (1%)	81	93
1	D	604/651~(93%)	602 (100%)	2 (0%)	92	86
All	All	$2469/2604\ (95\%)$	2454 (99%)	15 (1%)	86	95

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	406	TYR
1	D	179	ILE
1	С	38	PHE
1	D	406	TYR
1	С	406	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 14 ligands modelled in this entry, 4 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res	Link	В	ond leng	$\operatorname{gths}$	В	ond ang	gles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
4	CYS	С	803	-	5,6,6	1.04	1 (20%)	5,7,7	1.58	2 (40%)
3	HIS	A	802	2	6,11,11	0.98	1 (16%)	7,14,14	1.59	3 (42%)
5	PEG	С	804	-	6,6,6	0.49	0	5,5,5	0.24	0
3	HIS	С	802	2	6,11,11	0.98	1 (16%)	7,14,14	1.53	3 (42%)
4	CYS	A	803	-	5,6,6	1.09	1 (20%)	5,7,7	1.68	2 (40%)
5	PEG	В	804	-	6,6,6	0.49	0	5,5,5	0.23	0
4	CYS	D	803	-	5,6,6	1.03	1 (20%)	5,7,7	1.57	2 (40%)
4	CYS	В	803	-	5,6,6	1.00	1 (20%)	5,7,7	1.68	2 (40%)
3	HIS	D	802	2	6,11,11	0.99	1 (16%)	7,14,14	1.65	3 (42%)
3	HIS	В	802	2	6,11,11	0.95	1 (16%)	7,14,14	1.58	3 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CYS	С	803	-	-	3/6/6/6	-
3	HIS	A	802	2	-	1/8/8/8	0/1/1/1
5	PEG	С	804	-	-	0/4/4/4	-



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HIS	С	802	2	-	1/8/8/8	0/1/1/1
4	CYS	A	803	-	-	3/6/6/6	-
5	PEG	В	804	-	-	1/4/4/4	-
4	CYS	D	803	-	-	3/6/6/6	-
4	CYS	В	803	-	-	3/6/6/6	-
3	HIS	D	802	2	-	4/8/8/8	0/1/1/1
3	HIS	В	802	2	-	2/8/8/8	0/1/1/1

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	Observed(A)	$\operatorname{Ideal}( ext{\AA})$
3	D	802	HIS	OXT-C	-2.16	1.23	1.30
3	A	802	HIS	OXT-C	-2.15	1.23	1.30
3	С	802	HIS	OXT-C	-2.14	1.23	1.30
4	D	803	CYS	OXT-C	-2.14	1.23	1.30
4	С	803	CYS	OXT-C	-2.14	1.23	1.30

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}(^{o})$
3	D	802	HIS	OXT-C-O	-2.95	117.39	124.09
3	A	802	HIS	OXT-C-O	-2.81	117.71	124.09
4	С	803	CYS	OXT-C-O	-2.74	117.87	124.09
4	A	803	CYS	OXT-C-O	-2.70	117.95	124.09
3	В	802	HIS	OXT-C-O	-2.64	118.09	124.09

There are no chirality outliers.

5 of 21 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	В	802	HIS	CA-CB-CG-ND1
4	A	803	CYS	O-C-CA-N
4	В	803	CYS	O-C-CA-N
4	В	803	CYS	OXT-C-CA-N
4	С	803	CYS	OXT-C-CA-N


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In




addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
































### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



### 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$OWAB(A^2)$	Q < 0.9
1	A	700/728 (96%)	0.07	16 (2%) 60 62	38, 55, 90, 152	0
1	В	698/728 (95%)	0.07	19 (2%) 54 55	39, 58, 86, 162	0
1	С	699/728 (96%)	0.03	18 (2%) 56 57	40, 57, 86, 133	0
1	D	697/728 (95%)	0.41	47 (6%) 17 16	45, 83, 122, 169	0
All	All	2794/2912 (95%)	0.14	100 (3%) 42 42	38, 60, 109, 169	0

The worst 5 of 100 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	440	ALA	4.1
1	D	286	PHE	3.8
1	D	117	TRP	3.8
1	D	116	GLU	3.6
1	В	630	GLY	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
5	PEG	В	804	7/7	0.65	0.24	81,92,95,96	0
5	PEG	С	804	7/7	0.75	0.18	61,70,80,84	0
4	CYS	D	803	7/7	0.91	0.25	93,104,108,109	0
4	CYS	В	803	7/7	0.92	0.20	72,72,73,75	0
3	HIS	D	802	11/11	0.92	0.36	92,97,100,100	0
2	CO	D	801	1/1	0.93	0.10	89,89,89,89	0
3	HIS	A	802	11/11	0.93	0.23	53,57,63,64	0
4	CYS	С	803	7/7	0.93	0.22	72,72,72,73	0
2	CO	В	801	1/1	0.94	0.15	55,55,55,55	0
3	HIS	В	802	11/11	0.94	0.30	71,72,72,72	0
3	HIS	С	802	11/11	0.94	0.32	72,72,72,72	0
2	CO	A	801	1/1	0.95	0.12	57,57,57,57	0
4	CYS	A	803	7/7	0.96	0.16	61,70,75,76	0
2	CO	С	801	1/1	0.97	0.17	61,61,61,61	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






## Electron density around CYS B 803: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)



### 








# Electron density around HIS A 802: 2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)








## Electron density around CO B 801: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)








## Electron density around CO A 801: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)



## Electron density around CYS A 803: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





### 6.5 Other polymers (i)

There are no such residues in this entry.

