



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 16, 2023 – 02:40 AM JST

PDB ID : 6KLR
Title : Crystal structure of human WIPI3 in complex with the WIR-peptide from ATG2A
Authors : Ren, J.Q.; Liang, R.B.; Feng, W.
Deposited on : 2019-07-30
Resolution : 2.21 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5912 (2.24-2.20)
Clashscore	141614	6646 (2.24-2.20)
Ramachandran outliers	138981	6543 (2.24-2.20)
Sidechain outliers	138945	6544 (2.24-2.20)
RSRZ outliers	127900	5797 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	350	
1	B	350	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2485 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called chimera ATG2A and WIPI3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	297	2303	1465	395	425	18	0	2	0
1	B	22	164	108	28	28		0	0	0

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-32	GLY	-	expression tag	UNP Q2TAZO
A	-31	PRO	-	expression tag	UNP Q2TAZO
A	-30	GLY	-	expression tag	UNP Q2TAZO
A	-29	SER	-	expression tag	UNP Q2TAZO
A	3	GLY	-	linker	UNP Q2TAZO
A	4	SER	-	linker	UNP Q2TAZO
A	?	-	PRO	deletion	UNP Q5MNZ6
A	?	-	LYS	deletion	UNP Q5MNZ6
A	?	-	TYR	deletion	UNP Q5MNZ6
A	?	-	PRO	deletion	UNP Q5MNZ6
A	?	-	PRO	deletion	UNP Q5MNZ6
A	?	-	ASN	deletion	UNP Q5MNZ6
A	?	-	ARG	deletion	UNP Q5MNZ6
A	?	-	ASN	deletion	UNP Q5MNZ6
A	?	-	LYS	deletion	UNP Q5MNZ6
A	?	-	GLN	deletion	UNP Q5MNZ6
A	?	-	SER	deletion	UNP Q5MNZ6
A	?	-	SER	deletion	UNP Q5MNZ6
A	?	-	LEU	deletion	UNP Q5MNZ6
A	?	-	ALA	deletion	UNP Q5MNZ6
A	?	-	SER	deletion	UNP Q5MNZ6
A	?	-	ALA	deletion	UNP Q5MNZ6
A	?	-	SER	deletion	UNP Q5MNZ6
A	?	-	PHE	deletion	UNP Q5MNZ6
A	?	-	LEU	deletion	UNP Q5MNZ6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	PRO	deletion	UNP Q5MNZ6
A	?	-	LYS	deletion	UNP Q5MNZ6
A	?	-	TYR	deletion	UNP Q5MNZ6
A	?	-	PHE	deletion	UNP Q5MNZ6
A	?	-	SER	deletion	UNP Q5MNZ6
B	1370	GLY	-	expression tag	UNP Q2TAZ0
B	1371	PRO	-	expression tag	UNP Q2TAZ0
B	1372	GLY	-	expression tag	UNP Q2TAZ0
B	1373	SER	-	expression tag	UNP Q2TAZ0
B	1405	GLY	-	linker	UNP Q2TAZ0
B	1406	SER	-	linker	UNP Q2TAZ0
B	?	-	PRO	deletion	UNP Q5MNZ6
B	?	-	LYS	deletion	UNP Q5MNZ6
B	?	-	TYR	deletion	UNP Q5MNZ6
B	?	-	PRO	deletion	UNP Q5MNZ6
B	?	-	PRO	deletion	UNP Q5MNZ6
B	?	-	ASN	deletion	UNP Q5MNZ6
B	?	-	ARG	deletion	UNP Q5MNZ6
B	?	-	ASN	deletion	UNP Q5MNZ6
B	?	-	LYS	deletion	UNP Q5MNZ6
B	?	-	GLN	deletion	UNP Q5MNZ6
B	?	-	SER	deletion	UNP Q5MNZ6
B	?	-	SER	deletion	UNP Q5MNZ6
B	?	-	LEU	deletion	UNP Q5MNZ6
B	?	-	ALA	deletion	UNP Q5MNZ6
B	?	-	SER	deletion	UNP Q5MNZ6
B	?	-	ALA	deletion	UNP Q5MNZ6
B	?	-	SER	deletion	UNP Q5MNZ6
B	?	-	PHE	deletion	UNP Q5MNZ6
B	?	-	LEU	deletion	UNP Q5MNZ6
B	?	-	PRO	deletion	UNP Q5MNZ6
B	?	-	LYS	deletion	UNP Q5MNZ6
B	?	-	TYR	deletion	UNP Q5MNZ6
B	?	-	PHE	deletion	UNP Q5MNZ6
B	?	-	SER	deletion	UNP Q5MNZ6

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	16	Total O 16 16	0	0
2	B	2	Total O 2 2	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	75.12Å 121.06Å 37.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.52 – 2.21 24.52 – 2.21	Depositor EDS
% Data completeness (in resolution range)	98.4 (24.52-2.21) 98.4 (24.52-2.21)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.86 (at 2.22Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.213 , 0.255 0.217 , 0.257	Depositor DCC
R_{free} test set	864 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	54.4	Xtrriage
Anisotropy	0.673	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 56.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2485	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/2355	0.58	0/3186
1	B	0.38	0/170	0.55	0/235
All	All	0.40	0/2525	0.58	0/3421

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2303	0	2262	10	0
1	B	164	0	157	1	0
2	A	16	0	0	0	0
2	B	2	0	0	0	0
All	All	2485	0	2419	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 11 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:84:VAL:HG23	1:A:85:ILE:HG13	1.66	0.77

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:HIS:ND1	1:A:173:PRO:HA	2.10	0.66
1:A:156:HIS:CE1	1:A:173:PRO:HB3	2.35	0.62
1:A:156:HIS:HE1	1:A:173:PRO:HB3	1.68	0.57
1:A:295:PHE:HA	1:A:300:GLU:O	2.09	0.53

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	295/350 (84%)	285 (97%)	10 (3%)	0	100	100
1	B	20/350 (6%)	19 (95%)	1 (5%)	0	100	100
All	All	315/700 (45%)	304 (96%)	11 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	250/298 (84%)	246 (98%)	4 (2%)	62	75
1	B	18/298 (6%)	18 (100%)	0	100	100
All	All	268/596 (45%)	264 (98%)	4 (2%)	65	76

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	CYS
1	A	246	HIS
1	A	247	ILE
1	A	259	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	297/350 (84%)	0.42	20 (6%) 17 16	48, 79, 134, 168	0
1	B	22/350 (6%)	0.48	4 (18%) 1 1	70, 93, 108, 143	0
All	All	319/700 (45%)	0.43	24 (7%) 14 12	48, 81, 134, 168	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	255	SER	8.0
1	A	263	GLN	4.0
1	A	221	ALA	3.7
1	A	222	ALA	3.7
1	A	253	PRO	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.