



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 23, 2024 – 08:42 AM EDT

PDB ID : 3KXA  
Title : Crystal Structure of NGO0477 from Neisseria gonorrhoeae  
Authors : Ren, J.; Sainsbury, S.; Nettleship, J.E.; Owens, R.J.; Oxford Protein Production Facility (OPPF)  
Deposited on : 2009-12-02  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 1.20.1  
EDS : **FAILED**  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	4123 (2.80-2.80)
Ramachandran outliers	177936	4071 (2.80-2.80)
Sidechain outliers	177891	4073 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	A	141	
1	B	141	
1	C	141	
1	D	141	

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4160 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative uncharacterized protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	131	1020	639	193	183	2	3	0	0	0
1	B	130	1011	634	192	180	2	3	0	0	0
1	C	131	1020	639	193	183	2	3	0	0	0
1	D	130	1011	634	192	180	2	3	0	0	0

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MSE	-	expression tag	UNP Q5F9C2
A	-6	ALA	-	expression tag	UNP Q5F9C2
A	-5	HIS	-	expression tag	UNP Q5F9C2
A	-4	HIS	-	expression tag	UNP Q5F9C2
A	-3	HIS	-	expression tag	UNP Q5F9C2
A	-2	HIS	-	expression tag	UNP Q5F9C2
A	-1	HIS	-	expression tag	UNP Q5F9C2
A	0	HIS	-	expression tag	UNP Q5F9C2
B	-7	MSE	-	expression tag	UNP Q5F9C2
B	-6	ALA	-	expression tag	UNP Q5F9C2
B	-5	HIS	-	expression tag	UNP Q5F9C2
B	-4	HIS	-	expression tag	UNP Q5F9C2
B	-3	HIS	-	expression tag	UNP Q5F9C2
B	-2	HIS	-	expression tag	UNP Q5F9C2
B	-1	HIS	-	expression tag	UNP Q5F9C2
B	0	HIS	-	expression tag	UNP Q5F9C2
C	-7	MSE	-	expression tag	UNP Q5F9C2
C	-6	ALA	-	expression tag	UNP Q5F9C2
C	-5	HIS	-	expression tag	UNP Q5F9C2
C	-4	HIS	-	expression tag	UNP Q5F9C2
C	-3	HIS	-	expression tag	UNP Q5F9C2

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	HIS	-	expression tag	UNP Q5F9C2
C	-1	HIS	-	expression tag	UNP Q5F9C2
C	0	HIS	-	expression tag	UNP Q5F9C2
D	-7	MSE	-	expression tag	UNP Q5F9C2
D	-6	ALA	-	expression tag	UNP Q5F9C2
D	-5	HIS	-	expression tag	UNP Q5F9C2
D	-4	HIS	-	expression tag	UNP Q5F9C2
D	-3	HIS	-	expression tag	UNP Q5F9C2
D	-2	HIS	-	expression tag	UNP Q5F9C2
D	-1	HIS	-	expression tag	UNP Q5F9C2
D	0	HIS	-	expression tag	UNP Q5F9C2

- Molecule 2 is ASPARAGINE (three-letter code: ASN) (formula:  $C_4H_8N_2O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
2	A	1	Total	C	N	O	0	0
			9	4	2	3		
2	B	1	Total	C	N	O	0	0
			9	4	2	3		
2	C	1	Total	C	N	O	0	0
			9	4	2	3		
2	D	1	Total	C	N	O	0	0
			9	4	2	3		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	5	Total Cl 5 5	0	0
3	B	2	Total Cl 2 2	0	0
3	C	6	Total Cl 6 6	0	0

- Molecule 4 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	11	Total X 11 11	0	0
4	C	11	Total X 11 11	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	8	Total O 8 8	0	0
5	B	9	Total O 9 9	0	0
5	C	5	Total O 5 5	0	0
5	D	5	Total O 5 5	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.


- Molecule 1: Putative uncharacterized protein

Chain A: 



- Molecule 1: Putative uncharacterized protein

Chain B: 




- Molecule 1: Putative uncharacterized protein

Chain C: 



- Molecule 1: Putative uncharacterized protein

Chain D: 



M130
GLU
GLN
ALA

## 4 Data and refinement statistics i

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	124.82Å 124.82Å 137.58Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	26.31 – 2.80	Depositor
% Data completeness (in resolution range)	99.4 (26.31-2.80)	Depositor
$R_{merge}$	0.24	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.21 (at 2.80Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.230 , 0.280	Depositor
Wilson B-factor (Å <sup>2</sup> )	43.9	Xtrriage
Anisotropy	0.560	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.469 for h,-h-k,-l	Xtrriage
Total number of atoms	4160	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UNX, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/1033	0.99	6/1385 (0.4%)
1	B	0.41	0/1024	0.74	5/1373 (0.4%)
1	C	0.46	0/1033	0.75	4/1385 (0.3%)
1	D	0.41	0/1024	0.74	5/1373 (0.4%)
All	All	0.44	0/4114	0.81	20/5516 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	41	ARG	NE-CZ-NH2	17.02	128.81	120.30
1	A	41	ARG	NE-CZ-NH1	-16.58	112.01	120.30
1	A	36	ARG	NE-CZ-NH2	-11.17	114.71	120.30
1	C	36	ARG	NE-CZ-NH2	-10.88	114.86	120.30
1	A	36	ARG	NE-CZ-NH1	10.76	125.68	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	22	SER	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1020	0	1070	42	0
1	B	1011	0	1064	37	0
1	C	1020	0	1070	39	0
1	D	1011	0	1064	29	0
2	A	9	0	5	1	0
2	B	9	0	5	0	0
2	C	9	0	5	2	0
2	D	9	0	5	0	0
3	A	5	0	0	0	0
3	B	2	0	0	0	0
3	C	6	0	0	0	0
4	A	11	0	0	0	0
4	C	11	0	0	0	0
5	A	8	0	0	1	0
5	B	9	0	0	0	0
5	C	5	0	0	1	0
5	D	5	0	0	0	0
All	All	4160	0	4288	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 105 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:47:ARG:HG3	1:B:47:ARG:HH11	1.07	1.16
1:D:47:ARG:HH11	1:D:47:ARG:HG3	1.05	1.15
1:C:47:ARG:HH11	1:C:47:ARG:HG3	1.04	1.12
1:A:47:ARG:HH11	1:A:47:ARG:HG3	1.03	1.10
1:C:21:ILE:HG21	1:D:60:ALA:HB2	1.53	0.91

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	129/141 (92%)	124 (96%)	3 (2%)	2 (2%)	8	27
1	B	128/141 (91%)	123 (96%)	4 (3%)	1 (1%)	16	44
1	C	129/141 (92%)	123 (95%)	5 (4%)	1 (1%)	16	44
1	D	128/141 (91%)	123 (96%)	3 (2%)	2 (2%)	8	27
All	All	514/564 (91%)	493 (96%)	15 (3%)	6 (1%)	11	34

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	23	PRO
1	C	23	PRO
1	D	2	PRO
1	A	2	PRO
1	D	23	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	108/112 (96%)	104 (96%)	4 (4%)	29	63
1	B	107/112 (96%)	103 (96%)	4 (4%)	29	63
1	C	108/112 (96%)	103 (95%)	5 (5%)	23	55
1	D	107/112 (96%)	103 (96%)	4 (4%)	29	63
All	All	430/448 (96%)	413 (96%)	17 (4%)	27	60

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	40	ARG
1	D	125	ARG
1	B	125	ARG
1	C	21	ILE
1	C	25	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	98	GLN
1	D	5	HIS
1	C	5	HIS
1	B	98	GLN
1	C	98	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 39 ligands modelled in this entry, 13 are monoatomic and 22 are unknown - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ASN	C	140	-	7,8,8	0.85	1 (14%)	6,10,10	1.36	1 (16%)
2	ASN	D	140	-	7,8,8	1.07	1 (14%)	6,10,10	1.25	1 (16%)
2	ASN	B	140	-	7,8,8	1.05	1 (14%)	6,10,10	1.25	1 (16%)
2	ASN	A	140	-	7,8,8	0.86	1 (14%)	6,10,10	1.33	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ASN	C	140	-	-	4/8/8/8	-
2	ASN	D	140	-	-	6/8/8/8	-
2	ASN	B	140	-	-	6/8/8/8	-
2	ASN	A	140	-	-	4/8/8/8	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	140	ASN	OXT-C	-2.26	1.23	1.30
2	C	140	ASN	OXT-C	-2.16	1.23	1.30
2	A	140	ASN	OXT-C	-2.14	1.23	1.30
2	D	140	ASN	OXT-C	-2.13	1.23	1.30

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	140	ASN	OXT-C-O	-3.18	116.86	124.08
2	A	140	ASN	OXT-C-O	-3.17	116.88	124.08
2	D	140	ASN	OXT-C-O	-2.97	117.33	124.08
2	B	140	ASN	OXT-C-O	-2.96	117.38	124.08

There are no chirality outliers.

5 of 20 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	140	ASN	O-C-CA-N
2	C	140	ASN	O-C-CA-N
2	D	140	ASN	O-C-CA-N

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	C	140	ASN	OXT-C-CA-N
2	D	140	ASN	OXT-C-CA-N

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	140	ASN	2	0
2	A	140	ASN	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

### 6.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

### 6.4 Ligands

EDS failed to run properly - this section is therefore empty.

### 6.5 Other polymers

EDS failed to run properly - this section is therefore empty.