



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 12, 2024 – 12:38 PM EST

PDB ID : 3L35
Title : PIE12 D-peptide against HIV entry
Authors : Welch, B.D.; Redman, J.S.; Paul, S.; Whitby, F.G.; Weinstock, M.T.; Reeves, J.D.; Lie, Y.S.; Eckert, D.M.; Hill, C.P.; Root, M.J.; Kay, M.S.
Deposited on : 2009-12-16
Resolution : 1.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1935 (1.56-1.56)
Clashscore	180529	2073 (1.56-1.56)
Ramachandran outliers	177936	2037 (1.56-1.56)
Sidechain outliers	177891	2034 (1.56-1.56)
RSRZ outliers	164620	1935 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	47	 4% 79% 21%
1	B	47	 15% 70% 28% .
1	C	47	 6% 70% 26% .
2	H	18	 78% 22%
2	K	18	 72% 28%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	L	18	 83% 6% 11%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1789 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GP41 N-PEPTIDE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	47	Total 384	C 245	N 70	O 68	S 1	0	0	1
1	B	47	Total 404	C 256	N 75	O 72	S 1	0	2	1
1	C	47	Total 384	C 245	N 70	O 68	S 1	0	0	1

- Molecule 2 is a protein called HIV ENTRY INHIBITOR PIE12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	18	Total 144	C 94	N 23	O 25	S 2	0	0	1
2	K	18	Total 144	C 94	N 23	O 25	S 2	0	0	1
2	L	16	Total 132	C 86	N 21	O 23	S 2	0	0	1



- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	61	Total 61	O 61	0	0
3	B	45	Total 45	O 45	0	0
3	C	39	Total 39	O 39	0	0
3	H	20	Total 20	O 20	0	0
3	K	20	Total 20	O 20	0	0
3	L	12	Total 12	O 12	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GP41 N-PEPTIDE





- Molecule 1: GP41 N-PEPTIDE



- Molecule 1: GP41 N-PEPTIDE





- Molecule 2: HIV ENTRY INHIBITOR PIE12



- Molecule 2: HIV ENTRY INHIBITOR PIE12



- Molecule 2: HIV ENTRY INHIBITOR PIE12



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	30.85Å 40.49Å 80.31Å 90.00° 91.76° 90.00°	Depositor
Resolution (Å)	30.00 – 1.55 30.00 – 1.55	Depositor EDS
% Data completeness (in resolution range)	86.4 (30.00-1.55) 86.4 (30.00-1.55)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.84 (at 1.55Å)	Xtrriage
Refinement program	REFMAC 5.5.0062	Depositor
R, R_{free}	0.233 , 0.288 0.228 , 0.283	Depositor DCC
R_{free} test set	1273 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	23.0	Xtrriage
Anisotropy	0.047	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 33.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.022 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	1789	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DHI, DCY, DLY, NH2, ACE, DPR, DGL, DTR, DGN, DLE, DTY, DAS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/382	0.57	0/506
1	B	0.51	0/402	0.71	0/532
1	C	0.55	0/382	0.67	0/506
2	H	0.56	0/3	0.34	0/2
2	K	0.58	0/3	0.94	0/2
2	L	2.14	0/3	0.92	0/2
All	All	0.55	0/1175	0.65	0/1550

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	K	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	K	15	DGL	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	384	0	428	11	0
1	B	404	0	445	19	0
1	C	384	0	428	16	0
2	H	144	0	113	2	0
2	K	144	0	112	3	0
2	L	132	0	97	1	0
3	A	61	0	0	2	0
3	B	45	0	0	3	0
3	C	39	0	0	4	0
3	H	20	0	0	0	0
3	K	20	0	0	0	0
3	L	12	0	0	0	0
All	All	1789	0	1623	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 35 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:ILE:HG21	1:B:43[A]:ARG:HD2	1.19	1.11
1:C:28:LYS:HE3	3:C:132:HOH:O	1.54	1.03
1:C:22:GLU:HG3	3:C:50:HOH:O	1.69	0.93
1:A:44:ILE:CG2	1:B:43[A]:ARG:HD2	2.04	0.87
1:B:22[B]:GLU:HA	1:B:22[B]:GLU:OE1	1.84	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	45/47 (96%)	45 (100%)	0	0	100	100
1	B	47/47 (100%)	47 (100%)	0	0	100	100
1	C	45/47 (96%)	45 (100%)	0	0	100	100
2	H	1/18 (6%)	1 (100%)	0	0	100	100
2	K	1/18 (6%)	1 (100%)	0	0	100	100
All	All	139/177 (78%)	139 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	42/42 (100%)	42 (100%)	0	100	100
1	B	44/42 (105%)	40 (91%)	4 (9%)	7	0
1	C	42/42 (100%)	39 (93%)	3 (7%)	12	1
All	All	128/126 (102%)	121 (94%)	7 (6%)	21	2

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	45	LEU
1	C	1	ARG
1	C	43	ARG
1	C	17	LYS
1	B	43[B]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

44 non-standard protein/DNA/RNA residues are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	45/47 (95%)	0.46	2 (4%) 39 47	13, 22, 32, 36	0
1	B	45/47 (95%)	0.78	7 (15%) 6 7	9, 24, 35, 36	2 (4%)
1	C	45/47 (95%)	0.74	3 (6%) 25 29	12, 22, 41, 47	0
2	H	1/18 (5%)	0.52	0 100 100	23, 23, 23, 23	0
2	K	1/18 (5%)	0.37	0 100 100	20, 20, 20, 20	0
2	L	1/18 (5%)	0.26	0 100 100	20, 20, 20, 20	0
All	All	138/195 (70%)	0.65	12 (8%) 17 20	9, 23, 35, 47	2 (1%)

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2	MET	3.1
1	C	45	LEU	2.8
1	C	4	GLN	2.7
1	A	11	GLU	2.5
1	A	17	LYS	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	DGL	L	15	9/10	0.80	0.14	24,26,38,39	0
2	DGL	H	15	9/10	0.85	0.12	24,26,38,38	0
2	DLE	H	16	8/9	0.86	0.13	21,22,24,25	0
2	DCY	H	5	6/7	0.88	0.12	23,25,26,31	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	DGL	K	15	9/10	0.88	0.13	22,24,28,29	0
2	DAS	K	6	8/9	0.90	0.08	18,19,20,25	0
2	DGN	H	11	9/10	0.90	0.12	19,20,28,29	0
2	DHI	K	3	10/11	0.90	0.09	17,18,22,24	0
2	DLY	K	1	9/10	0.91	0.12	18,21,35,38	0
2	DPR	H	8	7/8	0.91	0.09	20,21,21,22	0
2	DCY	K	5	6/7	0.91	0.10	17,19,20,24	0
2	DTR	H	12	14/15	0.91	0.08	15,17,19,20	0
2	DLE	H	13	8/9	0.92	0.08	16,17,19,20	0
2	DCY	K	14	6/7	0.92	0.09	19,21,22,25	0
2	DGL	H	9	9/10	0.92	0.09	18,18,24,25	0
2	DGL	K	9	9/10	0.92	0.08	15,17,20,23	0
2	DHI	H	3	10/11	0.92	0.08	21,22,27,27	0
2	DAS	H	6	8/9	0.92	0.08	23,24,26,28	0
2	DLE	L	16	8/9	0.92	0.10	23,24,26,26	0
2	DTY	L	7	12/13	0.93	0.09	18,21,29,34	0
2	DGN	L	11	9/10	0.93	0.10	17,19,31,33	0
2	DLY	H	1	9/10	0.93	0.10	22,23,29,31	0
2	DCY	L	5	6/7	0.93	0.08	19,20,22,24	0
2	DLE	K	16	8/9	0.93	0.09	23,24,25,26	0
2	DTY	H	7	12/13	0.93	0.08	18,21,22,23	0
2	DLE	L	13	8/9	0.94	0.08	17,18,19,21	0
2	DCY	H	14	6/7	0.94	0.09	21,23,24,29	0
2	DTR	K	12	14/15	0.94	0.07	15,18,19,20	0
2	DCY	L	14	6/7	0.94	0.10	20,22,23,26	0
2	DAS	L	6	8/9	0.94	0.09	20,21,25,26	0
2	DGN	K	11	9/10	0.95	0.08	17,18,21,27	0
2	DPR	L	8	7/8	0.95	0.06	15,16,16,16	0
2	DHI	L	3	10/11	0.95	0.07	18,19,22,23	0
2	DTY	K	7	12/13	0.95	0.07	14,17,20,21	0
2	DTR	L	12	14/15	0.95	0.06	16,18,21,22	0
2	DGL	L	9	9/10	0.95	0.07	14,15,18,22	0
2	DTR	H	10	14/15	0.95	0.06	16,17,18,19	0
2	DPR	K	8	7/8	0.95	0.08	17,18,18,20	0
2	DTR	L	10	14/15	0.96	0.05	12,14,16,16	0
2	DLE	K	13	8/9	0.96	0.06	13,16,17,18	0
2	DPR	L	4	7/8	0.97	0.07	18,19,19,20	0
2	DTR	K	10	14/15	0.97	0.06	13,15,17,17	0
2	DPR	H	4	7/8	0.97	0.06	20,21,21,21	0
2	DPR	K	4	7/8	0.98	0.05	16,17,17,17	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.