



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 12, 2024 – 02:40 PM EDT

PDB ID : 1L5S
Title : Human liver glycogen phosphorylase complexed with uric acid, N-Acetyl-beta-D-glucopyranosylamine, and CP-403,700
Authors : Ekstrom, J.L.; Pauly, T.A.; Carty, M.D.; Soeller, W.C.; Culp, J.; Danley, D.E.; Hoover, D.J.; Treadway, J.L.; Gibbs, E.M.; Fletterick, R.J.; Day, Y.S.N.; Myszka, D.G.; Rath, V.L.
Deposited on : 2002-03-07
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	847	
1	B	847	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	MRD	B	903	-	-	X	-

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 13826 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycogen phosphorylase, liver form.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	793	Total 6437	C 4134	N 1092	O 1182	S 29	0	0	0
1	B	791	Total 6420	C 4125	N 1090	O 1176	S 29	0	0	0

- Molecule 2 is N-acetyl-beta-D-glucopyranosylamine (three-letter code: NBG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total 15	C 8	N 1	O 6	0	0
2	B	1	Total 15	C 8	N 1	O 6	0	0

- Molecule 3 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	15	8	1	5	1	0	0
3	B	1	15	8	1	5	1	0	0

- Molecule 4 is [5-CHLORO-1H-INDOL-2-CARBONYL-PHENYLALANINYL]-AZETIDINE-3-CARBOXYLIC ACID (three-letter code: 700) (formula: C₂₂H₂₀ClN₃O₄).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Cl	N	O		
4	A	1	30	22	1	3	4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Cl	N			O
4	B	1	30	22	1	3	4	0	0

- Molecule 5 is URIC ACID (three-letter code: URC) (formula: $C_5H_4N_4O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	12	5	4	3	0	0
5	B	1	12	5	4	3	0	0

- Molecule 6 is (4R)-2-METHYLPENTANE-2,4-DIOL (three-letter code: MRD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 8 6 2	0	0
6	A	1	Total C O 8 6 2	0	0
6	A	1	Total C O 8 6 2	0	0
6	B	1	Total C O 8 6 2	0	0
6	B	1	Total C O 8 6 2	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	433	Total O 433 433	0	0
7	B	352	Total O 352 352	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: Glycogen phosphorylase, liver form

Chain A: 



- Molecule 1: Glycogen phosphorylase, liver form

Chain B: 



ASN	V379	L489	V630	R770	ASP	S830	ASP	S830	ASP
LEU	L380	L502	M631	F771	LEU	L802	LEU	L802	LEU
ARG	P361	I503	M632	K772	LYS	S808	LYS	S808	LYS
PHE	E382	I507	D633	A775	GLU	G889	GLU	G889	GLU
ASN	E385	V512	V636	D785	ASN	K810	ASN	K810	ASN
VAL	R386	D527	G637	K786	THR	R815	THR	R815	THR
GLY	V392	D528	S638	S788	ILE	K818	ILE	K818	ILE
D261	E393	V529	K639	Q789	LEU	W827	LEU	W827	LEU
A272	E273	L644	F644	S789	SER	E828	SER	E828	SER
N274	L275	L645	L646	M792	ASN	P829	ASN	P829	ASN
I276	S276	E646	L652	H793	GLY	S830	GLY	S830	GLY
R277	R277	L662	L662	P794	ASN	ASP	ASN	ASP	ASN
F286	F286	L666	L666	L802	LEU	LEU	LEU	LEU	LEU
Q296	Q296	I666	I666	L802	LYS	LYS	LYS	LYS	LYS
L304	L304	S674	S674	L802	ILE	ILE	ILE	ILE	ILE
K420	K420	N678	N678	L802	SER	SER	SER	SER	SER
I308	I308	M679	M679	L802	LEU	LEU	LEU	LEU	LEU
R309	R309	K680	K680	L802	ASN	ASN	ASN	ASN	ASN
K315	K315	F681	F681	L802	GLU	GLU	GLU	GLU	GLU
F316	F316	M682	M682	L802	SER	SER	SER	SER	SER
GLY	GLY	M699	M699	L802	ASN	ASN	ASN	ASN	ASN
SER	SER	D715	D715	L802	GLY	GLY	GLY	GLY	GLY
THR	THR	D716	D716	L802	ASN	ASN	ASN	ASN	ASN
ARG	ARG	A719	A719	L802	LEU	LEU	LEU	LEU	LEU
GLY	GLY	A728	A728	L802	LYS	LYS	LYS	LYS	LYS
ALA	ALA	K729	K729	L802	VAL	VAL	VAL	VAL	VAL
GLY	GLY	Y732	Y732	L802	GLY	GLY	GLY	GLY	GLY
THR	THR	E733	E733	L802	ASN	ASN	ASN	ASN	ASN
V325	V325	K739	K739	L802	ASN	ASN	ASN	ASN	ASN
F326	F326	D743	D743	L802	ASN	ASN	ASN	ASN	ASN
F329	F329	P765	P765	L802	ASN	ASN	ASN	ASN	ASN
P330	P330	K769	K769	L802	ASN	ASN	ASN	ASN	ASN
A334	A334	D760	D760	L802	ASN	ASN	ASN	ASN	ASN
L337	L337	I761	I761	L802	ASN	ASN	ASN	ASN	ASN
I346	I346	M764	M764	L802	ASN	ASN	ASN	ASN	ASN
L349	L349	L766	L766	L802	ASN	ASN	ASN	ASN	ASN
I352	I352	V629	V629	L802	ASN	ASN	ASN	ASN	ASN
W361	W361	Y613	Y613	L802	ASN	ASN	ASN	ASN	ASN
S362	S362	H614	H614	L802	ASN	ASN	ASN	ASN	ASN
K363	K363	M615	M615	L802	ASN	ASN	ASN	ASN	ASN
A364	A364	L622	L622	L802	ASN	ASN	ASN	ASN	ASN
W365	W365	D628	D628	L802	ASN	ASN	ASN	ASN	ASN
E366	E366	V629	V629	L802	ASN	ASN	ASN	ASN	ASN
Q369	Q369	V629	V629	L802	ASN	ASN	ASN	ASN	ASN

4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	124.03Å 124.03Å 123.30Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.60 – 2.10	Depositor
% Data completeness (in resolution range)	87.8 (40.60-2.10)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.194 , 0.236	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	13826	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: URC, 700, MRD, PLP, NBG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/6581	0.70	3/8900 (0.0%)
1	B	0.47	0/6564	0.69	2/8877 (0.0%)
All	All	0.47	0/13145	0.69	5/17777 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	490	ARG	NE-CZ-NH2	-7.68	116.46	120.30
1	A	490	ARG	NE-CZ-NH2	-5.49	117.55	120.30
1	B	129	ALA	N-CA-C	-5.32	96.62	111.00
1	A	129	ALA	N-CA-C	-5.21	96.94	111.00
1	A	602	THR	N-CA-C	-5.08	97.28	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6437	0	6418	149	0
1	B	6420	0	6408	153	0
2	A	15	0	15	0	0
2	B	15	0	15	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	15	0	7	0	0
3	B	15	0	7	1	0
4	A	30	0	18	0	0
4	B	30	0	18	0	0
5	A	12	0	4	0	0
5	B	12	0	4	0	0
6	A	24	0	42	3	0
6	B	16	0	28	9	0
7	A	433	0	0	16	0
7	B	352	0	0	21	0
All	All	13826	0	12984	307	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 307 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:109:ASP:HB3	7:A:2741:HOH:O	1.53	1.05
1:A:49:ARG:HH22	6:A:905:MRD:H1C2	1.26	1.01
1:A:754:GLN:HG2	1:A:757:LEU:HB2	1.43	1.00
1:B:818:LYS:NZ	1:B:818:LYS:HB3	1.81	0.95
1:A:547:GLN:O	1:A:551:THR:HG23	1.72	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	787/847 (93%)	743 (94%)	38 (5%)	6 (1%)	19 15
1	B	785/847 (93%)	747 (95%)	33 (4%)	5 (1%)	25 21

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1572/1694 (93%)	1490 (95%)	71 (4%)	11 (1%)	22 18

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	421	ASP
1	A	555	VAL
1	B	421	ASP
1	B	435	GLY
1	A	829	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	694/740 (94%)	662 (95%)	32 (5%)	27 26
1	B	692/740 (94%)	655 (95%)	37 (5%)	22 20
All	All	1386/1480 (94%)	1317 (95%)	69 (5%)	24 23

5 of 69 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	615	MET
1	B	628	ASP
1	B	755	PRO
1	A	652	LEU
1	A	645	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	822	GLN
1	B	826	ASN
1	B	44	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	96	GLN
1	B	105	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NBG	A	861	-	15,15,15	1.44	4 (26%)	21,21,21	1.23	3 (14%)
5	URC	B	1863	-	8,13,13	2.31	4 (50%)	5,19,19	8.84	5 (100%)
6	MRD	A	905	-	7,7,7	0.74	0	9,10,10	0.90	0
5	URC	A	863	-	8,13,13	2.22	3 (37%)	5,19,19	8.87	5 (100%)
4	700	A	862	-	29,33,33	1.77	8 (27%)	37,47,47	1.82	10 (27%)
3	PLP	B	1860	1	15,15,16	1.97	4 (26%)	20,22,23	1.13	1 (5%)
6	MRD	A	1902	-	7,7,7	0.50	0	9,10,10	0.70	0
6	MRD	B	902	-	7,7,7	0.66	0	9,10,10	0.64	0
6	MRD	B	903	-	7,7,7	1.30	0	9,10,10	1.01	1 (11%)
6	MRD	A	904	-	7,7,7	0.93	0	9,10,10	0.87	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	700	B	1862	-	29,33,33	1.72	8 (27%)	37,47,47	1.74	8 (21%)
2	NBG	B	1861	-	15,15,15	1.67	4 (26%)	21,21,21	1.39	3 (14%)
3	PLP	A	860	1	15,15,16	2.26	8 (53%)	20,22,23	1.22	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NBG	A	861	-	-	0/6/26/26	0/1/1/1
5	URC	B	1863	-	-	-	0/2/2/2
6	MRD	A	905	-	-	2/5/5/5	-
5	URC	A	863	-	-	-	0/2/2/2
4	700	A	862	-	-	0/19/32/32	0/4/4/4
3	PLP	B	1860	1	-	2/6/6/8	0/1/1/1
6	MRD	A	1902	-	-	2/5/5/5	-
6	MRD	B	902	-	-	2/5/5/5	-
6	MRD	B	903	-	-	0/5/5/5	-
6	MRD	A	904	-	-	3/5/5/5	-
4	700	B	1862	-	-	2/19/32/32	0/4/4/4
2	NBG	B	1861	-	-	0/6/26/26	0/1/1/1
3	PLP	A	860	1	-	0/6/6/8	0/1/1/1

The worst 5 of 43 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1860	PLP	C4A-C4	-5.74	1.39	1.51
5	B	1863	URC	C5-C6	5.06	1.50	1.41
5	A	863	URC	C5-C6	4.56	1.49	1.41
3	A	860	PLP	C4A-C4	-4.23	1.42	1.51
4	B	1862	700	C5-C4	3.74	1.43	1.36

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	863	URC	C4-N9-C8	16.62	111.21	101.12
5	B	1863	URC	C4-N9-C8	16.41	111.08	101.12
5	B	1863	URC	C2-N1-C6	8.92	122.67	115.14
5	A	863	URC	C2-N1-C6	8.69	122.47	115.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
5	A	863	URC	C5-C6-N1	-5.43	116.00	123.43

There are no chirality outliers.

5 of 13 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
6	A	904	MRD	O2-C2-C3-C4
6	A	904	MRD	CM-C2-C3-C4
6	A	904	MRD	C2-C3-C4-C5
6	B	902	MRD	C2-C3-C4-O4
3	B	1860	PLP	C6-C5-C5A-O4P

There are no ring outliers.

6 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	905	MRD	2	0
3	B	1860	PLP	1	0
6	B	902	MRD	3	0
6	B	903	MRD	8	0
6	A	904	MRD	1	0
2	B	1861	NBG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.