



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 12, 2023 – 06:04 AM EDT

PDB ID : 4L7C
Title : Structure of keap1 kelch domain with 2-[[[(1S)-2-[[[(1R,2S)-2-(1H-tetrazol-5-yl)cyclohexyl]carbonyl]-1,2,3,4-tetrahydroisoquinolin-1-yl]methyl]-1H-isoindole-1,3(2H)-dione
Authors : Jnoff, E.; Brookfield, F.; Albrecht, C.; Barker, J.J.; Barker, O.; Beaumont, E.; Bromidge, S.; Brooks, M.; Ceska, T.; Courade, J.P.; Crabbe, T.; Duclos, S.; Fryatt, T.; Jigorel, E.; Kwong, J.; Sands, Z.; Smith, M.A.
Deposited on : 2013-06-13
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	
1	B	300	
1	C	300	

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 7023 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kelch-like ECH-associated protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	286	2250	1396	407	432	15	0	5	0
1	B	286	2221	1380	400	426	15	0	2	0
1	C	286	2228	1384	401	428	15	0	3	0

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	318	MET	-	expression tag	UNP Q14145
A	319	GLY	-	expression tag	UNP Q14145
A	320	HIS	-	expression tag	UNP Q14145
A	354	ASP	ARG	engineered mutation	UNP Q14145
A	610	LEU	-	expression tag	UNP Q14145
A	611	GLU	-	expression tag	UNP Q14145
A	612	HIS	-	expression tag	UNP Q14145
A	613	HIS	-	expression tag	UNP Q14145
A	614	HIS	-	expression tag	UNP Q14145
A	615	HIS	-	expression tag	UNP Q14145
A	616	HIS	-	expression tag	UNP Q14145
A	617	HIS	-	expression tag	UNP Q14145
B	318	MET	-	expression tag	UNP Q14145
B	319	GLY	-	expression tag	UNP Q14145
B	320	HIS	-	expression tag	UNP Q14145
B	354	ASP	ARG	engineered mutation	UNP Q14145
B	610	LEU	-	expression tag	UNP Q14145
B	611	GLU	-	expression tag	UNP Q14145
B	612	HIS	-	expression tag	UNP Q14145
B	613	HIS	-	expression tag	UNP Q14145
B	614	HIS	-	expression tag	UNP Q14145
B	615	HIS	-	expression tag	UNP Q14145
B	616	HIS	-	expression tag	UNP Q14145

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	617	HIS	-	expression tag	UNP Q14145
C	318	MET	-	expression tag	UNP Q14145
C	319	GLY	-	expression tag	UNP Q14145
C	320	HIS	-	expression tag	UNP Q14145
C	354	ASP	ARG	engineered mutation	UNP Q14145
C	610	LEU	-	expression tag	UNP Q14145
C	611	GLU	-	expression tag	UNP Q14145
C	612	HIS	-	expression tag	UNP Q14145
C	613	HIS	-	expression tag	UNP Q14145
C	614	HIS	-	expression tag	UNP Q14145
C	615	HIS	-	expression tag	UNP Q14145
C	616	HIS	-	expression tag	UNP Q14145
C	617	HIS	-	expression tag	UNP Q14145

- Molecule 2 is 2-[[[(1S)-2-[[[(1R,2S)-2-(1H-tetrazol-5-yl)cyclohexyl]carbonyl]-1,2,3,4-tetrahydroisoquinolin-1-yl]methyl]-1H-isoindole-1,3(2H)-dione (three-letter code: 1VW) (formula: C₂₆H₂₆N₆O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
2	A	1	Total	C	N	O	0	0
			35	26	6	3		
2	B	1	Total	C	N	O	0	0
			35	26	6	3		
2	C	1	Total	C	N	O	0	0
			35	26	6	3		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	82	Total O 82 82	0	0
4	B	30	Total O 30 30	0	0
4	C	87	Total O 87 87	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Kelch-like ECH-associated protein 1



- Molecule 1: Kelch-like ECH-associated protein 1



- Molecule 1: Kelch-like ECH-associated protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, α , β , γ	76.29Å 125.61Å 132.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.66 – 2.40 36.66 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (36.66-2.40) 100.0 (36.66-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.74 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.265 , 0.281 0.263 , 0.281	Depositor DCC
R_{free} test set	2573 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	40.8	Xtrriage
Anisotropy	0.155	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 19.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	7023	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 1VW, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/2303	0.47	0/3135
1	B	0.25	0/2274	0.47	0/3097
1	C	0.24	0/2284	0.46	0/3110
All	All	0.25	0/6861	0.47	0/9342

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2250	0	2130	13	0
1	B	2221	0	2107	8	0
1	C	2228	0	2111	6	0
2	A	35	0	26	0	0
2	B	35	0	26	0	0
2	C	35	0	26	0	0
3	A	4	0	3	0	0
3	B	8	0	6	0	0
3	C	8	0	6	0	0
4	A	82	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	30	0	0	0	0
4	C	87	0	0	0	0
All	All	7023	0	6441	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:470[B]:ARG:HH11	1:A:470[B]:ARG:HG3	0.96	1.12
1:A:470[B]:ARG:HH11	1:A:470[B]:ARG:CG	1.79	0.95
1:A:470[B]:ARG:HG3	1:A:470[B]:ARG:NH1	1.78	0.93
1:A:359[B]:GLN:HA	1:A:359[B]:GLN:OE1	1.77	0.82
1:C:466:ALA:HB1	1:C:514:VAL:HG23	1.86	0.56
1:B:466:ALA:HB1	1:B:514:VAL:HG23	1.89	0.54
1:A:366:ALA:HB1	1:A:418:VAL:HG22	1.89	0.53
1:A:387:ASN:HD22	1:A:387:ASN:N	2.07	0.53
1:A:470[B]:ARG:CG	1:A:470[B]:ARG:NH1	2.49	0.52
1:B:575:HIS:CD2	1:B:576:THR:HG23	2.44	0.52
1:A:446[B]:GLU:OE1	1:A:446[B]:GLU:N	2.26	0.51
1:B:527:GLY:HA3	1:C:481:THR:HG21	1.96	0.48
1:B:515:LEU:HD22	1:B:566:ILE:HG13	1.97	0.46
1:A:359[B]:GLN:OE1	1:A:359[B]:GLN:CA	2.59	0.46
1:A:454:ALA:HA	1:B:575:HIS:CE1	2.51	0.46
1:B:483:ARG:HD3	1:B:527:GLY:HA2	1.98	0.46
1:C:366:ALA:HB1	1:C:418:VAL:HG22	2.00	0.44
1:A:610:LEU:O	1:A:611:GLU:HB2	2.18	0.43
1:C:575:HIS:ND1	1:C:576:THR:HG23	2.34	0.43
1:A:560:THR:OG1	1:A:608:VAL:HG23	2.19	0.42
1:C:515:LEU:HD22	1:C:566:ILE:HG13	2.01	0.42
1:C:416:ILE:HD11	1:C:427:ALA:HB1	2.01	0.41
1:A:416:ILE:HD11	1:A:427:ALA:HB1	2.02	0.41
1:B:399:MET:SD	1:B:399:MET:N	2.91	0.41
1:B:366:ALA:HB1	1:B:418:VAL:HG22	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/300 (96%)	280 (97%)	9 (3%)	0	100	100
1	B	286/300 (95%)	272 (95%)	13 (4%)	1 (0%)	41	55
1	C	287/300 (96%)	279 (97%)	7 (2%)	1 (0%)	41	55
All	All	862/900 (96%)	831 (96%)	29 (3%)	2 (0%)	47	62

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	575	HIS
1	C	575	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/244 (98%)	227 (95%)	11 (5%)	27	43
1	B	235/244 (96%)	227 (97%)	8 (3%)	37	56
1	C	236/244 (97%)	230 (98%)	6 (2%)	47	67
All	All	709/732 (97%)	684 (96%)	25 (4%)	38	55

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	415	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	434	CYS
1	A	447	ARG
1	A	498[A]	ARG
1	A	498[B]	ARG
1	A	514	VAL
1	A	543	THR
1	A	557	LEU
1	A	575	HIS
1	A	594	VAL
1	A	610	LEU
1	B	326	ARG
1	B	399	MET
1	B	483	ARG
1	B	543[A]	THR
1	B	543[B]	THR
1	B	557	LEU
1	B	609	THR
1	B	610	LEU
1	C	434	CYS
1	C	447	ARG
1	C	557	LEU
1	C	596	ARG
1	C	609	THR
1	C	610	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	387	ASN
1	A	528	GLN
1	B	359	GLN
1	B	414	ASN
1	B	575	HIS
1	C	414	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	1VW	A	701	-	36,40,40	1.14	3 (8%)	53,58,58	2.32	12 (22%)
3	ACT	A	702	-	3,3,3	0.76	0	3,3,3	0.72	0
3	ACT	B	703	-	3,3,3	0.77	0	3,3,3	0.72	0
3	ACT	C	703	-	3,3,3	0.76	0	3,3,3	0.74	0
3	ACT	C	702	-	3,3,3	0.77	0	3,3,3	0.72	0
2	1VW	B	701	-	36,40,40	1.13	2 (5%)	53,58,58	2.33	13 (24%)
3	ACT	B	702	-	3,3,3	0.77	0	3,3,3	0.76	0
2	1VW	C	701	-	36,40,40	1.15	1 (2%)	53,58,58	2.32	12 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1VW	A	701	-	-	0/12/56/56	0/6/6/6
2	1VW	C	701	-	-	0/12/56/56	0/6/6/6
2	1VW	B	701	-	-	0/12/56/56	0/6/6/6

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	701	1VW	C2-N16	2.39	1.40	1.34
2	B	701	1VW	C2-N16	2.38	1.39	1.34
2	A	701	1VW	C2-N16	2.31	1.39	1.34
2	A	701	1VW	C25-N16	-2.11	1.45	1.47
2	A	701	1VW	C36-C37	-2.01	1.45	1.48
2	B	701	1VW	C36-C37	-2.00	1.45	1.48

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	701	1VW	N15-N14-N13	-7.48	104.65	109.53
2	B	701	1VW	N15-N14-N13	-7.45	104.66	109.53
2	A	701	1VW	O30-C29-N28	7.39	132.01	124.81
2	A	701	1VW	N15-N14-N13	-7.30	104.76	109.53
2	B	701	1VW	O30-C29-N28	7.24	131.86	124.81
2	C	701	1VW	O30-C29-N28	7.18	131.81	124.81
2	C	701	1VW	O38-C37-N28	7.05	131.67	124.81
2	A	701	1VW	O38-C37-N28	6.92	131.55	124.81
2	B	701	1VW	O38-C37-N28	6.92	131.55	124.81
2	B	701	1VW	C11-N15-N14	5.84	110.23	104.33
2	A	701	1VW	C11-N15-N14	5.81	110.19	104.33
2	C	701	1VW	C11-N15-N14	5.76	110.15	104.33
2	B	701	1VW	C9-C11-N12	3.20	127.80	121.25
2	A	701	1VW	C9-C11-N12	3.18	127.75	121.25
2	C	701	1VW	C9-C11-N12	3.11	127.61	121.25
2	A	701	1VW	C31-C29-N28	-3.08	103.69	105.88
2	C	701	1VW	C31-C29-N28	-3.04	103.72	105.88
2	B	701	1VW	C31-C29-N28	-2.97	103.77	105.88
2	A	701	1VW	C36-C37-N28	-2.90	103.82	105.88
2	C	701	1VW	N12-N13-N14	2.87	111.41	109.53
2	C	701	1VW	C36-C37-N28	-2.84	103.86	105.88
2	B	701	1VW	N12-N13-N14	2.79	111.36	109.53
2	B	701	1VW	C36-C37-N28	-2.69	103.97	105.88
2	A	701	1VW	C37-N28-C29	2.55	114.10	112.03
2	C	701	1VW	C37-N28-C29	2.53	114.08	112.03
2	A	701	1VW	N12-N13-N14	2.52	111.19	109.53
2	B	701	1VW	C37-N28-C29	2.37	113.96	112.03
2	A	701	1VW	O30-C29-C31	-2.25	124.30	128.68
2	C	701	1VW	C8-C9-C3	2.22	113.44	109.84
2	B	701	1VW	O30-C29-C31	-2.21	124.37	128.68
2	B	701	1VW	C9-C3-C2	2.21	114.47	111.19
2	B	701	1VW	C8-C9-C3	2.19	113.39	109.84
2	C	701	1VW	O38-C37-C36	-2.17	124.46	128.68

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	701	1VW	O30-C29-C31	-2.16	124.47	128.68
2	B	701	1VW	O38-C37-C36	-2.16	124.48	128.68
2	A	701	1VW	O38-C37-C36	-2.08	124.63	128.68
2	A	701	1VW	C8-C9-C3	2.06	113.18	109.84


There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	286/300 (95%)	0.10	10 (3%) 44 43	28, 38, 50, 58	0
1	B	286/300 (95%)	0.78	45 (15%) 2 1	33, 56, 77, 86	0
1	C	286/300 (95%)	-0.01	6 (2%) 63 61	27, 35, 46, 53	0
All	All	858/900 (95%)	0.29	61 (7%) 16 14	27, 40, 68, 86	0

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	611	GLU	4.7
1	B	353	LEU	4.3
1	B	402	GLN	4.1
1	A	611	GLU	4.0
1	B	326	ARG	4.0
1	B	399	MET	3.8
1	B	397	ASN	3.8
1	B	400	THR	3.8
1	A	326	ARG	3.7
1	A	385	ASP	3.6
1	A	575	HIS	3.6
1	B	372	GLY	3.5
1	B	539	VAL	3.3
1	B	470	ARG	3.2
1	B	540	GLU	3.2
1	B	356	ALA	3.1
1	C	326	ARG	3.1
1	B	348	SER	3.1
1	B	336	ARG	3.1
1	B	493	GLU	3.0
1	B	563	GLN	3.0
1	A	386	GLY	3.0
1	B	329	TYR	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	516	HIS	2.9
1	B	385	ASP	2.9
1	C	385	ASP	2.9
1	B	575	HIS	2.8
1	B	384	PRO	2.8
1	B	398	PRO	2.7
1	C	336	ARG	2.7
1	B	446	GLU	2.6
1	B	371	GLY	2.6
1	B	567	TYR	2.6
1	B	553	ARG	2.6
1	A	384	PRO	2.5
1	C	610	LEU	2.5
1	B	606	VAL	2.5
1	B	517	ASN	2.5
1	B	419	GLY	2.5
1	B	395	CYS	2.5
1	B	541	THR	2.5
1	A	447	ARG	2.4
1	B	340	SER	2.4
1	B	368	CYS	2.3
1	B	610	LEU	2.3
1	B	354	ASP	2.3
1	B	607	ALA	2.2
1	C	384	PRO	2.2
1	B	587	ASP	2.2
1	C	611	GLU	2.2
1	A	336	ARG	2.2
1	B	403	TRP	2.2
1	B	420	VAL	2.1
1	A	528	GLN	2.1
1	B	370	VAL	2.1
1	B	375	TYR	2.1
1	B	374	LEU	2.0
1	A	596	ARG	2.0
1	B	491	TYR	2.0
1	B	366	ALA	2.0
1	B	609	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ACT	C	702	4/4	0.88	0.19	40,40,40,40	0
2	1VW	B	701	35/35	0.92	0.13	45,46,46,46	0
3	ACT	A	702	4/4	0.92	0.34	53,53,53,53	0
2	1VW	A	701	35/35	0.92	0.14	39,39,40,40	0
3	ACT	B	702	4/4	0.93	0.17	44,44,44,44	0
2	1VW	C	701	35/35	0.93	0.13	32,33,33,33	0
3	ACT	C	703	4/4	0.93	0.20	35,36,36,36	0
3	ACT	B	703	4/4	0.94	0.17	37,38,38,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 1VW B 701:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 1VW A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.