



wwPDB X-ray Structure Validation Summary Report ⓘ

May 16, 2020 – 04:55 pm BST

PDB ID : 1LIL
Title : BENICE JONES PROTEIN CLE, A LAMBDA III IMMUNOGLOBULIN
LIGHT-CHAIN DIMER
Authors : Schiffer, M.; Huang, D.B.
Deposited on : 1996-05-13
Resolution : 2.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	212	 32% 48% 17% .
1	B	212	 25% 47% 25% .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3314 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LAMBDA III BENICE JONES PROTEIN CLE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	212	1595	994	267	328	6	0	0	0
1	B	212	1595	994	267	328	6	0	0	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	4	VAL	LEU	CONFLICT	UNP P01842
A	11	LEU	VAL	CONFLICT	UNP P01842
A	20	ARG	SER	CONFLICT	UNP P01842
A	26	GLU	ASP	CONFLICT	UNP P01842
A	27	LYS	THR	CONFLICT	UNP P01842
A	31	ALA	LYS	CONFLICT	UNP P01842
A	33	VAL	ALA	CONFLICT	UNP P01842
A	39	ARG	LYS	CONFLICT	UNP P01842
A	42	GLN	HIS	CONFLICT	UNP P01842
A	46	VAL	LEU	CONFLICT	UNP P01842
A	49	TYR	PHE	CONFLICT	UNP P01842
A	52	ASN	SER	CONFLICT	UNP P01842
A	53	ARG	LYS	CONFLICT	UNP P01842
A	66	SER	ASN	CONFLICT	UNP P01842
A	80	THR	ALA	CONFLICT	UNP P01842
A	81	LEU	MET	CONFLICT	UNP P01842
A	90	VAL	ALA	CONFLICT	UNP P01842
A	94	ASN	-	INSERTION	UNP P01842
A	95	ALA	-	INSERTION	UNP P01842
A	?	-	THR	DELETION	UNP P01842
A	96	VAL	ALA	CONFLICT	UNP P01842
B	4	VAL	LEU	CONFLICT	UNP P01842
B	11	LEU	VAL	CONFLICT	UNP P01842
B	20	ARG	SER	CONFLICT	UNP P01842
B	26	GLU	ASP	CONFLICT	UNP P01842

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	27	LYS	THR	CONFLICT	UNP P01842
B	31	ALA	LYS	CONFLICT	UNP P01842
B	33	VAL	ALA	CONFLICT	UNP P01842
B	39	ARG	LYS	CONFLICT	UNP P01842
B	42	GLN	HIS	CONFLICT	UNP P01842
B	46	VAL	LEU	CONFLICT	UNP P01842
B	49	TYR	PHE	CONFLICT	UNP P01842
B	52	ASN	SER	CONFLICT	UNP P01842
B	53	ARG	LYS	CONFLICT	UNP P01842
B	66	SER	ASN	CONFLICT	UNP P01842
B	80	THR	ALA	CONFLICT	UNP P01842
B	81	LEU	MET	CONFLICT	UNP P01842
B	90	VAL	ALA	CONFLICT	UNP P01842
B	94	ASN	-	INSERTION	UNP P01842
B	95	ALA	-	INSERTION	UNP P01842
B	?	-	THR	DELETION	UNP P01842
B	96	VAL	ALA	CONFLICT	UNP P01842


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	60	Total O 60 60	0	0
2	B	64	Total O 64 64	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: LAMBDA III BENGE JONES PROTEIN CLE



- Molecule 1: LAMBDA III BENGE JONES PROTEIN CLE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	113.17Å 72.54Å 49.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.65 9.42 – 2.66	Depositor EDS
% Data completeness (in resolution range)	71.0 (10.00-2.65) 73.0 (9.42-2.66)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtrriage
Refinement program	PROLSQ, X-PLOR	Depositor
R, R_{free}	0.190 , 0.340 0.178 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	34.1	Xtrriage
Anisotropy	0.575	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.24 , 135.6	EDS
L-test for twinning ¹	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3314	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.98	1/1634 (0.1%)	1.78	27/2234 (1.2%)
1	B	0.98	0/1634	1.87	36/2234 (1.6%)
All	All	0.98	1/3268 (0.0%)	1.83	63/4468 (1.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	38	GLN	CB-CG	7.52	1.72	1.52

The worst 5 of 63 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	54	ARG	NE-CZ-NH1	-12.58	114.01	120.30
1	B	39	ARG	NE-CZ-NH2	11.35	125.97	120.30
1	A	190	ARG	NE-CZ-NH1	-10.25	115.18	120.30
1	A	53	ARG	NE-CZ-NH1	10.15	125.38	120.30
1	B	53	ARG	NE-CZ-NH2	9.48	125.04	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1595	0	1533	144	0
1	B	1595	0	1533	179	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	60	0	0	9	2
2	B	64	0	0	12	1
All	All	3314	0	3066	319	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 51.

The worst 5 of 319 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:36:TYR:CE1	1:B:46:VAL:HG22	1.69	1.26
1:B:199:GLU:HA	2:B:409:HOH:O	1.35	1.25
1:A:47:VAL:O	1:A:48:ILE:HD13	1.43	1.14
1:A:87:TYR:HB3	1:A:98:PHE:CE1	1.83	1.13
1:A:87:TYR:HB3	1:A:98:PHE:HE1	1.00	1.07

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:375:HOH:O	2:B:479:HOH:O 2_655]	2.12	0.08
1:B:215:SER:O	2:A:310:HOH:O 4_456]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	210/212 (99%)	161 (77%)	33 (16%)	16 (8%)	1	0
1	B	210/212 (99%)	166 (79%)	27 (13%)	17 (8%)	1	0
All	All	420/424 (99%)	327 (78%)	60 (14%)	33 (8%)	1	0

5 of 33 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	28	LEU
1	A	30	ASP
1	A	40	PRO
1	B	40	PRO
1	B	69	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	181/181 (100%)	139 (77%)	42 (23%)	1 0
1	B	181/181 (100%)	139 (77%)	42 (23%)	1 0
All	All	362/362 (100%)	278 (77%)	84 (23%)	1 0

5 of 84 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	193	SER
1	B	30	ASP
1	B	187	LYS
1	A	208	THR
1	B	5	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	89	GLN
1	A	94	ASN
1	B	89	GLN
1	A	79	GLN
1	B	52	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	212/212 (100%)	-0.76	0 100 100	16, 16, 16, 16	0
1	B	212/212 (100%)	-0.76	0 100 100	16, 16, 16, 16	0
All	All	424/424 (100%)	-0.76	0 100 100	16, 16, 16, 16	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.